

DECOMPOSITION METHOD FOR THE *B*-BALANCED SHALLOW WATER EQUATION

Zhijun Qiao & Jules Kamdem Sadefo

ABSTRACT

The Adomian decomposition approach is used for the *b*-balanced shallow water wave equation $m_t + m_x u + b m u_x = 0$, $m = u - u_{xx}$. Approximate solutions are obtained for three smooth initial values. Compared with the existing method, our procedure just works with the polynomial and algebraic computations for this shallow water equation.

Key Words: Shallow water equation, Adomian decomposition method, Approximate solutions.

1. INTRODUCTION

Recently, the balanced water wave equation

$$m_t + m_x u + b m u_x = 0, m = u - u_{xx} \quad (1.1)$$

has arisen a lot of attractive attentions [5]. The family (1.1) is integrable only when $b = 2, 3$ [7], and it is reduced to the Camassa-Holm (*CH*) equation [2] as $b = 2$ and to the Degasperis-Procesi (*DP*) equation [3] as $b = 3$, respectively. The *DP* equation has bi-Hamiltonian structure and Lax pair [4], and can be also extended to a whole integrable hierarchy of equations [8] with parametric solutions under some constraints [9]. In addition, the *CH* equation also has new peaked and smooth solitons which are given in explicit forms [11].

In an earlier paper [6], we dealt with the decomposition of the *CH* equation using the Adomian method [1]. Very recently, Qiao [10] developed three new types of soliton solutions-M-shape peakon, dehisced soliton, and double dehisced 1-peak solions through studying the *DP* equation, and the most interesting is that a cuspon is a limit of the new peaked soliton solutions for the *DP* equation (see [10]). In the present paper, we discuss the *b*-balanced water wave equation (1.1) Using analysis procedure and the Adomian decomposition method, we are able to obtain the approximate solutions for three smooth initial values. Compared with the existing method, our procedure just works with the polynomial and algebraic computations for this family of water equations. Finally, we plot our approximate solutions in three dimensional space.

1.1. Analysis of the decomposition method. After an expansion, the b -balance water wave equation (1.1) for real $u(x, t)$ is rewritten as

$$L_t(u - c u_{xx}) = -m_x u - b m u_x, \quad (1.2)$$

where $L_t = \frac{\partial}{\partial t}$ is a linear operator.

Assuming the inverse operator L_t^{-1} exists and it can be taken as the definite integral with respect to t from t_0 to t , ie.

$$L_t^{-1} = \int_{t_0}^t (\cdot) dt$$

then applying the inverse operator L_t^{-1} on both sides of (1.2) yields

$$m(x, t) = m(x, t_0) + L_t^{-1} (-m_x u - b m u_x). \quad (1.3)$$

The ADM assumes that the unknown function $u(x, t)$ can be expressed by a sum of components defined by the decomposition series of the form:

$$u(x, t) = \sum_{n=0}^{\infty} u_n(x, t)$$

with u_0 defined as $u(x, 0)$ where $u(x, t)$ will be determined recursively. The nonlinear operator

$$\begin{aligned} NL(u) &:= m_x u + b m u_x \\ &:= m[u_x]u + b m[u]u_x \end{aligned} \quad (1.4)$$

can also be decomposed by an infinite series of polynomials given by

$$NL(u) = \sum_{n=0}^{\infty} A_n(u_0, u_1, \dots, u_n),$$

where $A_n(u_0, u_1, \dots, u_n)$ are the Adomian's polynomials which are defined as

$$A_n = \frac{1}{n!} \left[\frac{d}{d\lambda^n} NL \left(\sum_{i=0}^{\infty} \lambda^i u_i(x, t) \right) \right]_{\lambda=0}, \quad n = 0, 1, 2, \dots$$

$$= \frac{1}{n!}$$

since $m[\cdot]$ is

As we s
polynomial:

Algorithm

- Input da
 t_0 - initi
 $u(x, t_0)$ -
 c - posi
 b - con:
 k - num:
• Output:
- Step :
- Step :
- Step :

- Step :

$$= \frac{1}{n!} \left[\frac{d}{d\lambda^n} \left[m \left[\sum_{n=0}^{\infty} \lambda^j u_{jx} \right] \left(\sum_{n=0}^{\infty} \lambda^j u_j \right) + bm \left[\sum_{n=0}^{\infty} \lambda^j u_j \right] \left(\sum_{n=0}^{\infty} \lambda^j u_{jx} \right) \right] \right]_{\lambda=0}$$

$$\sum_{j=0}^n [m[u_{jx}] u_{n-j} + bm [u_{n-j}] u_{jx}], \tag{1.5}$$

since $m[\cdot]$ is a linear operator. Recursively, we then generates the formula of u_n .

$$\begin{cases} u_0 = u(x, t_0) = cu_{xx}(x, t_0) & b = 0 \\ u_{n+1} = cu_{nxx} - \int_{t_0}^t A_n ds & \text{if } n \neq 0. \end{cases} \tag{1.6}$$

As we see, it is easy to write a computer code for generating the Adomian polynomials. We summarize the whole procedure in the following algorithm

Algorithm

- Input data:

t_0 – initial time

$u(x, t_0)$ – initial condition,

c – positive constant parameter, i.e, $m[u](x, t_0) = u(x, t_0) - cu_{xx}(x, t_0) = J(x)$,

b – constant parameter of (1.1),

k – number of terms in the approximation.

- Output: $u_{approx}(x, t)$. The approximate solution

– Step 1: Set $u_0 = J(x)$ and $u_{approx}(x, t) = u_0$.

– Step 2: For $k = 0$ to $n - 1$, do Step 3, Step 4, and Step 5.

– Step 3: Compute

$$A_k = \sum_{j=0}^k [m[u_{jx}] u_{k-j} + bm [u_{k-j}] u_{jx}]$$

– Step 4: Compute

$$u_{k+1} = cu_{kxx} - \int_{t_0}^t A_k ds \quad \text{if } k \neq 0.$$

- Step 5: Compute $u_{approx} = u_{approx} + u_{k+1}$.
- Stop

2. CONVERGENCE ANALYSIS

In this section, we discuss the convergence property of the approximated solution for the b -balanced wave equation.

Let us consider the b -balanced wave equation in the Hilbert space $H = L^2((\alpha, \beta) \times [0, T])$.

$$H = \left\{ v : (\alpha, \beta) \times [0, T] \text{ with } \int_{(\alpha, \beta) \times [0, T]} v^2(x, s) ds dt < +\infty \right\}. \quad (2.7)$$

Then the operator is of the form

$$\begin{aligned} T(u) &= L_t(u - c u_{xx}) \\ &= -(1 + b) u_x u + u_{xxx} u + b u_{xx} u_x. \end{aligned} \quad (2.8)$$

The Adomian decomposition method is convergent if the following two hypotheses are satisfied¹ and some references therein for more details.

- (Hyp1): There exists a constant $k > 0$ such that the following inner product holds in H :

$$(T(u) - T(v), u - v) \geq k \|u - v\|, \quad \forall u, v \in H; \quad (2.9)$$

- (Hyp2): As long as both $u \in H$ and $v \in H$ are bounded (i.e. there is a positive number M such that $\|u\| \leq M$, $\|v\| \leq M$), there exists a constant $\theta(M) > 0$ such that

$$(T(u) - T(v), u - v) \leq \theta(M) \|u - v\| \|w\|, \quad \forall w \in H. \quad (2.10)$$

Theorem 2.1. (Sufficient conditions of convergence for the b -balanced wave equation) Let

$$\begin{aligned} T(u) &= L_t(u - c u_{xx}) \\ &= -(1 + b) u_x u + u_{xxx} u + b u_{xx} u_x, \\ &\text{with } b \leq 3, \quad L_t = \frac{\partial}{\partial t}, \end{aligned}$$

and consider the free initial and boundary conditions for the b -balanced equation. Then the Adomian decomposition method leads to a special solution of the b -balanced wave equation.

Proof. To prove the theorem, we just verify the conditions (Hyp1) and (Hyp2). For $\forall u, v \in H$, let us calculate:

$$\begin{aligned} T(u) - T(v) &= -(1+b)(u_x u - v_x v) + (u_{xxx} u - v_{xxx} v) + b(u_{xx} u_x - v_{xx} v_x) \\ &= \frac{1+b}{2}(u^2 - v^2)_x + \frac{1}{2}((u^2 - v^2)_{xxx} - 3(u_x^2 - v_x^2)_x) + \frac{b}{2}(u_x^2 - v_x^2)_x \\ &= \frac{1+b}{2}(u^2 - v^2)_x + \frac{1}{2}(u^2 - v^2)_{xxx} + \frac{b-3}{2}(u_x^2 - v_x^2)_x \end{aligned}$$

Therefore, we have the inner product

$$\begin{aligned} (T(u) - T(v), u - v) &= \frac{1+b}{2} \left(-\frac{\partial}{\partial x} (u^2 - v^2), u - v \right) - \frac{1}{2} \left(-\frac{\partial}{\partial x} (u_x^2 - v_x^2), u - v \right) \\ &\quad + \frac{3-b}{2} \left(-\frac{\partial^3}{\partial x^3} (u^2 - v^2), u - v \right). \end{aligned} \quad (2.11)$$

Let us assume that u, v are bounded and there is a constant $M > 0$ such that $(u, u), (v, v) < M^2$. By using Schwartz inequality

$$\left(-\frac{\partial}{\partial x} (u^2 - v^2), u - v \right) \leq \| (u^2 - v^2)_x \| \| u - v \|, \quad (2.12)$$

and since there exist θ_1 and θ_2 such that $\| (u - v)_x \| \leq \theta_1 \| u - v \|$, $\| (u + v)_x \| \leq \theta_2 \| u - v \|$ and $\| u + v \| \leq 2M$, we have

$$\left(\frac{\partial}{\partial x} (u^2 - v^2), u - v \right) \leq 2M\theta_1\theta_2 \| u - v \|^2.$$

\Leftrightarrow

$$\left(-\frac{\partial}{\partial x} (u^2 - v^2), u - v \right) \geq 2M\theta_1\theta_2 \| u - v \|^2. \quad (2.13)$$

Following the preceding procedure, we can calculate.

$$\left(\frac{\partial}{\partial x} (u_x^2 - v_x^2), u - v \right) \leq \| (u_x^2 - v_x^2)_x \| \| u - v \|^2$$

$$\begin{aligned} &\leq \theta_3 \|u_x + v_x\| \|u_x - v_x\| \|u - v\| \\ &\leq 2M\theta_3\theta_4\theta_5 \|u - v\|^2, \end{aligned}$$

⇔

$$\left(-\frac{\partial}{\partial x}(u_x^2 - v_x^2), u - v \right) \geq 2M\theta_3\theta_4\theta_5 \|u - v\|^2, \quad (2.14)$$

where θ_i ($i = 3, 4, 5$) are positive constants.

Moreover, the Cauchy-Schwartz-Buniakowski inequality yields

$$\left(\frac{\partial^3}{\partial x^3}(u^2 - v^2), u - v \right) \leq \|(u^2 - v^2)_{xxx}\| \|u - v\|, \quad (2.15)$$

then by using the mean value theorem, we have

$$\begin{aligned} \left(\frac{\partial^3}{\partial x^3}(u^2 - v^2), u - v \right) &\leq \theta_6\theta_7\theta_8 \|u^2 - v^2\| \|u - v\| \\ &\leq 2M\theta_6\theta_7\theta_8 \|u - v\|^2 \end{aligned}$$

⇔

$$\left(\frac{\partial^3}{\partial x^3}(u^2 - v^2), u - v \right) \geq 2M\theta_6\theta_7\theta_8 \|u - v\|^2 \quad (2.16)$$

where θ_j ($j = 6, 7, 8$) are three positive constants, and $\|(u^2 - v^2)_{xxx}\| \leq \theta_6\|(u^2 - v^2)_{xx}\|$, $\|(u + v)_{xx}\| \leq \theta_7\|(u + v)_x\|$ and $\|(u + v)_x\| \leq \theta_8\|u + v\|$.

Substituting (2.13), (2.14), (2.16) into (2.11) generates the following inner product:

$$\begin{aligned} (T(u) - T(v), u - v) &= \left(-\frac{\partial}{\partial x}(u^2 - v^2), u - v \right) \\ &\quad - (c - d) \left(\frac{\partial}{\partial x}(u_x^2 - v_x^2), u - v \right) - \frac{d}{2} \left(\frac{\partial^3}{\partial x^3}(u^2 - v^2), u - v \right) \\ &\geq k \|u - v\|^2, \end{aligned}$$

Decompositio

where $k = ($
 b -balanced

Let us r
 compute:

$(T(u) -$

where $\theta(M)$

$1 \cdot 10^{59}$
 $5 \cdot 10^{59}$

$-5 \cdot 10^{59}$

where $k = ((1 + b) \theta_1 \theta_2 - \theta_3 \theta_4 \theta_5 + (3 - b) \theta_6 \theta_7 \theta_8) M$. So, (Hyp1) is true for the b -balanced wave equation.

Let us now verify the hypotheses (Hyp2) for the operator $T(u)$. We directly compute:

$$(T(u) - T(v), w) = \frac{1+b}{2} \left(-b \frac{\partial}{\partial x} [u^2 - v^2], w \right) - \frac{1}{2} \left(-\frac{\partial}{\partial x} [u_x^2 - v_x^2], w \right) \\ - \frac{3-b}{2} \left(-\left[\frac{\partial^3}{\partial x^3} (u^2 - v^2) \right], w \right)$$

$$\leq \theta(M) \|u - v\| \|w\|$$

where $\theta(M) = 3M/2$. Therefore, (Hyp2) is correct as well.




Figure 1: Approximate solution for equation (3.17) with $b = -1$.

3. IMPLEMENTATION OF THE METHOD AND APPROXIMATE SOLUTIONS

In this section, we take some examples to show the procedure and present some approximate solutions $u(x, t) \simeq \sum_{j=0}^{10} u_n(x, t)$, for the b -balanced equation.

Example 3.1.

$$\begin{cases} m_t + m_x u + b m u_x = 0, & m = u - c u_{xx} \\ u_0 = u(x, 0) = -3e^{4x} + 2e^{5x}. \end{cases} \quad (3.17)$$

Example 3.2.

$$\begin{cases} m_t + m_x u + b m u_x = 0, & m = u - c u_{xx} \\ u_0 = u(x, 0) = \cosh(gx^2). \end{cases} \quad (3.18)$$

Example 3.3.

$$\begin{cases} m_t + m_x u + b m u_x = 0, & m = u - c u_{xx} \\ u_0 = u(u, 0) = 3 \cosh(c_1 x) + 2 \sin(c_2 x). \end{cases} \quad (3.19)$$




Figure 2: Approximate solution for equation (3.17) with $b = 0$.

4. CONCI

In this paper:
family of b
The initial

2.5

-2.5

5.1

-5.

-1.

MATE

it some




Figure 3: Approximate solution for equation (3.17) with $b = 0.5$

4. CONCLUSIONS

In this paper, we apply the Adomian polynomial decomposition method to solve the family of b -balanced shallow water equations in an explicitly approximate form. The initial values we adopted are smooth, but more interesting is: Some of




Figure 4: Approximate solution for equation (3.17) with $b = 3$




Figure 5: Approximate solution for equation (3.18) with $b = 1, c = 1, g = 2$.

approximate solutions are weak solutions with some peaks (see figures 1-9). In comparison with the existing method to obtain peaked solitary solutions, our procedure just works on the polynomial and algebraic computations. Our paper presents some peaked (i.e. Continuous but nonsmooth) explicit solutions for the b -balanced equation (1.1).




Figure 6: Approximate solution for equation (3.19) with $b = 0, c_1 = 4, c_2 = 5$.

-2.
-4
-6
-8

Fig

5. ACKNO

Qiaos work
Wafo Soh fo

5.10

-5.1
-1.1

Fig




Figure 7: Approximate solution for equation (3.19) with $b = -1, c_1 = 4, c_2 = 5$.

5. ACKNOWLEDGMENT

Qiaos work was partially supported by the *UTPA-FRC* grant. Sadefo thanks prof. C. Wafo Soh for some references concerning *ADM*.




Figure 8: Approximate solution for equation (3.19) with $b = -2, c_1 = 4, c_2 = 5$.

. In
our
per
e b-




Figure 9: Approximate solution for equation (3.19) with $b = -3$, $c_1 = 4$, $c_2 = 5$.

REFERENCES

- [1] G. Adomian, *Solving Frontier Problems of Physics: The Decomposition method*, Kluwer Academic, Dordrecht/Norwell, MA, 1994.
- [2] R. Camassa and D. D. Holm, An integrable shallow water equation with peaked solitons, *Phys. Rev. Lett.* **71** (1993), 1661-1664.
- [3] A. Degasperis and M. Procesi, Asymptotic integrability, in A. Degasperis and G. Gaeta (eds), *Symmetry and Perturbation Theory*, World Scientific, 1999, pp. 23-37.
- [4] A. Degasperis, D. D. Holm and A. N. W. Hone, A new integrable equation with peakon solutions, *Theoret. and Math. Phys.* **133** (2002), 1463-1474.
- [5] Darryl D. Holm and Martin F. Staley, Nonlinear balance and exchange of stability in dynamics of solitons, peakons, ramps/clis and leftons in a 1+1 nonlinear evolutionary PDE, *Physics Letters A* **308** (2003), 437-444.
- [6] J. Sadefo Kamdem and Zhijun Qiao, Decomposition method for the Camassa-Holm equation, *Chaos, Solitons and Fractals*, to appear in 2006.
- [7] A. V. Mikhailov and V. S. Novikov, Perturbative symmetry approach, *J. Phys. A: Math. Gen.* **35** (2002), 4775-4790.
- [8] Z. J. Qiao, Integrable hierarchy, 3×3 constrained systems, and parametric and stationary solutions, *Acta Applicandae Mathematicae* **83** (2004), 199-220.
- [9] Z. J. Qiao, Generalized r-matrix structure and algebro-geometric solutions for integrable systems, *Rev. Math. Phys.* **13** (2001), 545-586.

Decomposition

- [10] Z. J. Qiao, *Degasperis*
- [11] Z. J. Qiao, *Europhys.*

Zhijun Qiao
 Department of M
 The University of
 1201 W Univers
 E-mail: qiao@u

Jules Kamdem
 Laboratoire de r
 BP 1039 moulin
 E-mail: sadefo@

- [10] Z. J. Qiao, M-shape peakons, dehisced solitons, cuspons and new 1-peak solitons for the Degasperis-Procesi Equation, preprint, 2006.
- [11] Z. J. Qiao and G. Zhang, On peaked and smooth solitons for the Camassa-Holm equation, *Europhys. Lett.* **73** (2006), 657-663.

Zhijun Qiao

Department of Mathematics
The University of Texas-Pan American
1201 W University Drive, Edinburg, TX 78541, USA
E-mail: qiao@utpa.edu

Jules Kamdem Sadefo

Laboratoire de mathématiques, CNRS UMR 6056
BP 1039 moulin de la housse, 51687 Reims France
E-mail: sadefo@univ.reims.fr

Kluwer

solitons,

ta (eds),

peakon

ynamics

Physics

quation,

th. Gen.

tionary

egrable