

# Calculus, once again

---

David A. SANTOS  
dsantos@ccp.edu

November 6, 2007 Version

Copyright © 2007 David Anthony SANTOS. Permission is granted to copy, distribute and/or modify this document under the terms of the GNU Free Documentation License, Version 1.2 or any later version published by the Free Software Foundation; with no Invariant Sections, no Front-Cover Texts, and no Back-Cover Texts. A copy of the license is included in the section entitled “GNU Free Documentation License”.

# GNU Free Documentation License

Version 1.2, November 2002  
Copyright © 2000,2001,2002 Free Software Foundation, Inc.

51 Franklin St, Fifth Floor, Boston, MA 02110-1301 USA

Everyone is permitted to copy and distribute verbatim copies of this license document, but changing it is not allowed.

## Preamble

The purpose of this License is to make a manual, textbook, or other functional and useful document “free” in the sense of freedom: to assure everyone the effective freedom to copy and redistribute it, with or without modifying it, either commercially or noncommercially. Secondly, this License preserves for the author and publisher a way to get credit for their work, while not being considered responsible for modifications made by others.

This License is a kind of “copyleft”, which means that derivative works of the document must themselves be free in the same sense. It complements the GNU General Public License, which is a copyleft license designed for free software.

We have designed this License in order to use it for manuals for free software, because free software needs free documentation: a free program should come with manuals providing the same freedoms that the software does. But this License is not limited to software manuals; it can be used for any textual work, regardless of subject matter or whether it is published as a printed book. We recommend this License principally for works whose purpose is instruction or reference.

## 1. APPLICABILITY AND DEFINITIONS

This License applies to any manual or other work, in any medium, that contains a notice placed by the copyright holder saying it can be distributed under the terms of this License. Such a notice grants a world-wide, royalty-free license, unlimited in duration, to use that work under the conditions stated herein. The “**Document**”, below, refers to any such manual or work. Any member of the public is a licensee, and is addressed as “**you**”. You accept the license if you copy, modify or distribute the work in a way requiring permission under copyright law.

A “**Modified Version**” of the Document means any work containing the Document or a portion of it, either copied verbatim, or with modifications and/or translated into another language.

A “**Secondary Section**” is a named appendix or a front-matter section of the Document that deals exclusively with the relationship of the publishers or authors of the Document to the Document’s overall subject (or to related matters) and contains nothing that could fall directly within that overall subject. (Thus, if the Document is in part a textbook of mathematics, a Secondary Section may not explain any mathematics.) The relationship could be a matter of historical connection with the subject or with related matters, or of legal, commercial, philosophical, ethical or political position regarding them.

The “**Invariant Sections**” are certain Secondary Sections whose titles are designated, as being those of Invariant Sections, in the notice that says that the Document is released under this License. If a section does not fit the above definition of Secondary then it is not allowed to be designated as Invariant. The Document may contain zero Invariant Sections. If the Document does not identify any Invariant Sections then there are none.

The “**Cover Texts**” are certain short passages of text that are listed, as Front-Cover Texts or Back-Cover Texts, in the notice that says that the Document is released under this License. A Front-Cover Text may be at most 5 words, and a Back-Cover Text may be at most 25 words.

A “**Transparent**” copy of the Document means a machine-readable copy, represented in a format whose specification is available to the general public, that is suitable for revising the document straightforwardly with generic text editors or (for images composed of pixels) generic paint programs or (for drawings) some widely available drawing editor, and that is suitable for input to text formatters or for automatic translation to a variety of formats suitable for input to text formatters. A copy made in an otherwise Transparent file format whose markup, or absence of markup, has been arranged to thwart or discourage subsequent modification by readers is not Transparent. An image format is not Transparent if used for any substantial amount of text. A copy that is not “Transparent” is called “**Opaque**”.

Examples of suitable formats for Transparent copies include plain ASCII without markup, Texinfo input format, LaTeX input format, SGML or XML using a publicly available DTD, and standard-conforming simple HTML, PostScript or PDF designed for human modification. Examples of transparent image formats include PNG, XCF and JPG. Opaque formats include proprietary formats that can be read and edited only by proprietary word processors, SGML or XML for which the DTD and/or processing tools are not generally available, and the machine-generated HTML, PostScript or PDF produced by some word processors for output purposes only.

The “**Title Page**” means, for a printed book, the title page itself, plus such following pages as are needed to hold, legibly, the material this License requires to appear in the title page. For works in formats which do not have any title page as such, “Title Page” means the text near the most prominent appearance of the work’s title, preceding the beginning of the body of the text.

A section “**Entitled XYZ**” means a named subunit of the Document whose title either is precisely XYZ or contains XYZ in parentheses following text that translates XYZ in another language. (Here XYZ stands for a specific section name mentioned below, such as “**Acknowledgements**”, “**Dedications**”, “**Endorsements**”, or “**History**”). To “**Preserve the Title**” of such a section when you modify the Document means that it remains a section “Entitled XYZ” according to this definition.

The Document may include Warranty Disclaimers next to the notice which states that this License applies to the Document. These Warranty Disclaimers are considered to be included by reference in this License, but only as regards disclaiming warranties; any other implication that these Warranty Disclaimers may have is void and has no effect on the meaning of this License.

## 2. VERBATIM COPYING

You may copy and distribute the Document in any medium, either commercially or noncommercially, provided that this License, the copyright notices, and the license notice saying this License applies to the Document are reproduced in all copies, and that you add no other conditions whatsoever to those of this License. You may not use technical measures to obstruct or control the reading or further copying of the copies you make or distribute. However, you may accept compensation in exchange for copies. If you distribute a large enough number of copies you must also follow the conditions in section 3.

You may also lend copies, under the same conditions stated above, and you may publicly display copies.

## 3. COPYING IN QUANTITY

If you publish printed copies (or copies in media that commonly have printed covers) of the Document, numbering more than 100, and the Document’s license notice requires Cover Texts, you must enclose the copies in covers that carry, clearly and legibly, all these Cover Texts: Front-Cover Texts on the front cover, and Back-Cover Texts on the back cover. Both covers must also clearly and legibly identify you as the publisher of these copies. The front cover must present the full title with all words of the title equally prominent and visible. You may add other material on the covers in addition. Copying with changes limited to the covers, as long as they preserve the title of the Document and satisfy these conditions, can be treated as verbatim copying in other respects.

If the required texts for either cover are too voluminous to fit legibly, you should put the first ones listed (as many as fit reasonably) on the actual cover, and continue the rest onto adjacent pages.

If you publish or distribute Opaque copies of the Document numbering more than 100, you must either include a machine-readable Transparent copy along with each Opaque copy, or state in or with each Opaque copy a computer-network location from which the general network-using public has access to download using public-standard network protocols a complete Transparent copy of the Document, free of added material. If you use the latter option, you must take reasonably prudent steps, when you begin distribution of Opaque copies in quantity, to ensure that this Transparent copy will remain thus accessible at the stated location until at least one year after the last time you distribute an Opaque copy (directly or through your agents or retailers) of that edition to the public.

It is requested, but not required, that you contact the authors of the Document well before redistributing any large number of copies, to give them a chance to provide you with an updated version of the Document.

## 4. MODIFICATIONS

You may copy and distribute a Modified Version of the Document under the conditions of sections 2 and 3 above, provided that you release the Modified Version under precisely this License, with the Modified Version filling the role of the Document, thus licensing distribution and modification of the Modified Version to whoever possesses a copy of it. In addition, you must do these things in the Modified Version:

- Use in the Title Page (and on the covers, if any) a title distinct from that of the Document, and from those of previous versions (which should, if there were any, be listed in the History section of the Document). You may use the same title as a previous version if the original publisher of that version gives permission.
- List on the Title Page, as authors, one or more persons or entities responsible for authorship of the modifications in the Modified Version, together with at least five of the principal authors of the Document (all of its principal authors, if it has fewer than five), unless they release you from this requirement.
- State on the Title page the name of the publisher of the Modified Version, as the publisher.

- D. Preserve all the copyright notices of the Document.
- E. Add an appropriate copyright notice for your modifications adjacent to the other copyright notices.
- F. Include, immediately after the copyright notices, a license notice giving the public permission to use the Modified Version under the terms of this License, in the form shown in the Addendum below.
- G. Preserve in that license notice the full lists of Invariant Sections and required Cover Texts given in the Document's license notice.
- H. Include an unaltered copy of this License.
- I. Preserve the section Entitled "History", Preserve its Title, and add to it an item stating at least the title, year, new authors, and publisher of the Modified Version as given on the Title Page. If there is no section Entitled "History" in the Document, create one stating the title, year, authors, and publisher of the Document as given on its Title Page, then add an item describing the Modified Version as stated in the previous sentence.
- J. Preserve the network location, if any, given in the Document for public access to a Transparent copy of the Document, and likewise the network locations given in the Document for previous versions it was based on. These may be placed in the "History" section. You may omit a network location for a work that was published at least four years before the Document itself, or if the original publisher of the version it refers to gives permission.
- K. For any section Entitled "Acknowledgements" or "Dedications", Preserve the Title of the section, and preserve in the section all the substance and tone of each of the contributor acknowledgements and/or dedications given therein.
- L. Preserve all the Invariant Sections of the Document, unaltered in their text and in their titles. Section numbers or the equivalent are not considered part of the section titles.
- M. Delete any section Entitled "Endorsements". Such a section may not be included in the Modified Version.
- N. Do not retitle any existing section to be Entitled "Endorsements" or to conflict in title with any Invariant Section.
- O. Preserve any Warranty Disclaimers.

If the Modified Version includes new front-matter sections or appendices that qualify as Secondary Sections and contain no material copied from the Document, you may at your option designate some or all of these sections as invariant. To do this, add their titles to the list of Invariant Sections in the Modified Version's license notice. These titles must be distinct from any other section titles.

You may add a section Entitled "Endorsements", provided it contains nothing but endorsements of your Modified Version by various parties—for example, statements of peer review or that the text has been approved by an organization as the authoritative definition of a standard.

You may add a passage of up to five words as a Front-Cover Text, and a passage of up to 25 words as a Back-Cover Text, to the end of the list of Cover Texts in the Modified Version. Only one passage of Front-Cover Text and one of Back-Cover Text may be added by (or through arrangements made by) any one entity. If the Document already includes a cover text for the same cover, previously added by you or by arrangement made by the same entity you are acting on behalf of, you may not add another; but you may replace the old one, on explicit permission from the previous publisher that added the old one.

The author(s) and publisher(s) of the Document do not by this License give permission to use their names for publicity for or to assert or imply endorsement of any Modified Version.

## 5. COMBINING DOCUMENTS

You may combine the Document with other documents released under this License, under the terms defined in section 4 above for modified versions, provided that you include in the combination all of the Invariant Sections of all of the original documents, unmodified, and list them all as Invariant Sections of your combined work in its license notice, and that you preserve all their Warranty Disclaimers.

The combined work need only contain one copy of this License, and multiple identical Invariant Sections may be replaced with a single copy. If there are multiple Invariant Sections with the same name but different contents, make the title of each such section unique by adding at the end of it, in parentheses, the name of the original author or publisher of that section if known, or else a unique number. Make the same adjustment to the section titles in the list of Invariant Sections in the license notice of the combined work.

In the combination, you must combine any sections Entitled "History" in the various original documents, forming one section Entitled "History"; likewise combine any sections Entitled "Acknowledgements", and any sections Entitled "Dedications". You must delete all sections Entitled "Endorsements".

## 6. COLLECTIONS OF DOCUMENTS

You may make a collection consisting of the Document and other documents released under this License, and replace the individual copies of this License in the various documents with a single copy that is included in the collection, provided that you follow the rules of this License for verbatim copying of each of the documents in all other respects.

You may extract a single document from such a collection, and distribute it individually under this License, provided you insert a copy of this License into the extracted document, and follow this License in all other respects regarding verbatim copying of that document.

## 7. AGGREGATION WITH INDEPENDENT WORKS

A compilation of the Document or its derivatives with other separate and independent documents or works, in or on a volume of a storage or distribution medium, is called an "aggregate" if the copyright resulting from the compilation is not used to limit the legal rights of the compilation's users beyond what the individual works permit. When the Document is included in an aggregate, this License does not apply to the other works in the aggregate which are not themselves derivative works of the Document.

If the Cover Text requirement of section 3 is applicable to these copies of the Document, then if the Document is less than one half of the entire aggregate, the Document's Cover Texts may be placed on covers that bracket the Document within the aggregate, or the electronic equivalent of covers if the Document is in electronic form. Otherwise they must appear on printed covers that bracket the whole aggregate.

## 8. TRANSLATION

Translation is considered a kind of modification, so you may distribute translations of the Document under the terms of section 4. Replacing Invariant Sections with translations requires special permission from their copyright holders, but you may include translations of some or all Invariant Sections in addition to the original versions of these Invariant Sections. You may include a translation of this License, and all the license notices in the Document, and any Warranty Disclaimers, provided that you also include the original English version of this License and the original versions of those notices and disclaimers. In case of a disagreement between the translation and the original version of this License or a notice or disclaimer, the original version will prevail.

If a section in the Document is Entitled "Acknowledgements", "Dedications", or "History", the requirement (section 4) to Preserve its Title (section 1) will typically require changing the actual title.

## 9. TERMINATION

You may not copy, modify, sublicense, or distribute the Document except as expressly provided for under this License. Any other attempt to copy, modify, sublicense or distribute the Document is void, and will automatically terminate your rights under this License. However, parties who have received copies, or rights, from you under this License will not have their licenses terminated so long as such parties remain in full compliance.

## 10. FUTURE REVISIONS OF THIS LICENSE

The Free Software Foundation may publish new, revised versions of the GNU Free Documentation License from time to time. Such new versions will be similar in spirit to the present version, but may differ in detail to address new problems or concerns. See <http://www.gnu.org/copyleft/>.

Each version of the License is given a distinguishing version number. If the Document specifies that a particular numbered version of this License "or any later version" applies to it, you have the option of following the terms and conditions either of that specified version or of any later version that has been published (not as a draft) by the Free Software Foundation. If the Document does not specify a version number of this License, you may choose any version ever published (not as a draft) by the Free Software Foundation.

***Que a quien robe este libro, o lo tome prestado y no lo devuelva, se le convierta en una serpiente en las manos y lo venza. Que sea golpeado por la parálisis y todos sus miembros arruinados. Que languidezca de dolor gritando por piedad, y que no haya coto a su agonía hasta la última disolución. Que las polillas roan sus entrañas y, cuando llegue al final de su castigo, que arda en las llamas del Infierno para siempre.***  
*-Maldición anónima contra los ladrones de libros en el monasterio de San Pedro, Barcelona.*

# Contents

<b>GNU Free Documentation License</b>	<b>iii</b>
1. APPLICABILITY AND DEFINITIONS	iii
2. VERBATIM COPYING	iii
3. COPYING IN QUANTITY	iii
4. MODIFICATIONS	iii
5. COMBINING DOCUMENTS	iv
6. COLLECTIONS OF DOCUMENTS	iv
7. AGGREGATION WITH INDEPENDENT WORKS	iv
8. TRANSLATION	iv
9. TERMINATION	iv
10. FUTURE REVISIONS OF THIS LICENSE	iv
<b>Preface</b>	<b>viii</b>
<b>1 Preliminaries</b>	<b>1</b>
1.1 Sets	1
Homework	3
1.2 Numerical Functions	4
1.2.1 Injective and Surjective Functions	5
1.2.2 Algebra of Functions	6
1.2.3 Inverse Image	6
1.2.4 Inverse Function	7
Homework	8
1.3 Countability	9
Homework	10
1.4 Groups and Fields	10
Homework	12
1.5 Addition and Multiplication in $\mathbb{R}$	13
Homework	18
1.6 Order Axioms	19
1.6.1 Absolute Value	20
Homework	21
1.7 Classical Inequalities	21
1.7.1 Triangle Inequality	21
1.7.2 Bernoulli's Inequality	23
1.7.3 Rearrangement Inequality	23
1.7.4 Arithmetic Mean-Geometric Mean Inequality	24
1.7.5 Cauchy-Bunyakovsky-Schwarz Inequality	26
1.7.6 Minkowski's Inequality	27
Homework	28
1.8 Completeness Axiom	29
1.8.1 Greatest Integer Function	32
Homework	32

<b>2</b>	<b>Topology of <math>\mathbb{R}</math></b>	<b>34</b>
2.1	Intervals	34
2.2	Dense Sets	36
	Homework	37
2.3	Open and Closed Sets	37
2.4	Interior, Boundary, and Closure of a Set	39
2.5	Connected Sets	41
2.6	Compact Sets	41
	Homework	43
2.7	$\mathbb{R}$	44
2.8	Lebesgue Measure	45
2.9	The Cantor Set	45
<b>3</b>	<b>Sequences</b>	<b>47</b>
3.1	Limit of a Sequence	47
3.2	Convergence of Sequences	47
	Homework	51
3.3	Classical Limits of Sequences	53
	Homework	58
3.4	Averages of Sequences	59
	Homework	62
3.5	Orders of Infinity	62
	Homework	66
3.6	Cauchy Sequences	66
	Homework	67
3.7	Topology of sequences. Limit Superior and Limit Inferior	67
	Homework	68
<b>4</b>	<b>Series</b>	<b>70</b>
4.1	Convergence and Divergence of Series	70
	Homework	72
4.2	Convergence and Divergence of Series of Positive Terms	73
	Homework	78
4.3	Summation by Parts	79
4.4	Alternating Series	80
4.5	Absolute Convergence	80
<b>5</b>	<b>Real Functions of One Real Variable</b>	<b>81</b>
5.1	Limits of Functions	81
	Homework	85
5.2	Continuity	86
	Homework	89
5.3	Algebraic Operations with Continuous Functions	89
5.4	Monotonicity and Inverse Image	90
5.5	Convex Functions	91
5.5.1	Graphs of Functions	91
5.6	Classical Functions	92
5.6.1	Affine Functions	92
5.6.2	Quadratic Functions	92
5.6.3	Polynomial Functions	92
5.6.4	Exponential Functions	92
5.6.5	Logarithmic Functions	92
5.6.6	Trigonometric Functions	92
	Homework	93
5.6.7	Inverse Trigonometric Functions	93
5.7	Continuity of Some Standard Functions.	93
5.7.1	Continuity Polynomial Functions	93

5.7.2	Continuity of the Exponential and Logarithmic Functions . . . . .	93
5.7.3	Continuity of the Power Functions . . . . .	94
	Homework . . . . .	94
5.8	Inequalities Obtained by Continuity Arguments . . . . .	95
	Homework . . . . .	99
5.9	Intermediate Value Property . . . . .	99
	Homework . . . . .	101
5.10	Variation of a Function and Uniform Continuity . . . . .	101
	Homework . . . . .	105
5.11	Classical Limits . . . . .	105
<b>6</b>	<b>Differentiable Functions</b> . . . . .	<b>109</b>
6.1	Derivative at a Point . . . . .	109
	Homework . . . . .	109
6.2	Differentiation Rules . . . . .	110
	Homework . . . . .	115
6.3	Rolle's Theorem and the Mean Value Theorem . . . . .	116
	Homework . . . . .	117
6.4	Extrema . . . . .	118
	Homework . . . . .	119
6.5	Convex Functions . . . . .	119
	Homework . . . . .	122
6.6	Inequalities Obtained Through Differentiation . . . . .	123
	Homework . . . . .	125
6.7	Asymptotic Preponderance . . . . .	126
	Homework . . . . .	130
6.8	Asymptotic Equivalence . . . . .	130
	Homework . . . . .	135
6.9	Asymptotic Expansions . . . . .	136
	Homework . . . . .	138
<b>7</b>	<b>Integrable Functions</b> . . . . .	<b>139</b>
7.1	The Area Problem . . . . .	139
	Homework . . . . .	148
7.2	Integration . . . . .	148
	Homework . . . . .	153
7.3	Riemann-Stieltjes Integration . . . . .	155
7.4	Euler's Summation Formula . . . . .	155
<b>8</b>	<b>Sequences and Series of Functions</b> . . . . .	<b>156</b>
8.1	Pointwise Convergence . . . . .	156
8.2	Uniform Convergence . . . . .	156
8.3	Integrals and Derivatives of Sequences of Functions . . . . .	157
8.4	Power Series . . . . .	157
8.5	Maclaurin Expansions to know by inspection . . . . .	157
	Homework . . . . .	158
8.6	Comparison Tests . . . . .	158
	Homework . . . . .	158
8.7	Taylor Polynomials . . . . .	158
	Homework . . . . .	158
8.8	Abel's Theorem . . . . .	158
	Homework . . . . .	158
<b>A</b>	<b>Answers and Hints</b> . . . . .	<b>159</b>
	Answers and Hints . . . . .	159

# Preface

For many years I have been lucky enough to have students ask for more: more challenging problems, more illuminating proofs to different theorems, a deeper look at various topics, etc. To those students I normally recommend the books in the bibliography. Some of the same students have complained of not finding the books or wanting to buy them, but being impecunious, not being able to afford to buy them. Hence I have decided to make this compilation.

Here we take a semi-rigorous tour through Calculus. We don't construct the real numbers, but we examine closer the real number axioms and some of the basic theorems of Calculus. We also consider some Olympiad-level problems whose solution can be obtained through Calculus.

The reader is assumed to be familiar with proofs using mathematical induction, proofs by contradiction, and the mechanics of differentiation and integration.

David A. SANTOS  
[dsantos@ccp.edu](mailto:dsantos@ccp.edu)



# Chapter 1

## Preliminaries


**Why bother?** We will use the language of set theory throughout these notes. There are various elementary results that pop up in later proofs, among them, the De Morgan Laws and the Monotonicity Reversing of Complementation Rule.

The concept of a *function* lies at the core of mathematics. We will give a brief overview here of some basic properties of functions.

### 1.1 Sets

This section contains some of the set notation to be used throughout these notes. The one-directional arrow  $\implies$  reads “implies” and the two-directional arrow  $\iff$  reads “if and only if.”

**1 Definition** We will accept the notion of *set* as a primitive notion, that is, a notion that cannot be defined in terms of more elementary notions. By a *set* we will understand a well-defined collection of objects, which we will call the *elements* of the set. If the element  $x$  belongs to the set  $S$  we will write  $x \in S$ , and in the contrary case we will write  $x \notin S$ .<sup>1</sup> The *cardinality* of a set is the number of elements the set has. It can either be finite or infinite. We will denote the cardinality of the set  $S$  by  $\text{card}(S)$ .

 Some sets are used so often that merit special notation. We will denote by

$$\mathbb{N} = \{0, 1, 2, 3, \dots\}$$

the set of natural numbers, by

$$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$$
<sup>2</sup>

by  $\mathbb{Q}$  the set of rational numbers<sup>3</sup>, by  $\mathbb{R}$  the real numbers, and by  $\mathbb{C}$  the set of complex numbers. We will occasionally also use  $\alpha\mathbb{Z} = \{\dots, -3\alpha, -2\alpha, -\alpha, 0, \alpha, 2\alpha, 3\alpha, \dots\}$ , etc.

We will also denote the empty set, that is, the set having no elements by  $\emptyset$ .

**2 Definition** The *union* of two sets  $A$  and  $B$  is the set

$$A \cup B = \{x : (x \in A) \text{ or } (x \in B)\}.$$




This is read “ $A$  union  $B$ .” See figure 1.1. The *intersection* of two sets  $A$  and  $B$  is

$$A \cap B = \{x : (x \in A) \text{ and } (x \in B)\}.$$

<sup>1</sup> Georg Cantor(1845-1918), the creator of set theory, said “A set is any collection into a whole of definite, distinguishable objects, called **elements**, of our intuition or thought.”

<sup>2</sup> $\mathbb{Z}$  for the German word *Zählen* meaning “integer.”

<sup>3</sup> $\mathbb{Q}$  for “quotients.”

Figure 1.1:  $A \cup B$ Figure 1.2:  $A \cap B$ Figure 1.3:  $A \setminus B$ 

This is read “ $A$  intersection  $B$ .” See figure 1.2. The *set difference* of two sets  $A$  and  $B$  is

$$A \setminus B = \{x : (x \in A) \text{ and } (x \notin B)\}.$$

This is read “ $A$  set minus  $B$ .” See figure 1.3.

**3 Definition** Two sets  $A$  and  $B$  are *disjoint* if  $A \cap B = \emptyset$ .

**4 Example** Write  $A \cup B$  as the disjoint union of three sets.

**Solution:** Observe that

$$A \cup B = (A \setminus B) \cup (A \cap B) \cup (B \setminus A),$$

and that the sets on the dextral side are disjoint.

**5 Definition** A *subset*  $B$  of a set  $A$  is a subcollection of  $A$ , and we denote this by  $B \subseteq A$ .<sup>4</sup> This means that  $x \in B \implies x \in A$ .



$\emptyset$  and  $A$  are always subsets of any set  $A$ .

Observe that

$$A = B \iff (A \subseteq B) \text{ and } (B \subseteq A).$$

We use this observation on the next theorem.

**6 THEOREM (De Morgan Laws)** Let  $A, B, C$  be sets. Then

$$A \setminus (B \cap C) = (A \setminus B) \cup (A \setminus C), \quad A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C).$$

**Proof:** We have

$$\begin{aligned} x \in A \setminus (B \cup C) &\iff x \in A \text{ and } x \notin (B \text{ or } C) \\ &\iff (x \in A) \text{ and } ((x \notin B) \text{ and } (x \notin C)) \\ &\iff (x \in A \text{ and } x \notin B) \text{ and } (x \in A \text{ and } x \notin C) \\ &\iff (x \in A \setminus B) \text{ and } (x \in A \setminus C) \\ &\iff x \in (A \setminus B) \cap (A \setminus C). \end{aligned}$$

Also,

$$\begin{aligned} x \in A \setminus (B \cap C) &\iff x \in A \text{ and } x \notin (B \text{ and } C) \\ &\iff (x \in A) \text{ and } ((x \notin B) \text{ or } (x \notin C)) \\ &\iff (x \in A \text{ and } x \notin B) \text{ or } (x \in A \text{ and } x \notin C) \\ &\iff (x \in A \setminus B) \text{ or } (x \in A \setminus C) \\ &\iff x \in (A \setminus B) \cup (A \setminus C) \end{aligned}$$

□

<sup>4</sup>There seems not to be an agreement here by authors. Some use the notation  $\subset$  or  $\subseteq$  instead of  $\subseteq$ . Some see in the notation  $\subset$  the exclusion of equality. In these notes, we will always use the notation  $\subseteq$ , and if we wished to exclude equality we will write  $\subsetneq$ .

**7 THEOREM (Monotonicity Reversing of Complementation)** Let  $A, B, X$  be sets. Then

$$A \subseteq B \iff X \setminus B \subseteq X \setminus A.$$

**Proof:** We have

$$\begin{aligned} A \subseteq B &\iff (x \in A) \implies (x \in B) \\ &\iff (x \notin B) \implies (x \notin A) \\ &\iff (x \in X \text{ and } x \notin B) \implies (x \in X \text{ and } x \notin A) \\ &\iff X \setminus B \subseteq X \setminus A. \end{aligned}$$

□

**8 Definition** Let  $A_1, A_2, \dots, A_n$ , be sets. The *Cartesian Product* of these  $n$  sets is defined and denoted by

$$A_1 \times A_2 \times \dots \times A_n = \{(a_1, a_2, \dots, a_n) : a_k \in A_k\},$$

that is, the set of all ordered  $n$ -tuples whose elements belong to the given sets.



In the particular case when all the  $A_k$  are equal to a set  $A$ , we write

$$A_1 \times A_2 \times \dots \times A_n = A^n.$$

If  $a \in A$  and  $b \in A$  we write  $(a, b) \in A^2$ .

**9 Example** The Cartesian product is not necessarily commutative. For example,  $(\sqrt{2}, 1) \in \mathbb{R} \times \mathbb{Z}$  but  $(\sqrt{2}, 1) \notin \mathbb{Z} \times \mathbb{R}$ . Since  $\mathbb{R} \times \mathbb{Z}$  has an element that  $\mathbb{Z} \times \mathbb{R}$  does not,  $\mathbb{R} \times \mathbb{Z} \neq \mathbb{Z} \times \mathbb{R}$ .

**10 Example** Prove that if  $X \times X = Y \times Y$  then  $X = Y$ .

**Solution:** Let  $x \in X$ . Then  $(x, x) \in X \times X$ , which gives  $(x, x) \in Y \times Y$ , so  $y \in Y$ . Hence  $X \subseteq Y$ .

Similarly, if  $y \in Y$  then  $(y, y) \in Y \times Y$ , which gives  $(y, y) \in X \times X$ , so  $y \in X$ . Hence  $Y \subseteq X$ .

Thus  $X \subseteq Y$  and  $Y \subseteq X$  gives  $X = Y$ .

## Homework

**Problem 1.1.1** For a fixed  $n \in \mathbb{N}$  put  $A_n = \{nk : k \in \mathbb{N}\}$ .

1. Find  $A_2 \cap A_3$ .
2. Find  $\bigcap_{n=1}^{\infty} A_n$ .
3. Find  $\bigcup_{n=1}^{\infty} A_n$ .

**Problem 1.1.2** Prove the following properties of the empty set:

$$A \cap \emptyset = \emptyset, \quad A \cup \emptyset = A.$$

**Problem 1.1.3** Prove the following commutative laws:

$$A \cap B = B \cap A, \quad A \cup B = B \cup A.$$

**Problem 1.1.4** Prove by means of set inclusion the following distributive law:

$$(A \cup B) \cap C = (A \cap C) \cup (B \cap C).$$

**Problem 1.1.5** Prove the following associative laws:

$$A \cap (B \cap C) = (A \cap B) \cap C, \quad A \cup (B \cup C) = (A \cup B) \cup C.$$

**Problem 1.1.6** Prove that

$$A \cap B = A \iff A \subseteq B.$$

**Problem 1.1.7** Prove that

$$A \cup B = A \iff B \subseteq A.$$

**Problem 1.1.8** Prove that

$$A \subseteq B \implies A \cap C \subseteq B \cap C.$$

**Problem 1.1.9** Prove that

$$A \subseteq B \text{ and } C \subseteq B \implies A \cup C \subseteq B.$$

**Problem 1.1.10** Prove the following distributive laws:  
 $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$ ,  $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$ .

**Problem 1.1.11** Is there any difference between the sets  $\emptyset$ ,  $\{\emptyset\}$  and  $\{\{\emptyset\}\}$ ? Explain.

**Problem 1.1.12** Is the Cartesian product associative? Explain.

**Problem 1.1.13** Let  $A, B$ , and  $C$  be sets. Show that

$$A \times (B \setminus C) = (A \times B) \setminus (A \times C).$$

**Problem 1.1.14** Prove that a set with  $N \in \mathbb{N}$  elements has exactly  $2^N$  subsets.

## 1.2 Numerical Functions

**11 Definition** By a (numerical) function  $f : \text{Dom}(f) \rightarrow \text{Target}(f)$  we mean the collection of the following ingredients:

- ❶ a name for the function. Usually we use the letter  $f$ .
- ❷ a set of real number inputs called the domain of the function. The domain of  $f$  is denoted by  $\text{Dom}(f) \subseteq \mathbb{R}$ .
- ❸ an input parameter, also called independent variable or dummy variable. We usually denote a typical input by the letter  $x$ .
- ❹ a set of possible real number outputs of the function, called the target set of the function. The target set of  $f$  is denoted by  $\text{Target}(f) \subseteq \mathbb{R}$ .
- ❺ an assignment rule or formula, assigning to every input a unique output. This assignment rule for  $f$  is usually denoted by  $x \mapsto f(x)$ . The output of  $x$  under  $f$  is also referred to as the image of  $x$  under  $f$ , and is denoted by  $f(x)$ .

The notation<sup>5</sup>

$$f : \begin{array}{ccc} \text{Dom}(f) & \rightarrow & \text{Target}(f) \\ x & \mapsto & f(x) \end{array}$$

read “the function  $f$ , with domain  $\text{Dom}(f)$ , target set  $\text{Target}(f)$ , and assignment rule  $f$  mapping  $x$  to  $f(x)$ ” conveys all the above ingredients.



Oftentimes we will only need to mention the assignment rule of a function, without mentioning its domain or target set. In such instances we will sloppily say “the function  $f$ ” or more commonly, “the function  $x \mapsto f(x)$ ”; e.g., the square function  $x \mapsto x^2$ .<sup>6</sup>

**12 Definition** The image  $\text{Im}(f)$  of a function  $f$  is its set of actual outputs. In other words,

$$\text{Im}(f) = \{f(a) : a \in \text{Dom}(f)\}.$$

Observe that we always have  $\text{Im}(f) \subseteq \text{Target}(f)$ . For a set  $A$ , we also define

$$f(A) = \{f(a) : a \in A\}.$$

**13 THEOREM** Let  $f : X \rightarrow Y$  be a function and let  $A \subseteq X$ ,  $A' \subseteq X$ . Then

1.  $A \subseteq A' \implies f(A) \subseteq f(A')$
2.  $f(A \cup A') = f(A) \cup f(A')$
3.  $f(A \cap A') \subseteq f(A) \cap f(A')$
4.  $f(A) \setminus f(A') \subseteq f(A \setminus A')$

**Proof:**

<sup>5</sup>Notice the difference in the arrows. The straight arrow  $\rightarrow$  is used to mean that a certain set is associated with another set, whereas the arrow  $\mapsto$  (read “maps to”) is used to denote that an input becomes a certain output.

<sup>6</sup>This corresponds to the even sloppier American usage “the function  $f(x) = x^2$ .”

1.  $x \in A \implies x \in A'$  and hence  $f(x) \in f(A) \implies f(x) \in f(A') \implies f(A) \subseteq f(A')$
2. Since  $A \subseteq A \cup A'$  and  $A' \subseteq A \cup A'$ , we have  $f(A) \subseteq f(A \cup A')$  and  $f(A') \subseteq f(A \cup A')$ , by part (1) and thus  $f(A) \subseteq f(A') \subseteq f(A \cup A')$ . Moreover, if  $y \in f(A \cup A')$ , then  $\exists x \in A \cup A'$  such that  $y = f(x)$ . Then either  $x \in A$  and so  $f(x) \in f(A)$  or  $x \in A'$  and so  $f(x) \in f(A')$ . Either way,  $f(x) \in f(A) \cup f(A')$  and

$$y \in f(A \cup A') \implies y \in f(A) \cup f(A') \implies f(A \cup A') \subseteq f(A) \cup f(A').$$

Hence

$$f(A \cup A') = f(A) \cup f(A').$$

3. Let  $y \in f(A \cap A')$ . Then  $\exists x \in A \cap A'$  such that  $f(x) = y$ . Thus we have both  $x \in A \implies f(x) \in f(A)$  and  $x \in A' \implies f(x) \in f(A')$ . Therefore  $f(x) \in f(A) \cap f(A')$  and we conclude that  $f(A \cap A') \subseteq f(A) \cap f(A')$ .
4. Let  $y \in f(A) \setminus f(A')$ . Then  $y \in f(A)$  and  $y \notin f(A')$ . Thus  $\exists x \in A$  such that  $f(x) = y$ . Since  $y \notin f(A')$ , then  $x \notin A'$ . Therefore  $x \in A \setminus A'$  and finally,  $y \in f(A \setminus A')$ . This means that  $f(A) \setminus f(A') \subseteq f(A \setminus A')$  as claimed.

□

### 1.2.1 Injective and Surjective Functions

**14 Definition** A function is *injective* or *one-to-one* whenever two different values of its domain generate two different values in its image. A function is *surjective* or *onto* if every element of its target set is hit, that is, the target set is the same as the image of the function. A function is *bijective* if it is both injective and surjective.

**15 Example** The function

$$a: \begin{array}{ccc} \mathbb{R} & \rightarrow & \mathbb{R} \\ x & \mapsto & x^2 \end{array}$$

is neither injective nor surjective.

The function

$$b: \begin{array}{ccc} \mathbb{R} & \rightarrow & [0; +\infty[ \\ x & \mapsto & x^2 \end{array}$$

is surjective but not injective.

The function

$$c: \begin{array}{ccc} [0; +\infty[ & \rightarrow & \mathbb{R} \\ x & \mapsto & x^2 \end{array}$$

is injective but not surjective.

The function

$$d: \begin{array}{ccc} [0; +\infty[ & \rightarrow & [0; +\infty[ \\ x & \mapsto & x^2 \end{array}$$

is a bijection.

A bijection between two sets essentially tells us that the two sets have the same size. We will make this statement more precise now for finite sets.

**16 THEOREM** Let  $f: A \rightarrow B$  be a function, and let  $A$  and  $B$  be finite. If  $f$  is injective, then  $\text{card}(A) \leq \text{card}(B)$ . If  $f$  is surjective then  $\text{card}(B) \leq \text{card}(A)$ . If  $f$  is bijective, then  $\text{card}(A) = \text{card}(B)$ .

**Proof:** Put  $n = \text{card}(A)$ ,  $A = \{x_1, x_2, \dots, x_n\}$  and  $m = \text{card}(B)$ ,  $B = \{y_1, y_2, \dots, y_m\}$ .

If  $f$  were injective then  $f(x_1), f(x_2), \dots, f(x_n)$  are all distinct, and among the  $y_k$ . Hence  $n \leq m$ .

If  $f$  were surjective then each  $y_k$  is hit, and for each, there is an  $x_i$  with  $f(x_i) = y_k$ . Thus there are at least  $m$  different images, and so  $n \geq m$ . □

### 1.2.2 Algebra of Functions

**17 Definition** Let  $f : \text{Dom}(f) \rightarrow \text{Target}(f)$  and  $g : \text{Dom}(g) \rightarrow \text{Target}(g)$ . Then  $\text{Dom}(f \pm g) = \text{Dom}(f) \cap \text{Dom}(g)$  and the sum (respectively, difference) function  $f + g$  (respectively,  $f - g$ ) is given by

$$f \pm g : \begin{array}{ccc} \text{Dom}(f) \cap \text{Dom}(g) & \rightarrow & \text{Target}(f \pm g) \\ x & \mapsto & f(x) \pm g(x) \end{array} .$$

In other words, if  $x$  belongs both to the domain of  $f$  and  $g$ , then

$$(f \pm g)(x) = f(x) \pm g(x).$$

**18 Definition** Let  $f : \text{Dom}(f) \rightarrow \text{Target}(f)$  and  $g : \text{Dom}(g) \rightarrow \text{Target}(g)$ . Then  $\text{Dom}(fg) = \text{Dom}(f) \cap \text{Dom}(g)$  and the product function  $fg$  is given by

$$fg : \begin{array}{ccc} \text{Dom}(f) \cap \text{Dom}(g) & \rightarrow & \text{Target}(fg) \\ x & \mapsto & f(x) \cdot g(x) \end{array} .$$

In other words, if  $x$  belongs both to the domain of  $f$  and  $g$ , then

$$(fg)(x) = f(x) \cdot g(x).$$

**19 Definition** Let  $g : \text{Dom}(g) \rightarrow \text{Target}(g)$  be a function. The *support* of  $g$ , denoted by  $\text{supp}(g)$  is the set of elements in  $\text{Dom}(g)$  where  $g$  does not vanish, that is

$$\text{supp}(g) = \{x \in \text{Dom}(g) : g(x) \neq 0\}.$$

**20 Definition** Let  $f : \text{Dom}(f) \rightarrow \text{Target}(f)$  and  $g : \text{Dom}(g) \rightarrow \text{Target}(f)$ . Then  $\text{Dom}\left(\frac{f}{g}\right) = \text{Dom}(f) \cap \text{supp}(g)$  and the quotient function  $\frac{f}{g}$  is given by

$$\frac{f}{g} : \begin{array}{ccc} \text{Dom}(f) \cap \text{supp}(g) & \rightarrow & \text{Target}(f/g) \\ x & \mapsto & \frac{f(x)}{g(x)} \end{array} .$$

In other words, if  $x$  belongs both to the domain of  $f$  and  $g$  and  $g(x) \neq 0$ , then  $\frac{f}{g}(x) = \frac{f(x)}{g(x)}$ .

**21 Definition** Let  $f : \text{Dom}(f) \rightarrow \text{Target}(f)$ ,  $g : \text{Dom}(g) \rightarrow \text{Target}(g)$  and let  $U = \{x \in \text{Dom}(g) : g(x) \in \text{Dom}(f)\}$ . We define the *composition* function of  $f$  and  $g$  as

$$f \circ g : \begin{array}{ccc} U & \rightarrow & \text{Target}(f \circ g) \\ x & \mapsto & f(g(x)) \end{array} . \tag{1.1}$$

We read  $f \circ g$  as “ $f$  composed with  $g$ .”

### 1.2.3 Inverse Image

**22 Definition** Let  $X$  and  $Y$  be subsets of  $\mathbb{R}$  and let  $f : X \rightarrow Y$  be a function. Let  $B \subseteq Y$ . The *inverse image* of  $B$  by  $f$  is the set

$$f^{-1}(B) = \{x \in X : f(x) \in B\}.$$

If  $B = \{b\}$  consists of only one element, we write, abusing notation,  $f^{-1}(\{b\}) = f^{-1}(b)$ . It is clear that  $f^{-1}(Y) = X$  and  $f^{-1}(\emptyset) = \emptyset$ .

**23 Example** Let

$$f : \begin{array}{ccc} \{-2, -1, 0, 1, 3\} & \rightarrow & \{0, 1, 4, 5, 9\} \\ x & \mapsto & x^2 \end{array} .$$

Then  $f^{-1}(\{0, 1\}) = \{0, -1, 1\}$ ,  $f^{-1}(1) = \{-1, 1\}$ ,  $f^{-1}(5) = \emptyset$ ,  $f^{-1}(4) = 2$ ,  $f^{-1}(0) = 0$ , etc. Notice that we have abused notation in all but the first example.

**24 THEOREM** Let  $f : X \rightarrow Y$  be a function and let  $B \subseteq Y, B' \subseteq Y$ . Then

1.  $B \subseteq B' \implies f^{-1}(B) \subseteq f^{-1}(B')$
2.  $f^{-1}(B \cup B') = f^{-1}(B) \cup f^{-1}(B')$
3.  $f^{-1}(B \cap B') = f^{-1}(B) \cap f^{-1}(B')$
4.  $f^{-1}(B) \setminus f^{-1}(B') = f^{-1}(B \setminus B')$

**Proof:**

1. Assume  $x \in f^{-1}(B)$ . Then there is  $y \in B \subseteq B'$  such that  $f(x) = y$ . But  $y$  is also in  $B'$  so  $x \in f^{-1}(B')$ . Thus  $f^{-1}(B) \subseteq f^{-1}(B')$ .
2. Since  $B \subseteq B \cup B'$  and  $B' \subseteq B \cup B'$ , we have  $f^{-1}(B) \subseteq f^{-1}(B \cup B')$  and  $f^{-1}(B') \subseteq f^{-1}(B \cup B')$ , by part (1). Thus  $f^{-1}(B) \cup f^{-1}(B') \subseteq f^{-1}(B \cup B')$ . Now, let  $x \in f^{-1}(B \cup B')$ . There is  $y \in B \cup B'$  such that  $f(x) = y$ . Either  $y \in B$  and so  $y \in B \implies x \in f^{-1}(B)$  or  $y \in B'$  and so  $y \in B' \implies x \in f^{-1}(B')$ . Either way,  $x \in f^{-1}(B) \cup f^{-1}(B')$ . Thus  $f^{-1}(B \cup B') \subseteq f^{-1}(B) \cup f^{-1}(B')$ . We conclude that  $f^{-1}(B \cup B') = f^{-1}(B) \cup f^{-1}(B')$ .
3. Let  $x \in f^{-1}(B \cap B')$ . Then  $\exists y \in B \cap B'$  such that  $f(x) = y$ . Thus we have both  $y \in B \implies x \in f^{-1}(B)$  and  $y \in B' \implies x \in f^{-1}(B')$ . Therefore  $x \in f^{-1}(B) \cap f^{-1}(B')$  and we conclude that  $f^{-1}(B \cap B') \subseteq f^{-1}(B) \cap f^{-1}(B')$ . Now, let  $x \in f^{-1}(B) \cap f^{-1}(B')$ . Then  $x \in f^{-1}(B)$  and  $x \in f^{-1}(B')$ . Then  $f(x) \in B$  and  $f(x) \in B'$ . Thus  $f(x) \in B \cap B'$  and so  $x \in f^{-1}(B \cap B')$ . Hence  $f^{-1}(B) \cap f^{-1}(B') \subseteq f^{-1}(B \cap B')$  also, and we conclude that  $f^{-1}(B) \cap f^{-1}(B') = f^{-1}(B \cap B')$ .
4. Let  $x \in f^{-1}(B) \setminus f^{-1}(B')$ . Then  $x \in f^{-1}(B)$  and  $x \notin f^{-1}(B')$ . Thus  $f(x) \in B$  and  $f(x) \notin B'$ . Thus  $f(x) \in B \setminus B'$  and therefore  $x \in f^{-1}(B \setminus B')$ , giving  $f^{-1}(B) \setminus f^{-1}(B') \subseteq f^{-1}(B \setminus B')$ . Now, let  $x \in f^{-1}(B \setminus B')$ . Then  $f(x) \in B \setminus B'$ , which means that  $f(x) \in B$  but  $f(x) \notin B'$ . Thus  $x \in f^{-1}(B)$  but  $x \notin f^{-1}(B')$ , which gives  $x \in f^{-1}(B) \setminus f^{-1}(B')$  and so  $f^{-1}(B \setminus B') \subseteq f^{-1}(B) \setminus f^{-1}(B')$ . This establishes the desired equality.

□

**25 THEOREM** Let  $f : X \rightarrow Y$  be a function. Let  $A \times B \subseteq X \times Y$ . Then

1.  $A \subseteq (f^{-1} \circ f)(A)$
2.  $(f \circ f^{-1})(B) \subseteq B$

**Proof:** We have

1. Let  $x \in A$ . Then  $\exists y \in Y$  such that  $y = f(x)$ . Thus  $y \in f(A)$ . Therefore  $x \in f^{-1}(f(A))$ .
2.  $y \in (f \circ f^{-1})(B)$ . Then  $\exists x \in f^{-1}(B)$  such that  $f(x) = y$ . Thus  $x \in f^{-1}(y)$ . Hence  $f(x) \in B$ . Therefore  $y \in B$ .

□

### 1.2.4 Inverse Function

**26 Definition** Let  $A \times B \subseteq \mathbb{R}^2$ . A function  $F : A \rightarrow B$  is said to be *invertible* if there exists a function  $F^{-1}$  (called the *inverse* of  $F$ ) such that  $F \circ F^{-1} = \text{Id}_B$  and  $F^{-1} \circ F = \text{Id}_A$ . Here  $\text{Id}_S$  is the identity on the set  $S$  function with rule  $\text{Id}_S(x) = x$ .

The central question is now: given a function  $F : A \rightarrow B$ , when is  $F^{-1} : B \rightarrow A$  a function? The answer is given in the next theorem.

**27 THEOREM** Let  $A \times B \subseteq \mathbb{R}^2$ . A function  $f : A \rightarrow B$  is invertible if and only if it is a bijection. That is,  $f^{-1} : B \rightarrow A$  is a function if and only if  $f$  is bijective.

**Proof:** Assume first that  $f$  is invertible. Then there is a function  $f^{-1} : B \rightarrow A$  such that

$$f \circ f^{-1} = \text{Id}_B \text{ and } f^{-1} \circ f = \text{Id}_A. \tag{1.2}$$

Let us prove that  $f$  is injective and surjective. Let  $s, t$  be in the domain of  $f$  and such that  $f(s) = f(t)$ . Applying  $f^{-1}$  to both sides of this equality we get  $(f^{-1} \circ f)(s) = (f^{-1} \circ f)(t)$ . By the definition of inverse function,  $(f^{-1} \circ f)(s) = s$  and  $(f^{-1} \circ f)(t) = t$ . Thus  $s = t$ . Hence  $f(s) = f(t) \implies s = t$  implying that  $f$  is injective. To prove that  $f$  is surjective we must show that for every  $b \in f(A) \exists a \in A$  such that  $f(a) = b$ . We take  $a = f^{-1}(b)$  (observe that  $f^{-1}(b) \in A$ ). Then  $f(a) = f(f^{-1}(b)) = (f \circ f^{-1})(b) = b$  by definition of inverse function. This shows that  $f$  is surjective. We conclude that if  $f$  is invertible then it is also a bijection.

Assume now that  $f$  is a bijection. For every  $b \in B$  there exists a unique  $a$  such that  $f(a) = b$ . This makes the rule  $g : B \rightarrow A$  given by  $g(b) = a$  a function. It is clear that  $g \circ f = \text{Id}_A$  and  $f \circ g = \text{Id}_B$ . We may thus take  $f^{-1} = g$ . This concludes the proof.  $\square$

### Homework

**Problem 1.2.1** Find all functions with domain  $\{a, b\}$  and target set  $\{c, d\}$ .

**Problem 1.2.2** Let  $A, B$  be finite sets with  $\text{card}(A) = n$  and  $\text{card}(B) = m$ . Prove that

- The number of functions from  $A$  to  $B$  is  $m^n$ .
- If  $n \leq m$ , the number of injective functions from  $A$  to  $B$  is  $m(m-1)(m-2)\cdots(m-n+1)$ . If  $n > m$  there are no injective functions from  $A$  to  $B$ .

**Problem 1.2.3** Let  $A$  and  $B$  be two finite sets with  $\text{card}(A) = n$  and  $\text{card}(B) = m$ . If  $n < m$  prove that there are no surjections from  $A$  to  $B$ . If  $n \geq m$  prove that the number of surjective functions from  $A$  to  $B$  is

$$m^n - \binom{m}{1}(m-1)^n + \binom{m}{2}(m-2)^n - \binom{m}{3}(m-3)^n + \cdots + (-1)^{m-1} \binom{m}{m-1} (1)^n.$$

**Problem 1.2.4** Let  $h : \mathbb{R} \rightarrow \mathbb{R}$  be given by  $h(1-x) = 2x$ . Find  $h(3x)$ .

**Problem 1.2.5** Consider the polynomial

$$(1-x^2+x^4)^{2003} = a_0 + a_1x + a_2x^2 + \cdots + a_{8012}x^{8012}.$$

Find

- ①  $a_0$
- ②  $a_0 + a_1 + a_2 + \cdots + a_{8012}$
- ③  $a_0 - a_1 + a_2 - a_3 + \cdots - a_{8011} + a_{8012}$
- ④  $a_0 + a_2 + a_4 + \cdots + a_{8010} + a_{8012}$
- ⑤  $a_1 + a_3 + \cdots + a_{8009} + a_{8011}$

**Problem 1.2.6** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$ , be a function such that  $\forall x \in ]0; +\infty[$ ,

$$[f(x^3 + 1)]^{\sqrt{x}} = 5,$$

find the value of

$$\left[ f\left(\frac{27+y^3}{y^3}\right) \right]^{\sqrt{\frac{27}{y}}}$$

for  $y \in ]0; +\infty[$ .

**Problem 1.2.7** Let  $f$  satisfy  $f(n+1) = (-1)^{n+1}n - 2f(n), n \geq 1$  If  $f(1) = f(1001)$  find

$$f(1) + f(2) + f(3) + \cdots + f(1000).$$

**Problem 1.2.8** If  $f(a)f(b) = f(a+b) \forall a, b \in \mathbb{R}$  and  $f(x) > 0 \forall x \in \mathbb{R}$ , find  $f(0)$ . Also, find  $f(-a)$  and  $f(2a)$  in terms of  $f(a)$ .

**Problem 1.2.9** Prove that  $f : \begin{matrix} \mathbb{R} \setminus \{-1\} & \rightarrow & \mathbb{R} \setminus \{1\} \\ x & \rightarrow & \frac{x-1}{x+1} \end{matrix}$  is a bijection

and find  $f^{-1}$ .

**Problem 1.2.10** Let  $f^{[1]}(x) = f(x) = x+1, f^{[n+1]} = f \circ f^{[n]}, n \geq 1$ . Find a closed formula for  $f^{[n]}$

$(1)^n$ .

**Problem 1.2.11** Let  $f, g : [0; 1] \rightarrow \mathbb{R}$  be functions. Demonstrate that there exist  $(a, b) \in [0; 1]^2$  such that  $\frac{1}{4} \leq |f(a) + g(b) - ab|$ .

**Problem 1.2.12** Demonstrate that there is no function  $f : \mathbb{R} \setminus \{1/2\} \rightarrow \mathbb{R}$  such that

$$x \in \mathbb{R} \setminus \{1/2\} \implies f(x) \left( f\left(\frac{x-1}{2x-1}\right) \right) = x^2 + x + 1$$

**Problem 1.2.13** Find all functions  $f : \mathbb{R} \setminus \{-1, 0\} \rightarrow \mathbb{R}$  such that

$$x \in \mathbb{R} \setminus \{-1, 0\} \implies f(x) + f\left(\frac{-1}{x+1}\right) = 3x + 2.$$

**Problem 1.2.14** Let  $f^{[1]}(x) = f(x) = 2x, f^{[n+1]} = f \circ f^{[n]}, n \geq 1$ . Find a closed formula for  $f^{[n]}$

**Problem 1.2.15** Find all functions  $g : \mathbb{R} \rightarrow \mathbb{R}$  that satisfy  $g(x+y) + g(x-y) = 2x^2 + 2y^2$ .

**Problem 1.2.16** Find all the functions  $f : \mathbb{R} \rightarrow \mathbb{R}$  that satisfy  $f(xy) = yf(x)$ .



**Problem 1.2.17** Find all functions  $f: \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}$  for which

$$f(x) + 2f\left(\frac{1}{x}\right) = x.$$

**Problem 1.2.18** Find all functions  $f: \mathbb{R} \setminus \{-1\} \rightarrow \mathbb{R}$  such that

$$(f(x))^2 \cdot f\left(\frac{1-x}{1+x}\right) = 64x.$$

**Problem 1.2.19** Let  $f^{[1]} = f$  be given by  $f(x) = \frac{1}{1-x}$ . Find

- (i)  $f^{[2]}(x) = (f \circ f)(x)$ ,  
(ii)  $f^{[3]}(x) = (f \circ f \circ f)(x)$ , and  
(iii)  $f^{[69]} = \underbrace{(f \circ f \circ \dots \circ f)}_{69 \text{ compositions with itself}}(x)$ .

**Problem 1.2.20** Let  $f: A \rightarrow B$  and  $g: B \rightarrow C$  be functions. Show that (i) if  $g \circ f$  is injective, then  $f$  is injective. (ii) if  $g \circ f$  is surjective, then  $g$  is surjective.

## 1.3 Countability

**28 Definition** A set  $X$  is countable if either it is finite or if there is a bijection  $f: X \rightarrow \mathbb{N}$ , that is, the set  $X$  has as many elements as  $\mathbb{N}$ .

Any countable set can be thus enumerated a sequence

$$x_1, x_2, x_3, \dots$$

Thus the strictly positive integers can be enumerated as customarily:

$$1, 2, 3, \dots$$

Another possible enumeration<sup>7</sup> is the following

$$3, 5, 7, 9, \dots, \quad , 2 \cdot 3, 2 \cdot 5, 2 \cdot 7, 2 \cdot 9, \dots, \quad , 2^2 \cdot 3, 2^2 \cdot 5, 2^2 \cdot 7, 2^2 \cdot 9, \dots, \quad , \dots, 2^4, 2^3, 2^2, 2, 1,$$

that is, we start with the odd integers in increasing order, then  $2$  times the odd integers,  $2^2$  times the odd integers, etc., and at the end we put the powers of  $2$  in decreasing order.

**29 LEMMA** Any subset  $X \subseteq \mathbb{N}$  is countable.

**Proof:** If  $X$  is finite, then there is nothing to prove. If  $X$  is infinite, we can arrange the elements of  $X$  increasing order, say,

$$x_1 < x_2 < x_3 < \dots$$

We then map the smallest element  $x_1 \in S$  to  $1$ , the next smallest  $x_2$  to  $2$ , etc.  $\square$



Hence, even though  $2\mathbb{N} \subsetneq \mathbb{N}$ , the sets  $2\mathbb{N}$  and  $\mathbb{N}$  have the same number of elements. This can also be seen by noticing that  $f: \mathbb{N} \rightarrow 2\mathbb{N}$  given by  $x_n = 2n$  is a bijection.

**30 LEMMA** A set  $X$  is countable if and only if there is an injection  $f: X \rightarrow \mathbb{N}$ .

**Proof:** The assertion is evident if  $X$  is finite. Hence assume  $X$  is infinite. If  $f: X \rightarrow \mathbb{N}$  is an injection then  $f(X)$  is an infinite subset of  $\mathbb{N}$ . Hence there is a bijection  $g: f(X) \rightarrow \mathbb{N}$  by virtue of Lemma 29. Thus  $(g \circ f): X \rightarrow \mathbb{N}$  is a bijection.  $\square$



An obvious consequence of the above lemma is that if  $X'$  is countable and there is an injection  $f: X \rightarrow X'$  then  $X$  is countable.

**31 THEOREM**  $\mathbb{Z}$  is countable.

<sup>7</sup>Which is relevant in chaos theory, for Sarkovkii's Theorem.

**Proof:** One can take, as a bijection between the two sets, for example,  $f: \mathbb{Z} \rightarrow \mathbb{N}$ ,

$$f(x) = \begin{cases} 2x+1 & \text{if } x \geq 0 \\ -2x & \text{if } x < 0. \end{cases}$$

□


**32 THEOREM**  $\mathbb{Q}$  is countable.

**Proof:** Consider  $f: \mathbb{Q} \rightarrow \mathbb{N}$  given

$$f\left(\frac{a}{b}\right) = 2^{|a|} 3^b 5^{1+\text{signum}(a)},$$

where  $\frac{a}{b}$  is in least terms, and  $b > 0$ . By the uniqueness of the prime factorisation of an integer,  $f$  is an injection.

□

 The above theorem means that there are as many rational numbers as natural numbers. Thus the rationals can be enumerated as

$$q_1, q_2, q_3, \dots,$$

**33 THEOREM (Cantor's Diagonal Argument)**  $\mathbb{R}$  is uncountable.

**Proof:** Assume  $\mathbb{R}$  were countable so that its complete set of elements may be enumerated, say, as in the list

$$\begin{aligned} r_1 &= n_1.d_{11}d_{12}d_{13}\dots \\ r_2 &= n_2.d_{21}d_{22}d_{23}\dots \\ r_3 &= n_3.d_{31}d_{32}d_{33}\dots, \end{aligned}$$

where we have used decimal notation. Define the new real  $r = 0.d_1d_2d_3\dots$  by  $d_i = 0$  if  $d_{ii} \neq 0$  and  $d_i = 1$  if  $d_{ii} = 0$ . This is a real number (as it is a decimal), but it differs from  $r_i$  in the  $i^{\text{th}}$  decimal place. It follows that the list is incomplete and the reals are uncountable. □

**34 THEOREM** The interval  $] -1; 1[$  is uncountable.

**Proof:** Observe that the map  $f: ] -1; 1[ \rightarrow \mathbb{R}$  given by  $f(x) = \tan \frac{\pi x}{2}$  is a bijection. □

## Homework

**Problem 1.3.1** Prove that there are as many numbers in  $[0; 1]$  as in any interval  $[a; b]$  with  $a < b$ . **Problem 1.3.2** Prove that there are as many numbers in  $] -\infty; +\infty[$  as in  $] 0; +\infty[$ .

## 1.4 Groups and Fields

Here we observe the rules of the game for the operations of addition and multiplication in  $\mathbb{R}$ .

**35 Definition** Let  $S, T$  be sets. A *binary operation* is a function

$$\otimes: \begin{array}{l} S \times S \rightarrow T \\ (a, b) \mapsto \otimes(a, b) \end{array}.$$

We usually use the “infix” notation  $a \otimes b$  rather than the “prefix” notation  $\otimes(a, b)$ . If  $S = T$  then we say that the binary operation is *internal* or *closed* and if  $S \neq T$  then we say that it is *external*.

**36 Example** Ordinary addition is a closed binary operation on the sets  $\mathbb{N}$ ,  $\mathbb{Z}$ ,  $\mathbb{Q}$ ,  $\mathbb{R}$ . Ordinary subtraction is a binary operation on these sets. It is not closed on  $\mathbb{N}$ , since for example  $1 - 2 = -1 \notin \mathbb{N}$ , but it is closed in the remaining sets.

**37 Example** The operation  $\otimes : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$  given by  $a \otimes b = 1 + a \cdot b$ , where  $\cdot$  is the ordinary multiplication of real numbers is commutative but not associative. To see commutativity we have

$$a \otimes b = 1 + ab = 1 + ba = b \otimes a.$$

Now,

$$1 \otimes (1 \otimes 2) = 1 \otimes (1 + 1 \cdot 2) = 1 \otimes (3) = 1 + 1 \cdot 3 = 4, \quad \text{but} \quad (1 \otimes 1) \otimes 2 = (1 + 1 \cdot 1) \otimes 2 = 2 \otimes 2 = 1 + 2 \cdot 2 = 5,$$

so the operation is not associative.

**38 Definition** Let  $G$  be a non-empty set and  $\otimes$  be a binary operation on  $G \times G$ . Then  $\langle G, \otimes \rangle$  is called a *group* if the following axioms hold:

**G1:**  $\otimes$  is closed, that is,

$$\forall (a, b) \in G^2, \quad a \otimes b \in G,$$

**G2:**  $\otimes$  is associative, that is,

$$\forall (a, b, c) \in G^3, \quad a \otimes (b \otimes c) = (a \otimes b) \otimes c,$$

**G3:**  $G$  has an identity element, that is

$$\exists e \in G \text{ such that } \forall a \in G, \quad e \otimes a = a \otimes e = a,$$

**G4:** Every element of  $G$  is invertible, that is

$$\forall a \in G, \quad \exists a^{-1} \in G \text{ such that } a \otimes a^{-1} = a^{-1} \otimes a = e.$$



From now on, we drop the sign  $\otimes$  and rather use juxtaposition for the underlying binary operation in a given group. Thus we will say a "group  $G$ " rather than the more precise "a group  $\langle G, \otimes \rangle$ ."

**39 Definition** A group  $G$  is *abelian* if its binary operation is commutative, that is,  $\forall (a, b) \in G^2, a \otimes b = b \otimes a$ .

**40 Example**  $\langle \mathbb{Z}, + \rangle$ ,  $\langle \mathbb{Q}, + \rangle$ ,  $\langle \mathbb{R}, + \rangle$ ,  $\langle \mathbb{C}, + \rangle$  are all abelian groups under addition. The identity element is  $0$  and the inverse of  $a$  is  $-a$ .

**41 Example**  $\langle \mathbb{Q} \setminus \{0\}, \cdot \rangle$ ,  $\langle \mathbb{R} \setminus \{0\}, \cdot \rangle$ ,  $\langle \mathbb{C} \setminus \{0\}, \cdot \rangle$  are all abelian groups under multiplication. The identity element is  $1$  and the inverse of  $a$  is  $\frac{1}{a}$ .

**42 Example**  $\langle \mathbb{Z} \setminus \{0\}, \cdot \rangle$  is not a group. For example the element  $2$  does not have a multiplicative inverse.

**43 Example** Let  $V_4 = \{e, a, b, c\}$  and define  $\otimes$  by the table below.

$\otimes$	$e$	$a$	$b$	$c$
$e$	$e$	$a$	$b$	$c$
$a$	$a$	$e$	$c$	$b$
$b$	$b$	$c$	$e$	$a$
$c$	$c$	$b$	$a$	$e$

It is an easy exercise to check that  $V_4$  is an abelian group, called the *Klein Viergruppe*.

**44 THEOREM** Let  $G$  be a group. Then

1. There is only one identity element, the identity element is unique.
2. The inverse of each element is unique.
3.  $\forall (a, b) \in G^2$  we have

$$(ab)^{-1} = b^{-1}a^{-1}.$$

**Proof:**

1. Let  $e$  and  $e'$  be identity elements. Since  $e$  is an identity,  $e = ee'$ . Since  $e'$  is an identity,  $e' = ee'$ . This gives  $e = ee' = e'$ .
2. Let  $b$  and  $b'$  be inverses of  $a$ . Then  $e = ab$  and  $b'a = e$ . This gives

$$b = eb = (b'a)b = b'(ab) = b'e = b'.$$

3. We have

$$(ab)(b^{-1}a^{-1}) = a(bb^{-1})a^{-1} = a(e)a^{-1} = aa^{-1} = e.$$

Thus  $b^{-1}a^{-1}$  works as a right inverse for  $ab$ . A similar calculation shows also that it works as a left inverse. Since inverses are unique, we must have

$$(ab)^{-1} = b^{-1}a^{-1}.$$

This completes the proof.  $\square$


**45 Definition** Let  $n \in \mathbb{Z}$  and let  $G$  be a group. If  $a \in G$ , we define

$$a^0 = e,$$

$$a^{|n|} = \underbrace{a \cdot a \cdots a}_{|n| \text{ a's}}$$

and

$$a^{-|n|} = \underbrace{a^{-1} \cdot a^{-1} \cdots a^{-1}}_{|n| \text{ a}^{-1}\text{'s}}.$$


 If  $(m, n) \in \mathbb{Z}^2$ , then by associativity

$$(a^n)(a^m) = (a^m)(a^n) = a^{m+n}.$$

**46 Definition** Let  $F$  be a set having at least two elements  $\mathbf{0}_F$  and  $\mathbf{1}_F$  ( $\mathbf{0}_F \neq \mathbf{1}_F$ ) together with two binary operations  $\cdot$  (field multiplication) and  $+$  (field addition). A *field*  $\langle F, \cdot, + \rangle$  is a triplet such that  $\langle F, + \rangle$  is an abelian group with identity  $\mathbf{0}_F$ ,  $\langle F \setminus \{\mathbf{0}_F\}, \cdot \rangle$  is an abelian group with identity  $\mathbf{1}_F$  and the operations  $\cdot$  and  $+$  satisfy

$$a \cdot (b + c) = (a \cdot b) + (a \cdot c),$$

that is, field multiplication distributes over field addition.

 We will continue our practice of denoting multiplication by juxtaposition, hence the  $\cdot$  sign will be dropped.

**47 Example**  $\langle \mathbb{Q}, \cdot, + \rangle$ ,  $\langle \mathbb{R}, \cdot, + \rangle$ , and  $\langle \mathbb{C}, \cdot, + \rangle$  are all fields. The multiplicative identity in each case is  $\mathbf{1}$  and the additive identity is  $\mathbf{0}$ .

## Homework

**Problem 1.4.1** Is the set of real irrational numbers closed under addition? Under multiplication?

**Problem 1.4.2** Let

$$S = \{x \in \mathbb{Z} : \exists(a, b) \in \mathbb{Z}^2, x = a^3 + b^3 + c^3 - 3abc\}.$$

Prove that  $S$  is closed under multiplication, that is, if  $x \in S$  and  $y \in S$  then  $xy \in S$ .

**Problem 1.4.3 (Putnam, 1971)** Let  $S$  be a set and let  $\circ$  be a binary operation on  $S$  satisfying the two laws

$$(\forall x \in S)(x \circ x = x),$$

and

$$(\forall(x, y, z) \in S^3)((x \circ y) \circ z = (y \circ z) \circ x).$$

Shew that  $\circ$  is commutative.

**Problem 1.4.4 (Putnam, 1972)** Let  $\mathcal{S}$  be a set and let  $*$  be a binary operation of  $\mathcal{S}$  satisfying the laws  $\forall(x, y) \in \mathcal{S}^2$

$$x * (x * y) = y, \quad (1.3)$$

$$(y * x) * x = y. \quad (1.4)$$

Shew that  $*$  is commutative, but not necessarily associative.

**Problem 1.4.5** On  $\mathbb{Q} \cap ]-1; 1[$  define the binary operation  $\otimes$  by

$$a \otimes b = \frac{a + b}{1 + ab},$$

where juxtaposition means ordinary multiplication and  $+$  is the ordinary addition of real numbers. Prove that  $(\mathbb{Q} \cap ]-1; 1[, \otimes)$  is an abelian group by following these steps.

1. Prove that  $\otimes$  is a closed binary operation on  $\mathbb{Q} \cap ]-1; 1[$ .
2. Prove that  $\otimes$  is both commutative and associative.
3. Find an element  $e \in \mathbb{Q} \cap ]-1; 1[$  such that  $(\forall a \in \mathbb{Q} \cap ]-1; 1[)(e \otimes a = a)$ .
4. Given  $e$  as above and an arbitrary element  $a \in \mathbb{Q} \cap ]-1; 1[$ , solve the equation  $a \otimes b = e$  for  $b$ .

**Problem 1.4.6** Let  $G$  be a group satisfying  $(\forall a \in G)$

$$a^2 = e.$$

Prove that  $G$  is an abelian group.

**Problem 1.4.7** Let  $G$  be a group where  $(\forall(a, b) \in G^2)$

$$((ab)^3 = a^3 b^3) \quad \text{and} \quad ((ab)^5 = a^5 b^5).$$

Shew that  $G$  is abelian.

**Problem 1.4.8** Suppose that in a group  $G$  there exists a pair  $(a, b) \in G^2$  satisfying

$$(ab)^k = a^k b^k$$

for three consecutive integers  $k = i, i + 1, i + 2$ . Prove that  $ab = ba$ .

## 1.5 Addition and Multiplication in $\mathbb{R}$

Since  $\mathbb{R}$  is a field, it satisfies the following list of axioms, which we list for future reference.

**48 Axiom (Arithmetical Axioms of  $\mathbb{R}$ )**  $(\mathbb{R}, \cdot, +)$ —that is, the set of real numbers endowed with multiplication  $\cdot$  and addition  $+$ —is a field. This entails that  $+$  and  $\cdot$  verify the following properties.

**R1:**  $+$  and  $\cdot$  are closed binary operations, that is,

$$\forall(a, b) \in \mathbb{R}^2, \quad a + b \in \mathbb{R}, \quad a \cdot b \in \mathbb{R},$$

**R2:**  $+$  and  $\cdot$  are associative binary operations, that is,

$$\forall(a, b, c) \in \mathbb{R}^3, \quad a + (b + c) = (a + b) + c, \quad a \cdot (b \cdot c) = (a \cdot b) \cdot c$$

**R3:**  $+$  and  $\cdot$  are commutative binary operations, that is,

$$\forall(a, b) \in \mathbb{R}^2, \quad a + b = b + a, \quad a \cdot b = b \cdot a,$$

**R4:**  $\mathbb{R}$  has an additive identity element  $\mathbf{0}$ , and a multiplicative identity element  $\mathbf{1}$ , with  $\mathbf{0} \neq \mathbf{1}$ , such that

$$\forall a \in \mathbb{R}, \quad \mathbf{0} + a = a + \mathbf{0} = a, \quad \mathbf{1} \cdot a = a \cdot \mathbf{1} = a,$$

**R5:** Every element of  $\mathbb{R}$  has an additive inverse, and every element of  $\mathbb{R} \setminus \{\mathbf{0}\}$  has a multiplicative inverse, that is,

$$\forall a \in \mathbb{R}, \quad \exists(-a) \in \mathbb{R} \text{ such that } a + (-a) = (-a) + a = \mathbf{0},$$

$$\forall b \in \mathbb{R} \setminus \{\mathbf{0}\}, \quad \exists b^{-1} \in \mathbb{R} \setminus \{\mathbf{0}\} \text{ such that } b \cdot b^{-1} = b^{-1} \cdot b = \mathbf{1},$$


**R6:**  $+$  and  $\cdot$  satisfy the following distributive law:

$$\forall (a, b, c) \in \mathbb{R}^3, \quad a \cdot (b + c) = a \cdot b + a \cdot c.$$

Since  $+$  and  $\cdot$  are associative in  $\mathbb{R}$ , we may write a sum  $a_1 + a_2 + \dots + a_n$  or a product  $a_1 a_2 \dots a_n$  of real numbers without risking ambiguity. We often use the following shortcut notation.

**49 Definition** For real numbers  $a_i$  we define

$$a_1 + a_2 + \dots + a_n = \sum_{k=1}^n a_k \quad \text{and} \quad a_1 a_2 \dots a_n = \prod_{k=1}^n a_k.$$

 By convention  $\sum_{k \in \emptyset} a_k = 0$  and  $\prod_{k \in \emptyset} a_k = 1$ .

**50 THEOREM (Lagrange's Identity)** Let  $a_k, b_k$  be real numbers. Then

$$\left( \sum_{k=1}^n a_k b_k \right)^2 = \left( \sum_{k=1}^n a_k^2 \right) \left( \sum_{k=1}^n b_k^2 \right) - \sum_{1 \leq k < j \leq n} (a_k b_j - a_j b_k)^2.$$

**Proof:** For  $j = k$ ,  $a_k b_j - a_j b_k = 0$ , so we may relax the inequality in the last sum. We have

$$\begin{aligned} \sum_{1 \leq k < j \leq n} (a_k b_j - a_j b_k)^2 &= \sum_{1 \leq k \leq j \leq n} (a_k^2 b_j^2 - 2a_k b_k a_j b_j + a_j^2 b_k^2) \\ &= \sum_{1 \leq k \leq j \leq n} a_k^2 b_j^2 - 2 \sum_{1 \leq k \leq j \leq n} a_k b_k a_j b_j + \sum_{1 \leq k \leq j \leq n} a_j^2 b_k^2 \\ &= \sum_{k=1}^n \sum_{j=1}^n a_k^2 b_j^2 - \left( \sum_{k=1}^n a_k b_k \right)^2, \end{aligned}$$

proving the theorem.  $\square$

Recall that the factorial symbol  $!$  is defined by

$$0! = 1; \quad k! = k(k-1)! \quad \text{if } k \geq 1.$$

**51 Definition (Binomial Coefficients)** Let  $n \in \mathbb{N}$  We define  $\binom{n}{0} = 1 = \binom{n}{n}$  and for  $1 \leq k \leq n$ ,

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}.$$

If  $k > n$  we take  $\binom{n}{k} = 0$ .

**52 LEMMA (Pascal's Identity)** For  $n \geq 1$  and  $1 \leq k \leq n$ ,

$$\binom{n}{k} = \binom{n-1}{k} + \binom{n-1}{k-1}.$$

**Proof:** We have

$$\begin{aligned} \binom{n-1}{k} + \binom{n-1}{k-1} &= \frac{(n-1)!}{k!(n-1-k)!} + \frac{(n-1)!}{(k-1)!(n-k)!} \\ &= \frac{(n-1)!}{(k-1)!(n-1-k)!} \left( \frac{1}{k} + \frac{1}{n-k} \right) \\ &= \frac{(n-1)!}{(k-1)!(n-1-k)!} \left( \frac{n}{k(n-k)} \right) \\ &= \frac{n!}{k!(n-k)!} = \binom{n}{k}. \end{aligned}$$

□

Using Pascal's Identity we obtain *Pascal's Triangle*.

$$\begin{array}{cccccccc}
 & & & & \binom{0}{0} & & & & \\
 & & & & \binom{1}{0} & & \binom{1}{1} & & \\
 & & & \binom{2}{0} & & \binom{2}{1} & & \binom{2}{2} & \\
 & & \binom{3}{0} & & \binom{3}{1} & & \binom{3}{2} & & \binom{3}{3} \\
 & \binom{4}{0} & & \binom{4}{1} & & \binom{4}{2} & & \binom{4}{3} & \binom{4}{4} \\
 \binom{5}{0} & & \binom{5}{1} & & \binom{5}{2} & & \binom{5}{3} & & \binom{5}{4} & \binom{5}{5} \\
 \vdots & & \vdots & & \vdots & & \vdots & & \vdots & \vdots
 \end{array}$$

When the numerical values are substituted, the triangle then looks like this.

$$\begin{array}{ccccccc}
 & & & & 1 & & & \\
 & & & & 1 & & 1 & \\
 & & & 1 & 2 & & 1 & \\
 & & 1 & 3 & 3 & & 1 & \\
 & 1 & 4 & 6 & 4 & & 1 & \\
 1 & 5 & 10 & 10 & 5 & & 1 & \\
 \vdots & \vdots & \vdots & \vdots & \vdots & & \vdots & \vdots
 \end{array}$$

We see from Pascal's Triangle that binomial coefficients are symmetric. This symmetry is easily justified by the identity  $\binom{n}{k} = \binom{n}{n-k}$ . We also notice that the binomial coefficients tend to increase until they reach the middle, and that then they decrease symmetrically.

**53 THEOREM (Binomial Theorem)** For  $n \in \mathbb{N}$ ,

$$(x+y)^n = \sum_{k=0}^n \binom{n}{k} x^k y^{n-k}.$$

**Proof:** The theorem is obvious for  $n = 0$  (defining  $(x+y)^0 = 1$ ),  $n = 1$  (as  $(x+y)^1 = x+y$ ), and  $n = 2$  (as  $(x+y)^2 = x^2 + 2xy + y^2$ ). Assume  $n \geq 3$ . The induction hypothesis is that  $(x+y)^n = \sum_{k=0}^n \binom{n}{k} x^k y^{n-k}$ . Then we

have

$$\begin{aligned}
(x+y)^{n+1} &= (x+y)(x+y)^n \\
&= (x+y) \left( \sum_{k=0}^n \binom{n}{k} x^k y^{n-k} \right) \\
&= \sum_{k=0}^n \binom{n}{k} x^{k+1} y^{n-k} + \sum_{k=0}^n \binom{n}{k} x^k y^{n-k+1} \\
&= x^{n+1} + \sum_{k=0}^{n-1} \binom{n}{k} x^{k+1} y^{n-k} + \sum_{k=1}^n \binom{n}{k} x^k y^{n-k+1} + y^{n+1} \\
&= x^{n+1} + \sum_{k=1}^n \binom{n}{k-1} x^k y^{n-k+1} + \sum_{k=1}^n \binom{n}{k} x^k y^{n-k+1} + y^{n+1} \\
&= x^{n+1} + \sum_{k=1}^n \left( \binom{n}{k-1} + \binom{n}{k} \right) x^k y^{n-k+1} + y^{n+1} \\
&= x^{n+1} + \sum_{k=1}^n \binom{n+1}{k} x^k y^{n-k+1} + y^{n+1} \\
&= \sum_{k=0}^{n+1} \binom{n+1}{k} x^k y^{n-k+1},
\end{aligned}$$

proving the theorem.  $\square$

**54 LEMMA** If  $a \in \mathbb{R}$ ,  $a \neq 1$  and  $n \in \mathbb{N} \setminus \{0\}$ , then

$$1 + a + a^2 + \cdots + a^{n-1} = \frac{1 - a^n}{1 - a}.$$

**Proof:** For, put  $S = 1 + a + a^2 + \cdots + a^{n-1}$ . Then  $aS = a + a^2 + \cdots + a^{n-1} + a^n$ . Thus

$$S - aS = (1 + a + a^2 + \cdots + a^{n-1}) - (a + a^2 + \cdots + a^{n-1} + a^n) = 1 - a^n,$$

and from  $(1 - a)S = S - aS = 1 - a^n$  we obtain the result.  $\square$

**55 THEOREM** Let  $n$  be a strictly positive integer. Then

$$y^n - x^n = (y - x)(y^{n-1} + y^{n-2}x + \cdots + yx^{n-2} + x^{n-1}).$$

**Proof:** By making the substitution  $a = \frac{x}{y}$  in Lemma 54 we see that

$$1 + \frac{x}{y} + \left(\frac{x}{y}\right)^2 + \cdots + \left(\frac{x}{y}\right)^{n-1} = \frac{1 - \left(\frac{x}{y}\right)^n}{1 - \frac{x}{y}}$$

we obtain

$$\left(1 - \frac{x}{y}\right) \left(1 + \frac{x}{y} + \left(\frac{x}{y}\right)^2 + \cdots + \left(\frac{x}{y}\right)^{n-1}\right) = 1 - \left(\frac{x}{y}\right)^n,$$

or equivalently,

$$\left(1 - \frac{x}{y}\right) \left(1 + \frac{x}{y} + \frac{x^2}{y^2} + \cdots + \frac{x^{n-1}}{y^{n-1}}\right) = 1 - \frac{x^n}{y^n}.$$

Multiplying by  $y^n$  both sides,

$$y \left(1 - \frac{x}{y}\right) y^{n-1} \left(1 + \frac{x}{y} + \frac{x^2}{y^2} + \cdots + \frac{x^{n-1}}{y^{n-1}}\right) = y^n \left(1 - \frac{x^n}{y^n}\right),$$

which is

$$y^n - x^n = (y - x)(y^{n-1} + y^{n-2}x + \cdots + yx^{n-2} + x^{n-1}),$$

yielding the result.  $\square$



**56 THEOREM**  $1 + 2 + \cdots + n = \frac{n(n+1)}{2}$ .

**First Proof:** Observe that

$$k^2 - (k-1)^2 = 2k - 1.$$

From this

$$\begin{array}{rcl} 1^2 - 0^2 & = & 2 \cdot 1 - 1 \\ 2^2 - 1^2 & = & 2 \cdot 2 - 1 \\ 3^2 - 2^2 & = & 2 \cdot 3 - 1 \\ \vdots & & \vdots \\ n^2 - (n-1)^2 & = & 2 \cdot n - 1 \end{array}$$

Adding both columns,

$$n^2 - 0^2 = 2(1 + 2 + 3 + \cdots + n) - n.$$

Solving for the sum,

$$1 + 2 + 3 + \cdots + n = n^2/2 + n/2 = \frac{n(n+1)}{2}.$$

□

**Second Proof:** We may utilise Gauss' trick: If

$$A_n = 1 + 2 + 3 + \cdots + n$$

then

$$A_n = n + (n-1) + \cdots + 1.$$

Adding these two quantities,

$$\begin{array}{rcl} A_n & = & 1 + 2 + \cdots + n \\ A_n & = & n + (n-1) + \cdots + 1 \\ \hline 2A_n & = & (n+1) + (n+1) + \cdots + (n+1) \\ & = & n(n+1), \end{array}$$

since there are  $n$  summands. This gives  $A_n = \frac{n(n+1)}{2}$ , that is,

$$1 + 2 + \cdots + n = \frac{n(n+1)}{2}.$$

Applying Gauss's trick to the general arithmetic sum

$$(a) + (a+d) + (a+2d) + \cdots + (a+(n-1)d)$$

we obtain

$$(a) + (a+d) + (a+2d) + \cdots + (a+(n-1)d) = \frac{n(2a+(n-1)d)}{2} \quad (1.5)$$

□

**57 THEOREM**  $1^2 + 2^2 + 3^2 + \cdots + n^2 = \frac{n(n+1)(2n+1)}{6}$ .

**Proof:** Observe that

$$k^3 - (k-1)^3 = 3k^2 - 3k + 1.$$

Hence

$$\begin{aligned} 1^3 - 0^3 &= 3 \cdot 1^2 - 3 \cdot 1 + 1 \\ 2^3 - 1^3 &= 3 \cdot 2^2 - 3 \cdot 2 + 1 \\ 3^3 - 2^3 &= 3 \cdot 3^2 - 3 \cdot 3 + 1 \\ \vdots &\quad \quad \quad \vdots \\ n^3 - (n-1)^3 &= 3 \cdot n^2 - 3 \cdot n + 1 \end{aligned}$$

Adding both columns,

$$n^3 - 0^3 = 3(1^2 + 2^2 + 3^2 + \dots + n^2) - 3(1 + 2 + 3 + \dots + n) + n.$$

From the preceding example  $1 + 2 + 3 + \dots + n = \frac{n^2}{2} + \frac{n}{2} = \frac{n(n+1)}{2}$  so

$$n^3 - 0^3 = 3(1^2 + 2^2 + 3^2 + \dots + n^2) - \frac{3}{2} \cdot n(n+1) + n.$$

Solving for the sum,

$$1^2 + 2^2 + 3^2 + \dots + n^2 = \frac{n^3}{3} + \frac{1}{2} \cdot n(n+1) - \frac{n}{3}.$$

After simplifying we obtain

$$1^2 + 2^2 + 3^2 + \dots + n^2 = \frac{n(n+1)(2n+1)}{6}.$$

□

### Homework

**Problem 1.5.1** Prove that for  $n \geq 1$ ,

$$2^n = \sum_{k=0}^n \binom{n}{k}; \quad 0 = \sum_{k=0}^n (-1)^k \binom{n}{k}, \quad 2^{n-1} = \sum_{\substack{0 \leq k \leq n \\ k \text{ even}}} \binom{n}{k} = \sum_{\substack{1 \leq k \leq n \\ k \text{ odd}}} \binom{n}{k}.$$

**Problem 1.5.2** Given that 1002004008016032 has a prime factor  $p > 250000$ , find it.

**Problem 1.5.3** Prove that  $(a + b + c)^2 = a^2 + b^2 + c^2 + 2ab + 2bc + 2ca$ .

**Problem 1.5.4** Let  $a, b, c$  be real numbers. Prove that

$$a^3 + b^3 + c^3 - 3abc = (a + b + c)(a^2 + b^2 + c^2 - ab - bc - ca).$$

**Problem 1.5.5** Prove that

$$\binom{n}{k} = \frac{n}{k} \binom{n-1}{k-1}.$$

**Problem 1.5.6** Prove that

$$\binom{n}{k} = \frac{n}{k} \cdot \frac{n-1}{k-1} \cdot \binom{n-2}{k-2}.$$

**Problem 1.5.7** Prove that

$$\sum_{k=1}^n k \binom{n}{k} p^k (1-p)^{n-k} = np.$$

**Problem 1.5.8** Prove that

$$\sum_{k=2}^n k(k-1) \binom{n}{k} p^k (1-p)^{n-k} = n(n-1)p^2.$$

**Problem 1.5.9** Demonstrate that

$$\sum_{k=0}^n (k-np)^2 \binom{n}{k} p^k (1-p)^{n-k} = np(1-p).$$

**Problem 1.5.10** Let  $x \in \mathbb{R} \setminus \{1\}$  and let  $n \in \mathbb{N} \setminus \{0\}$ . Prove that

$$\sum_{k=0}^n \frac{2^k}{x^{2^k} + 1} = \frac{1}{x-1} - \frac{2^{n+1}}{x^{2^{n+1}} + 1}.$$

**Problem 1.5.11** Consider the  $n^k$   $k$ -tuples  $(a_1, a_2, \dots, a_k)$  which can be formed by taking  $a_i \in \{1, 2, \dots, n\}$ , repetitions allowed. Demonstrate that

$$\sum_{a_i \in \{1, 2, \dots, n\}} \min(a_1, a_2, \dots, a_k) = 1^k + 2^k + \dots + n^k.$$

## 1.6 Order Axioms



**Vocabulary Alert!** We will call a number  $x$  positive if  $x \geq 0$  and strictly positive if  $x > 0$ . Similarly, we will call a number  $y$  negative if  $y \leq 0$  and strictly negative if  $y < 0$ . This usage differs from most Anglo-American books, who prefer such terms as non-negative and non-positive.

We assume  $\mathbb{R}$  endowed with a relation  $>$  which satisfies the following axioms.

**58 Axiom (Trichotomy Law)**  $\forall (x, y) \in \mathbb{R}^2$  exactly one of the following holds:

$$x > y, \quad x = y, \quad \text{or} \quad y > x.$$

**59 Axiom (Transitivity of Order)**  $\forall (x, y, z) \in \mathbb{R}^3$ ,

$$\text{if } x > y \text{ and } y > z \text{ then } x > z.$$

**60 Axiom (Preservation of Inequalities by Addition)**  $\forall (x, y, z) \in \mathbb{R}^3$ ,

$$\text{if } x > y \text{ then } x + z > y + z.$$

**61 Axiom (Preservation of Inequalities by Positive Factors)**  $\forall (x, y, z) \in \mathbb{R}^3$ ,

$$\text{if } x > y \text{ and } z > 0 \text{ then } xz > yz.$$



$x < y$  means that  $y > x$ .  $x \leq y$  means that either  $y > x$  or  $y = x$ , etc.

**62 THEOREM** The square of any real number is positive, that is,  $\forall a \in \mathbb{R}, \quad a^2 \geq 0$ . In fact, if  $a \neq 0$  then  $a^2 > 0$ .

**Proof:** If  $a = 0$ , then  $0^2 = 0$  and there is nothing to prove. Assume now that  $a \neq 0$ . By trichotomy, either  $a > 0$  or  $a < 0$ . Assume first that  $a > 0$ . Applying Axiom 61 with  $x = z = a$  and  $y = 0$  we have

$$aa > a0 \implies a^2 > 0,$$

so the theorem is proved if  $a > 0$ .

If  $a < 0$  then  $-a > 0$  and we apply the result just obtained:

$$-a > 0 \implies (-a)^2 > 0 \implies 1 \cdot a^2 > 0 \implies a^2 > 0,$$

so the result is true regardless the sign of  $a$ .  $\square$

Theorem 62 will prove to be extremely powerful and will be the basis for many of the classical inequalities that follow.

**63 THEOREM** If  $(x, y) \in \mathbb{R}^2$ ,

$$x > y \iff x - y > 0.$$

**Proof:** This is a direct consequence of Axiom 60 upon taking  $z = -y$ .  $\square$

**64 THEOREM** If  $(x, y, a, b) \in \mathbb{R}^4$ ,

$$x > y \text{ and } a \geq b \implies x + a > y + b.$$

**Proof:** We have

$$x > y \implies x + a > y + a, \quad y + a \geq y + b,$$

by Axiom 60 and so by Axiom 59  $x + a > y + b$ .  $\square$

**65 THEOREM** If  $(x, y, a, b) \in \mathbb{R}^4$ ,

$$x > y > 0 \quad \text{and} \quad a \geq b > 0 \implies xa > yb.$$

**Proof:** *Indeed*

$$x > y \implies xa > ya, \quad ya \geq yb,$$

by Axiom 61 and so by Axiom 59  $xa > yb$ .  $\square$

**66 THEOREM**  $1 > 0$ .

**Proof:** By definition of  $\mathbb{R}$  being a field  $0 \neq 1$ . Assume that  $1 < 0$  then  $1^2 > 0$  by Theorem 62. But  $1^2 = 1$  and so  $1 > 0$ , a contradiction to our original assumption.  $\square$

**67 THEOREM**  $x > 0 \implies -x < 0$  and  $x^{-1} > 0$ .

**Proof:** *Indeed,  $-1 < 0$  since  $-1 \neq 0$  and assuming  $-1 > 0$  would give  $0 = -1 + 1 > 1$ , which contradicts Theorem 66. Thus*

$$-x = -1 \cdot x < 0.$$

Similarly, assuming  $x^{-1} < 0$  would give  $1 = x^{-1}x < 0$ .  $\square$

**68 THEOREM**  $x > 1 \implies x^{-1} < 1$ .

**Proof:** Since  $x^{-1} \neq 1$ , assuming  $x^{-1} > 1$  would give  $1 = xx^{-1} > 1 \cdot 1 = 1$ , a contradiction.  $\square$

### 1.6.1 Absolute Value

**69 Definition (The Signum (Sign) Function)** Let  $x$  be a real number. We define  $\mathbf{signum}(x) = \begin{cases} -1 & \text{if } x < 0, \\ 0 & \text{if } x = 0, \\ +1 & \text{if } x > 0. \end{cases}$

**70 LEMMA** The signum function is multiplicative, that is, if  $(x, y) \in \mathbb{R}^2$  then  $\mathbf{signum}(x \cdot y) = \mathbf{signum}(x) \mathbf{signum}(y)$ .

**Proof:** *Immediate from the definition of signum.*  $\square$

**71 Definition (Absolute Value)** Let  $x \in \mathbb{R}$ . The *absolute value* of  $x$  is defined and denoted by

$$|x| = \mathbf{signum}(x) x.$$

**72 THEOREM** Let  $x \in \mathbb{R}$ . Then

$$1. \quad |x| = \begin{cases} -x & \text{if } x < 0, \\ x & \text{if } x \geq 0. \end{cases}$$

$$2. \quad |x| \geq 0,$$

$$3. \quad |x| = \max(x, -x),$$

$$4. \quad |-x| = |x|,$$

$$5. \quad -|x| \leq x \leq |x|.$$

$$6. \quad \sqrt{x^2} = |x|$$

$$7. \quad |x|^2 = |x^2| = x^2$$

$$8. \quad x = \mathbf{signum}(x) |x|$$

**Proof:** These are immediate from the definition of  $|x|$ .  $\square$

**73 THEOREM** ( $\forall (x, y) \in \mathbb{R}^2$ ),

$$|xy| = |x| |y|.$$

**Proof:** We have

$$|xy| = \text{signum}(xy) xy = (\text{signum}(x)x)(\text{signum}(y)y) = |x| |y|,$$

where we have used Lemma 70.  $\square$

**74 THEOREM** Let  $t \geq 0$ . Then

$$|x| \leq t \iff -t \leq x \leq t.$$

**Proof:** Either  $|x| = x$  or  $|x| = -x$ . If  $|x| = x$ ,

$$|x| \leq t \iff x \leq t \iff -t \leq 0 \leq x \leq t.$$

If  $|x| = -x$ ,

$$|x| \leq t \iff -x \leq t \iff -t \leq x \leq 0 \leq t.$$

$\square$

**75 THEOREM** If  $(x, y) \in \mathbb{R}^2$ ,  $\max(x, y) = \frac{x + y + |x - y|}{2}$  and  $\min(x, y) = \frac{x + y - |x - y|}{2}$ .

**Proof:** Observe that  $\max(x, y) + \min(x, y) = x + y$ , since one of these quantities must be the maximum and the other the minimum, or else, they are both equal.

Now, either  $|x - y| = x - y$ , and so  $x \geq y$ , meaning that  $\max(x, y) - \min(x, y) = x - y$ , or  $|x - y| = -(x - y) = y - x$ , which means that  $y \geq x$  and so  $\max(x, y) - \min(x, y) = y - x$ . In either case we get  $\max(x, y) - \min(x, y) = |x - y|$ . Solving now the system of equations

$$\begin{aligned} \max(x, y) + \min(x, y) &= x + y \\ \max(x, y) - \min(x, y) &= |x - y|, \end{aligned}$$

for  $\max(x, y)$  and  $\min(x, y)$  gives the result.  $\square$

## Homework

**Problem 1.6.1** Let  $x, y$  be real numbers. Then

$$0 \leq x < y \iff x^2 < y^2.$$

**Problem 1.6.2** Let  $t \geq 0$ . Prove that

$$|x| \geq t \iff (x \geq t) \text{ or } (x \leq -t).$$

**Problem 1.6.3** Let  $(x, y) \in \mathbb{R}^2$ . Prove that  $\max(x, y) = -\min(-x, -y)$ .

**Problem 1.6.4** Let  $x, y, z$  be real numbers. Prove that

$$\max(x, y, z) = x + y + z - \min(x, y) - \min(y, z) - \min(z, x) + \min(x, y, z).$$

**Problem 1.6.5** Let  $a < b$ . Demonstrate that

$$|x - a| < |x - b| \iff x < \frac{a + b}{2}.$$

## 1.7 Classical Inequalities

### 1.7.1 Triangle Inequality

**76 THEOREM (Triangle Inequality)** Let  $(a, b) \in \mathbb{R}^2$ . Then

$$|a + b| \leq |a| + |b|. \tag{1.6}$$

**Proof:** From 5 in Theorem 72, by addition,

$$-|a| \leq a \leq |a|$$

to

$$-|b| \leq b \leq |b|$$

we obtain

$$-(|a| + |b|) \leq a + b \leq (|a| + |b|),$$

whence the theorem follows by applying Theorem 74.  $\square$

By induction, we obtain the following generalisation to  $n$  terms.

**77 COROLLARY** Let  $x_1, x_2, \dots, x_n$  be real numbers. Then

$$|x_1 + x_2 + \dots + x_n| \leq |x_1| + |x_2| + \dots + |x_n|.$$

**Proof:** We apply Theorem 76  $n-1$  times

$$\begin{aligned} |x_1 + x_2 + \dots + x_n| &\leq |x_1| + |x_2 + \dots + x_{n-1} + x_n| \\ &\leq |x_1| + |x_2| + |x_3 + \dots + x_{n-1} + x_n| \\ &\vdots \\ &\leq |x_1| + |x_2| + \dots + |x_{n-1} + x_n| \\ &\leq |x_1| + |x_2| + \dots + |x_{n-1}| + |x_n|. \end{aligned}$$

$\square$

**78 COROLLARY** Let  $(a, b) \in \mathbb{R}^2$ . Then

$$\boxed{||a| - |b|| \leq |a - b|}. \quad (1.7)$$

**Proof:** We have

$$|a| = |a - b + b| \leq |a - b| + |b|,$$

giving

$$|a| - |b| \leq |a - b|.$$

Similarly,

$$|b| = |b - a + a| \leq |b - a| + |a| = |a - b| + |a|,$$

gives

$$|b| - |a| \leq |a - b| \implies -|a - b| \leq |a| - |b|.$$

Thus

$$-|a - b| \leq |a| - |b| \leq |a - b|,$$

and we now apply Theorem 74.  $\square$

**79 THEOREM** Let  $b_i > 0$  for  $1 \leq i \leq n$ . Then

$$\min\left(\frac{a_1}{b_1}, \frac{a_2}{b_2}, \dots, \frac{a_n}{b_n}\right) \leq \frac{a_1 + a_2 + \dots + a_n}{b_1 + b_2 + \dots + b_n} \leq \max\left(\frac{a_1}{b_1}, \frac{a_2}{b_2}, \dots, \frac{a_n}{b_n}\right).$$

**Proof:** For every  $k$ ,  $1 \leq k \leq n$ ,

$$\min\left(\frac{a_1}{b_1}, \frac{a_2}{b_2}, \dots, \frac{a_n}{b_n}\right) \leq \frac{a_k}{b_k} \leq \max\left(\frac{a_1}{b_1}, \frac{a_2}{b_2}, \dots, \frac{a_n}{b_n}\right) \implies b_k \min\left(\frac{a_1}{b_1}, \frac{a_2}{b_2}, \dots, \frac{a_n}{b_n}\right) \leq a_k \leq b_k \max\left(\frac{a_1}{b_1}, \frac{a_2}{b_2}, \dots, \frac{a_n}{b_n}\right).$$

Adding all these inequalities for  $1 \leq k \leq n$ ,

$$(b_1 + b_2 + \dots + b_n) \min\left(\frac{a_1}{b_1}, \frac{a_2}{b_2}, \dots, \frac{a_n}{b_n}\right) \leq a_1 + a_2 + \dots + a_n \leq (b_1 + b_2 + \dots + b_n) \max\left(\frac{a_1}{b_1}, \frac{a_2}{b_2}, \dots, \frac{a_n}{b_n}\right),$$

from where the result is obtained.  $\square$

## 1.7.2 Bernoulli's Inequality

**80 THEOREM** If  $0 \leq a < b$ ,  $n \geq 1 \in \mathbb{N}$

$$na^{n-1} < \frac{b^n - a^n}{b - a} < nb^{n-1}.$$

**Proof:** By Theorem 55,

$$\begin{aligned} \frac{b^n - a^n}{b - a} &= b^{n-1} + b^{n-2}a + b^{n-3}a^2 + \cdots + b^2a^{n-3} + ba^{n-2} + a^{n-1} \\ &< b^{n-1} + b^{n-1} + \cdots + b^{n-1} + b^{n-1} \\ &= nb^{n-1}, \end{aligned}$$

from where the dextral inequality follows. The sinistral inequality can be established similarly.  $\square$

**81 THEOREM (Bernoulli's Inequality)** If  $x > -1$ ,  $x \neq 0$ , and if  $n \in \mathbb{N} \setminus \{0\}$  then

$$(1 + x)^n > 1 + nx.$$

**Proof:** Set  $b = 1 + x$ ,  $a = 1$  in Theorem 80 and use the sinistral inequality.  $\square$



If  $x > 0$  then Bernoulli's Inequality is an easy consequence of the Binomial Theorem, as

$$(1 + x)^n = 1 + \binom{n}{1}x + \binom{n}{2}x^2 + \cdots > 1 + \binom{n}{1}x = 1 + nx.$$

## 1.7.3 Rearrangement Inequality

**82 Definition** Given a set of real numbers  $\{x_1, x_2, \dots, x_n\}$  denote by

$$\check{x}_1 \geq \check{x}_2 \geq \cdots \geq \check{x}_n$$

the decreasing rearrangement of the  $x_i$  and denote by

$$\hat{x}_1 \leq \hat{x}_2 \leq \cdots \leq \hat{x}_n$$

the increasing rearrangement of the  $x_i$ .


**83 Definition** Given two sequences of real numbers  $\{x_1, x_2, \dots, x_n\}$  and  $\{y_1, y_2, \dots, y_n\}$  of the same length  $n$ , we say that they are *similarly sorted* if they are both increasing or both decreasing, and *differently sorted* if one is increasing and the other decreasing.

**84 Example** The sequences  $1 \leq 2 \leq \cdots \leq n$  and  $1^2 \leq 2^2 \leq \cdots \leq n^2$  are similarly sorted, and the sequences  $\frac{1}{1^2} \geq \frac{1}{2^2} \geq \cdots \geq \frac{1}{n^2}$  and  $1^3 \leq 2^3 \leq \cdots \leq n^3$  are differently sorted.

**85 THEOREM (Rearrangement Inequality)** Given sets of real numbers  $\{a_1, a_2, \dots, a_n\}$  and  $\{b_1, b_2, \dots, b_n\}$  we have

$$\sum_{1 \leq k \leq n} \check{a}_k \hat{b}_k \leq \sum_{1 \leq k \leq n} a_k b_k \leq \sum_{1 \leq k \leq n} \hat{a}_k \hat{b}_k.$$

Thus the sum  $\sum_{1 \leq k \leq n} a_k b_k$  is minimised when the sequences are differently sorted, and maximised when the sequences are similarly sorted.

 Observe that

$$\sum_{1 \leq k \leq n} \check{a}_k \hat{b}_k = \sum_{1 \leq k \leq n} \hat{a}_k \check{b}_k \quad \text{and} \quad \sum_{1 \leq k \leq n} \hat{a}_k \hat{b}_k = \sum_{1 \leq k \leq n} \check{a}_k \check{b}_k.$$

**Proof:** Let  $\{\sigma(1), \sigma(2), \dots, \sigma(n)\}$  be a reordering of  $\{1, 2, \dots, n\}$ . If there are two sub-indices  $i, j$ , such that the sequences pull in opposite directions, say,  $a_i > a_j$  and  $b_{\sigma(i)} < b_{\sigma(j)}$ , then consider the sums

$$\begin{aligned} S &= a_1 b_{\sigma(1)} + a_2 b_{\sigma(2)} + \dots + a_i b_{\sigma(i)} + \dots + a_j b_{\sigma(j)} + \dots + a_n b_{\sigma(n)} \\ S' &= a_1 b_{\sigma(1)} + a_2 b_{\sigma(2)} + \dots + a_i b_{\sigma(j)} + \dots + a_j b_{\sigma(i)} + \dots + a_n b_{\sigma(n)} \end{aligned}$$

Then

$$S' - S = (a_i - a_j)(b_{\sigma(j)} - b_{\sigma(i)}) > 0.$$

This last inequality shows that the closer the  $a$ 's and the  $b$ 's are to pulling in the same direction the larger the sum becomes. This proves the result.  $\square$

### 1.7.4 Arithmetic Mean-Geometric Mean Inequality

**86 THEOREM (Arithmetic Mean-Geometric Mean Inequality)** Let  $a_1, \dots, a_n$  be positive real numbers. Then their geometric mean is at most their arithmetic mean, that is,

$$\sqrt[n]{a_1 \cdots a_n} \leq \frac{a_1 + \dots + a_n}{n},$$

with equality if and only if  $a_1 = \dots = a_n$ .

We will provide multiple proofs of this important inequality. Some other proofs will be found in latter chapters.

**First Proof:** Our first proof uses the Rearrangement Inequality (Theorem 85) in a rather clever way. We may assume that the  $a_k$  are strictly positive. Put

$$x_1 = \frac{a_1}{(a_1 a_2 \cdots a_n)^{1/n}}, \quad x_2 = \frac{a_1 a_2}{(a_1 a_2 \cdots a_n)^{2/n}}, \quad \dots, \quad x_n = \frac{a_1 a_2 \cdots a_n}{(a_1 a_2 \cdots a_n)^{n/n}} = 1,$$

and

$$y_1 = \frac{1}{x_1}, \quad y_2 = \frac{1}{x_2}, \quad \dots, \quad y_n = \frac{1}{x_n} = 1.$$

Observe that for  $2 \leq k \leq n$ ,

$$x_k y_{k-1} = \frac{a_1 a_2 \cdots a_k}{(a_1 a_2 \cdots a_n)^{k/n}} \cdot \frac{(a_1 a_2 \cdots a_n)^{(k-1)/n}}{a_1 a_2 \cdots a_{k-1}} = \frac{a_k}{(a_1 a_2 \cdots a_n)^{1/n}}.$$

The  $x_k$  and  $y_k$  are differently sorted, so by virtue of the Rearrangement Inequality we gather

$$\begin{aligned} 1 + 1 + \dots + 1 &= x_1 y_1 + x_2 y_2 + \dots + x_n y_n \\ &\leq x_1 y_n + x_2 y_1 + \dots + x_n y_{n-1} \\ &= \frac{a_1}{(a_1 a_2 \cdots a_n)^{1/n}} + \frac{a_2}{(a_1 a_2 \cdots a_n)^{1/n}} + \dots + \frac{a_n}{(a_1 a_2 \cdots a_n)^{1/n}}, \end{aligned}$$

or

$$n \leq \frac{a_1 + a_2 + \dots + a_n}{(a_1 a_2 \cdots a_n)^{1/n}},$$

from where we obtain the result.  $\square$

**Second Proof:** This second proof is a clever induction argument due to Cauchy. It proves the inequality first for powers of 2 and then interpolates for numbers between consecutive powers of 2.

Since the square of a real number is always positive, we have, for positive real numbers  $a, b$

$$(\sqrt{a} - \sqrt{b})^2 \geq 0 \implies \sqrt{ab} \leq \frac{a+b}{2},$$



proving the inequality for  $k = 2$ . Observe that equality happens if and only if  $\mathbf{a} = \mathbf{b}$ . Assume now that the inequality is valid for  $k = 2^{n-1} > 2$ . This means that for any positive real numbers  $x_1, x_2, \dots, x_{2^{n-1}}$  we have

$$(x_1 x_2 \cdots x_{2^{n-1}})^{1/2^{n-1}} \leq \frac{x_1 + x_2 + \cdots + x_{2^{n-1}}}{2^{n-1}}. \quad (1.8)$$

Let us prove the inequality for  $2k = 2^n$ . Consider any any positive real numbers  $y_1, y_2, \dots, y_{2^n}$ . Notice that there are  $2^n - 2^{n-1} = 2^{n-1}(2 - 1) = 2^{n-1}$  integers in the interval  $[2^{n-1} + 1; 2^n]$ . We have

$$\begin{aligned} (y_1 y_2 \cdots y_{2^n})^{1/2^n} &= \sqrt{(y_1 y_2 \cdots y_{2^{n-1}})^{1/2^{n-1}} (y_{2^{n-1}+1} \cdots y_{2^n})^{1/2^{n-1}}} \\ &\leq \frac{(y_1 y_2 \cdots y_{2^{n-1}})^{1/2^{n-1}} + (y_{2^{n-1}+1} \cdots y_{2^n})^{1/2^{n-1}}}{2} \\ &\leq \frac{\frac{y_1 + y_2 + \cdots + y_{2^{n-1}}}{2^{n-1}} + \frac{2}{2^{n-1}} \frac{y_{2^{n-1}+1} + \cdots + y_{2^n}}{2^{n-1}}}{2} \\ &= \frac{y_1 + \cdots + y_{2^n}}{2^n}, \end{aligned}$$

where the first inequality follows by the Case  $n = 2$  and the second by the induction hypothesis (1.8). The theorem is thus proved for powers of 2.

Assume now that  $2^{n-1} < k < 2^n$ , and consider the  $k$  positive real numbers  $a_1, a_2, \dots, a_k$ . The trick is to pad this collection of real numbers up to the next highest power of 2, the added real numbers being the average of the existing ones. Hence consider the  $2^n$  real numbers

$$a_1, a_2, \dots, a_k, a_{k+1}, \dots, a_{2^n}$$

with  $a_{k+1} = \dots = a_{2^n} = \frac{a_1 + a_2 + \cdots + a_k}{k}$ . Since we have already proved the theorem for  $2^n$  we have

$$\left( a_1 a_2 \cdots a_k \left( \frac{a_1 + a_2 + \cdots + a_k}{k} \right)^{2^n - k} \right)^{1/2^n} \leq \frac{a_1 + a_2 + \cdots + a_k + (2^n - k) \left( \frac{a_1 + a_2 + \cdots + a_k}{k} \right)}{2^n},$$

whence

$$(a_1 a_2 \cdots a_k)^{1/2^n} \left( \frac{a_1 + a_2 + \cdots + a_k}{k} \right)^{1 - k/2^n} \leq \frac{k \frac{a_1 + a_2 + \cdots + a_k}{k} + (2^n - k) \left( \frac{a_1 + a_2 + \cdots + a_k}{k} \right)}{2^n},$$

which implies

$$(a_1 a_2 \cdots a_k)^{1/2^n} \left( \frac{a_1 + a_2 + \cdots + a_k}{k} \right)^{1 - k/2^n} \leq \left( \frac{a_1 + a_2 + \cdots + a_k}{k} \right),$$

Solving for  $\frac{a_1 + a_2 + \cdots + a_k}{k}$  gives the desired inequality.  $\square$

**Third Proof:** As in the second proof, the Case  $k = 2$  is easily established. Put

$$A_k = \frac{a_1 + a_2 + \cdots + a_k}{k}, \quad G_k = (a_1 a_2 \cdots a_k)^{1/k}.$$

Observe that

$$a_{k+1} = (k+1)A_{k+1} - kA_k.$$

The inductive hypothesis is that  $A_k \geq G_k$  and we must shew that  $A_{k+1} \geq G_{k+1}$ . Put

$$A = \frac{a_{k+1} + (k-1)A_{k+1}}{k}, \quad G = \left( a_{k+1} A_{k+1}^{k-1} \right)^{1/k}.$$

By the inductive hypothesis  $A \geq G$ . Now,

$$\frac{A + A_k}{2} = \frac{(k+1)A_{k+1} - kA_k + (k-1)A_{k+1} + A_k}{2} = A_{k+1}.$$

Hence

$$\begin{aligned} A_{k+1} &= \frac{A + A_k}{2} \\ &\geq (AA_k)^{1/2} \\ &\geq (GG_k)^{1/2} \\ &= \left(G_{k+1}^{k+1} A_{k+1}^{k-1}\right)^{1/2k} \end{aligned}$$

We have established that

$$A_{k+1} \geq \left(G_{k+1}^{k+1} A_{k+1}^{k-1}\right)^{1/2k} \implies A_{k+1} \geq G_{k+1},$$

completing the induction.  $\square$

**Fourth Proof:** We will make a series of substitutions that preserve the sum

$$a_1 + a_2 + \dots + a_n$$

while strictly increasing the product

$$a_1 a_2 \dots a_n.$$

At the end, the  $a_i$  will all be equal and the arithmetic mean  $A$  of the numbers will be equal to their geometric mean  $G$ . If the  $a_i$  where all  $> A$  then  $\frac{a_1 + a_2 + \dots + a_n}{n} > \frac{nA}{n} = A$ , impossible. Similarly, the  $a_i$  cannot be all  $< A$ . Hence there must exist two indices say  $i, j$ , such that  $a_i < A < a_j$ . Put  $a'_i = A$ ,  $a'_j = a_i + a_j - A$ . Observe that  $a_i + a_j = a'_i + a'_j$ , so replacing the original  $a$ 's with the primed  $a$ 's does not alter the arithmetic mean. On the other hand,

$$a'_i a'_j = A(a_i + a_j - A) = a_i a_j + (a_j - A)(A - a_i) > a_i a_j$$

since  $a_j - A > 0$  and  $A - a_i > 0$ .

This change has replaced one of the  $a$ 's by a quantity equal to the arithmetic mean, has not changed the arithmetic mean, and made the geometric mean larger. Since there at most  $n$   $a$ 's to be replaced, the procedure must eventually terminate when all the  $a$ 's are equal (to their arithmetic mean). Strict inequality then holds when at least two of the  $a$ 's are unequal.  $\square$

### 1.7.5 Cauchy-Bunyakovsky-Schwarz Inequality

**87 THEOREM (Cauchy-Bunyakovsky-Schwarz Inequality)** Let  $x_k, y_k$  be real numbers,  $1 \leq k \leq n$ . Then

$$\left| \sum_{k=1}^n x_k y_k \right| \leq \left( \sum_{k=1}^n x_k^2 \right)^{1/2} \left( \sum_{k=1}^n y_k^2 \right)^{1/2},$$

with equality if and only if

$$(a_1, a_2, \dots, a_n) = t(b_1, b_2, \dots, b_n)$$

for some real constant  $t$ .

**First Proof:** The inequality follows at once from Lagrange's Identity

$$\left( \sum_{k=1}^n x_k y_k \right)^2 = \left( \sum_{k=1}^n x_k^2 \right) \left( \sum_{k=1}^n y_k^2 \right) - \sum_{1 \leq k < j \leq n} (x_k y_j - x_j y_k)^2$$

(Theorem 50), since  $\sum_{1 \leq k < j \leq n} (x_k y_j - x_j y_k)^2 \geq 0$ .  $\square$

**Second Proof:** Put  $a = \sum_{k=1}^n x_k^2$ ,  $b = \sum_{k=1}^n x_k y_k$ , and  $c = \sum_{k=1}^n y_k^2$ . Consider the quadratic polynomial

$$at^2 + bt + c = t^2 \sum_{k=1}^n x_k^2 - 2t \sum_{k=1}^n x_k y_k + \sum_{k=1}^n y_k^2 = \sum_{k=1}^n (tx_k - y_k)^2 \geq 0,$$

where the inequality follows because a sum of squares of real numbers is being summed. Thus this quadratic polynomial is positive for all real  $t$ , so it must have complex roots. Its discriminant  $b^2 - 4ac$  must be negative, from where we gather

$$4 \left( \sum_{k=1}^n x_k y_k \right)^2 \leq 4 \left( \sum_{k=1}^n x_k^2 \right) \left( \sum_{k=1}^n y_k^2 \right),$$

which gives the inequality  $\square$

For our third proof of the CBS Inequality we need the following lemma.

**88 LEMMA** For  $(a, b, x, y) \in \mathbb{R}^4$  with  $x > 0$  and  $y > 0$  the following inequality holds:

$$\frac{a^2}{x} + \frac{b^2}{y} \geq \frac{(a+b)^2}{x+y}.$$

Equality holds if and only if  $\frac{a}{x} = \frac{b}{y}$ .

**Proof:** Since the square of a real number is always positive, we have

$$\begin{aligned} (ay - bx)^2 \geq 0 &\implies a^2 y^2 - 2abxy + b^2 x^2 \geq 0 \\ &\implies a^2 y(x+y) + b^2 x(x+y) \geq (a+b)^2 xy \\ &\implies \frac{a^2}{x} + \frac{b^2}{y} \geq \frac{(a+b)^2}{x+y}. \end{aligned}$$

Equality holds if and only if the first inequality is 0.  $\square$



Iterating the result on Lemma 88,

$$\frac{a_1^2}{b_1} + \frac{a_2^2}{b_2} + \dots + \frac{a_n^2}{b_n} \geq \frac{(a_1 + a_2 + \dots + a_n)^2}{b_1 + b_2 + \dots + b_n},$$

with equality if and only if  $\frac{a_1}{b_1} = \frac{a_2}{b_2} = \dots = \frac{a_n}{b_n}$ .

**Third Proof:** By the preceding remark, we have

$$\begin{aligned} x_1^2 + x_2^2 + \dots + x_n^2 &= \frac{x_1^2 y_1^2}{y_1^2} + \frac{x_2^2 y_2^2}{y_2^2} + \dots + \frac{x_n^2 y_n^2}{y_n^2} \\ &\geq \frac{(x_1 y_1 + x_2 y_2 + \dots + x_n y_n)^2}{y_1^2 + y_2^2 + \dots + y_n^2}, \end{aligned}$$

and upon rearranging, CBS is once again obtained.  $\square$

### 1.7.6 Minkowski's Inequality

**89 THEOREM (Minkowski's Inequality)** Let  $x_k, y_k$  be any real numbers. Then

$$\left( \sum_{k=1}^n (x_k + y_k)^2 \right)^{1/2} \leq \left( \sum_{k=1}^n x_k^2 \right)^{1/2} + \left( \sum_{k=1}^n y_k^2 \right)^{1/2}.$$

**Proof:** We have

$$\begin{aligned} \sum_{k=1}^n (x_k + y_k)^2 &= \sum_{k=1}^n x_k^2 + 2 \sum_{k=1}^n x_k y_k + \sum_{k=1}^n y_k^2 \\ &\leq \sum_{k=1}^n x_k^2 + 2 \left( \sum_{k=1}^n x_k^2 \right)^{1/2} \left( \sum_{k=1}^n y_k^2 \right)^{1/2} + \sum_{k=1}^n y_k^2 \\ &= \left( \left( \sum_{k=1}^n x_k^2 \right)^{1/2} + \left( \sum_{k=1}^n y_k^2 \right)^{1/2} \right)^2, \end{aligned}$$

where the inequality follows from the CBS Inequality.  $\square$

## Homework

**Problem 1.7.1** Let  $(a, b, c, d) \in \mathbb{R}^4$ . Prove that

$$||a - c| - |b - c|| \leq |a - b| \leq |a - c| + |b - c|.$$

**Problem 1.7.2** Let  $(x_1, x_2, \dots, x_n) \in \mathbb{R}^n$  be such that

$$x_1^2 + x_2^2 + \dots + x_n^2 = x_1^3 + x_2^3 + \dots + x_n^3 = x_1^4 + x_2^4 + \dots + x_n^4.$$

Prove that  $x_k \in \{0, 1\}$ .

**Problem 1.7.3** Let  $n \geq 2$  an integer. Let  $(x_1, x_2, \dots, x_n) \in \mathbb{R}^n$  be such that

$$x_1^2 + x_2^2 + \dots + x_n^2 = x_1 x_2 + x_2 x_3 + \dots + x_{n-1} x_n + x_n x_1.$$

Prove that  $x_1 = x_2 = \dots = x_n$ .

**Problem 1.7.4** If  $b > 0$  and  $B > 0$  prove that

$$\frac{a}{b} < \frac{A}{B} \implies \frac{a}{b} < \frac{a+A}{b+B} < \frac{A}{B}.$$

Further, if  $p$  and  $q$  are positive integers such that

$$\frac{7}{10} < \frac{p}{q} < \frac{11}{15},$$

what is the least value of  $q$ ?

**Problem 1.7.5** Prove that if  $r \geq s \geq t$  then

$$r^2 - s^2 + t^2 \geq (r - s + t)^2.$$

**Problem 1.7.6** Assume that  $a_k, b_k, c_k, k = 1, \dots, n$ , are positive real numbers. Shew that

$$\left( \sum_{k=1}^n a_k b_k c_k \right)^4 \leq \left( \sum_{k=1}^n a_k^4 \right) \left( \sum_{k=1}^n b_k^4 \right) \left( \sum_{k=1}^n c_k^2 \right)^2.$$

**Problem 1.7.7** Prove that for integer  $n > 1$ ,

$$n! < \left( \frac{n+1}{2} \right)^n.$$

**Problem 1.7.8** Prove that for integer  $n > 2$ ,

$$n^{n/2} < n!.$$

**Problem 1.7.9** Prove that for all integers  $n \geq 0$  the inequality  $n(n-1) < 2^{n+1}$  is verified.

**Problem 1.7.10** Prove that  $\forall (a, b, c) \in \mathbb{R}^3$ ,

$$a^2 + b^2 + c^2 \geq ab + bc + ca.$$

**Problem 1.7.11** Prove that  $\forall (a, b, c) \in \mathbb{R}^3$ , with  $a \geq 0, b \geq 0, c \geq 0$ , the following inequalities hold:

$$a^3 + b^3 + c^3 \geq \max(a^2 b + b^2 c + c^2 a, a^2 c + b^2 a + c^2 b),$$

$$a^3 + b^3 + c^3 \geq 3abc,$$

$$a^3 + b^3 + c^3 \geq \frac{1}{2} (a^2(b+c) + b^2(c+a) + c^2(a+b)).$$

**Problem 1.7.12 (Chebyshev's Inequality)** Given sets of real numbers  $\{a_1, a_2, \dots, a_n\}$  and  $\{b_1, b_2, \dots, b_n\}$  prove that

$$\frac{1}{n} \sum_{1 \leq k \leq n} \check{a}_k \hat{b}_k \leq \left( \frac{1}{n} \sum_{1 \leq k \leq n} a_k \right) \left( \frac{1}{n} \sum_{1 \leq k \leq n} b_k \right) \leq \frac{1}{n} \sum_{1 \leq k \leq n} \hat{a}_k \hat{b}_k.$$

**Problem 1.7.13** If  $x > 0$ , from

$$\sqrt{x+1} - \sqrt{x} = \frac{1}{\sqrt{x+1} + \sqrt{x}},$$

prove that

$$\frac{1}{2\sqrt{x+1}} < \sqrt{x+1} - \sqrt{x} < \frac{1}{2\sqrt{x}}.$$

Use this to prove that if  $n > 1$  is a positive integer, then

$$2\sqrt{n+1} - 2 < 1 + \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{3}} + \dots + \frac{1}{\sqrt{n}} < 2\sqrt{n} - 1$$

**Problem 1.7.14** If  $0 < a \leq b$ , shew that

$$\frac{1}{8} \cdot \frac{(b-a)^2}{b} \leq \frac{a+b}{2} - \sqrt{ab} \leq \frac{1}{8} \cdot \frac{(b-a)^2}{a}$$

**Problem 1.7.15** Shew that

$$\frac{1}{2} \cdot \frac{3}{4} \cdot \frac{5}{6} \dots \frac{9999}{10000} < \frac{1}{100}.$$

**Problem 1.7.16** Prove that for all  $x > 0$ ,

$$\sum_{k=1}^n \frac{1}{(x+k)^2} < \frac{1}{x} - \frac{1}{x+n}.$$

**Problem 1.7.17** Let  $x_i \in \mathbb{R}$  such that  $\sum_{i=1}^n |x_i| = 1$  and  $\sum_{i=1}^n x_i = 0$ .

Prove that

$$\left| \sum_{i=1}^n \frac{x_i}{i} \right| \leq \frac{1}{2} \left( 1 - \frac{1}{n} \right).$$

**Problem 1.7.18** Let  $n$  be a strictly positive integer. Let  $x_i \geq 0$ . Prove that

$$\prod_{k=1}^n (1 + x_k) \geq 1 + \sum_{k=1}^n x_k.$$

When does equality hold?

**Problem 1.7.19 (Nesbitt's Inequality)** Let  $a, b, c$  be strictly positive real numbers. Then

$$\frac{a}{b+c} + \frac{b}{c+a} + \frac{c}{a+b} \geq \frac{3}{2}.$$

**Problem 1.7.20** Let  $a > 0$ . Use mathematical induction to prove that

$$\sqrt{a + \sqrt{a + \sqrt{a + \cdots + \sqrt{a}}}} < \frac{1 + \sqrt{4a + 1}}{2},$$

where the left member contains an arbitrary number of radicals.

**Problem 1.7.21** Let  $a, b, c$  be positive real numbers. Prove that

$$(a + b)(b + c)(c + a) \geq 8abc.$$

**Problem 1.7.22 (IMO, 1978)** Let  $a_k$  be a sequence of pairwise distinct positive integers. Prove that

$$\sum_{k=1}^n \frac{a_k}{k^2} \geq \sum_{k=1}^n \frac{1}{k}.$$

**Problem 1.7.23 (Harmonic Mean-Geometric Mean Inequality)**

Let  $x_i > 0$  for  $1 \leq i \leq n$ . Then

$$\frac{n}{\frac{1}{x_1} + \frac{1}{x_2} + \cdots + \frac{1}{x_n}} \leq (x_1 x_2 \cdots x_n)^{1/n},$$

with equality iff  $x_1 = x_2 = \cdots = x_n$ .

**Problem 1.7.24 (Arithmetic Mean-Quadratic Mean Inequality)**

Let  $x_i \geq 0$  for  $1 \leq i \leq n$ . Then

$$\frac{x_1 + x_2 + \cdots + x_n}{n} \leq \left( \frac{x_1^2 + x_2^2 + \cdots + x_n^2}{n} \right)^{1/2},$$

with equality iff  $x_1 = x_2 = \cdots = x_n$ .

**Problem 1.7.25** Given a set of real numbers  $\{a_1, a_2, \dots, a_n\}$  prove that there is an index  $m \in \{0, 1, \dots, n\}$  such that

$$\left| \sum_{1 \leq k \leq m} a_k - \sum_{m < k \leq n} a_k \right| \leq \max_{1 \leq k \leq n} |a_k|.$$

If  $m = 0$  the first sum is to be taken as 0 and if  $m = n$  the second one will be taken as 0.

**Problem 1.7.26** Give a purely geometric proof of Minkowski's inequality for  $n = 2$ . That is, prove that if  $(a, b), (c, d) \in \mathbb{R}^2$ , then

$$\sqrt{(a+c)^2 + (b+d)^2} \leq \sqrt{a^2 + b^2} + \sqrt{c^2 + d^2}.$$

Equality occurs if and only if  $ad = bc$ .

**Problem 1.7.27** Let  $x_k \in [0; 1]$  for  $1 \leq k \leq n$ . Demonstrate that

$$\min \left( \prod_{k=1}^n x_k, \prod_{k=1}^n (1 - x_k) \right) \leq \frac{1}{2^n}.$$

**Problem 1.7.28** If  $n > 0$  is an integer and if  $a_k > 0$ ,  $1 \leq k \leq n$  are real numbers, demonstrate that

$$\left( \sum_{k=1}^n \frac{a_k}{k} \right)^2 \leq \sum_{j=1}^n \sum_{k=1}^n \frac{a_j a_k}{j+k-1}.$$

**Problem 1.7.29** Let  $n$  be a strictly positive integer, let  $a_k \geq 0$ ,  $1 \leq k \leq n$  be real numbers such that  $a_1 \geq a_2 \geq \cdots \geq a_n$ , and let  $b_k$ ,  $1 \leq k \leq n$  be real numbers. Assume that for all indices  $k \in \{1, 2, \dots, n\}$ ,

$$\sum_{i=1}^k a_i \leq \sum_{i=1}^k b_i.$$

Prove that

$$\sum_{i=1}^n a_i^2 \leq \sum_{i=1}^n b_i^2$$

**Problem 1.7.30** Let  $n \geq 2$  an integer and let  $a_k$ ,  $1 \leq k \leq n$  be real numbers such that  $a_1 \leq a_2 \leq \cdots \leq a_n$ . Prove that there is an index  $k \in \{1, 2, \dots, n\}$  such that

$$(a_{k+1} - a_k)^2 \leq \frac{12}{n(n^2 - 1)} (a_1^2 + a_2^2 + \cdots + a_n^2).$$

**Problem 1.7.31 (AIME 1991)** Let  $P = \{a_1, a_2, \dots, a_n\}$  be a collection of points with

$$0 < a_1 < a_2 < \cdots < a_n < 17.$$

Consider


$$S_n = \min_P \sum_{k=1}^n \sqrt{(2k-1)^2 + a_k^2},$$

where the minimum runs over all such partitions  $P$ . Show that exactly one of  $S_2, S_3, \dots, S_n, \dots$  is an integer, and find which one it is.

## 1.8 Completeness Axiom

**Why bother?** We saw that both  $\mathbb{Q}$  and  $\mathbb{R}$  are fields, and hence they both satisfy the same arithmetical axioms. Why the need then for  $\mathbb{R}$ ? In this section we will study a property of  $\mathbb{R}$  that is not shared with  $\mathbb{Q}$ , that of completeness. It essentially means that there are no 'holes' on the real line.

**90 Definition** A number  $u$  is an *upper bound* for a set of numbers  $A \subseteq \mathbb{R}$  if for all  $a \in A$  we have  $a \leq u$ . The smallest such upper bound is called the *supremum* or *least upper bound* of the set  $A$ , and is denoted by  $\sup A$ . If  $\sup A \in A$  then we say that  $A$  has a *maximum* and we denote it by  $\max A (= \sup A)$ . Similarly, a number  $l$  is a *lower bound* for a set of numbers  $B \subseteq \mathbb{R}$  if for all  $b \in B$  we have  $l \leq b$ . The largest such lower bound is called the *infimum* or *greatest lower bound* of the set  $B$ , and is denoted by  $\inf B$ . If  $\inf B \in B$  then we say that  $B$  has a *minimum* and we denote it by  $\inf B (= \min B)$ .

 We define  $\inf(\mathbb{R}) = -\infty$ ,  $\sup(\mathbb{R}) = +\infty$ ,  $\inf(\emptyset) = +\infty$  and  $\sup(\emptyset) = -\infty$ .

**91 Definition** A set of numbers  $A$  is said to be *complete* if every non-empty subset of  $A$  which is bounded above has a supremum lying in  $A$ .

**92 Axiom (Completeness of  $\mathbb{R}$ )** Any non-empty set of real numbers which is bounded above has a supremum. Any non-empty set of real numbers which is bounded below has an infimum.

**93 THEOREM (Approximation Property of the Supremum and Infimum)** Let  $A \neq \emptyset$  be a set of real numbers possessing a supremum  $\sup A$ . Then

$$\forall \epsilon > 0 \quad \exists a \in A \quad \text{such that} \quad \sup A - \epsilon \leq a.$$


Let  $B \neq \emptyset$  be a set of real numbers possessing an infimum  $\inf B$ . Then

$$\forall \epsilon > 0 \quad \exists b \in B \quad \text{such that} \quad \inf B + \epsilon \geq b.$$

**Proof:** If  $\forall a \in A$ ,  $\sup A - \epsilon > a$  then  $\sup A - \epsilon$  would be an upper bound smaller than the least upper bound, a contradiction to the definition of  $\sup A$ . Hence there must be a rogue  $a \in A$  such that  $\sup A - \epsilon \leq a$ .

If  $\forall b \in B$ ,  $\inf B + \epsilon < b$  then  $\inf B + \epsilon$  would be a lower bound greater than the greatest lower bound, a contradiction to the definition of  $\inf B$ . Hence there must be a rogue  $b \in B$  such that  $\inf B + \epsilon \geq b$ .

□

 The above result should be intuitively clear.  $\sup A$  sits on the fence, just to the right of  $A$ , so that going just a bit to the left should put  $\sup A - \epsilon$  within  $A$ , etc.

**94 THEOREM (Monotonicity Property of the Supremum and Infimum)** Let  $\emptyset \subsetneq A \subseteq B \subseteq \mathbb{R}$  and suppose that both  $A$  and  $B$  have a supremum and an infimum. Then  $\sup A \leq \sup B$  and  $\inf B \leq \inf A$ .

**Proof:** Assume  $B$  is bounded above with supremum  $\sup B$ . Suppose  $x \in A$ . Then  $x \in B$  and so  $x \leq \sup B$ . Thus  $\sup B$  is an upper bound for the elements of  $A$ , and so by definition,  $\sup A \leq \sup B$ .

Assume  $B$  is bounded below with infimum  $\inf B$ . Suppose  $x \in A$ . Then  $x \in B$  and so  $x \geq \inf B$ . Thus  $\inf B$  is a lower bound for the elements of  $A$  and so by definition,  $\inf A \geq \inf B$ . □

**95 LEMMA** Let  $a, b$  be real numbers and assume that for all numbers  $\epsilon > 0$  the following inequality holds:

$$a < b + \epsilon.$$

Then  $a \leq b$ .

**Proof:** Assume contrariwise that  $a > b$ . Hence  $\frac{a-b}{2} > 0$ . Since the inequality  $a < b + \epsilon$  holds for every  $\epsilon > 0$  in particular it holds for  $\epsilon = \frac{a-b}{2}$ . This implies that

$$a < b + \frac{a-b}{2} \quad \text{or} \quad a < b.$$

Thus starting with the assumption that  $a > b$  we reach the incompatible conclusion that  $a < b$ . The original assumption must be wrong. We therefore conclude that  $a \leq b$ . □

**96 THEOREM (Additive Property of the Supremum)** Let  $\emptyset \subsetneq A \subseteq \mathbb{R}$ , and  $B \subseteq \mathbb{R}$ . Put

$$A + B = \{x + y : (x, y) \in A \times B\}$$

and suppose that both  $A$  and  $B$  have a supremum. Then  $A + B$  has also a supremum and

$$\sup(A + B) = \sup A + \sup B.$$

**Proof:** If  $t \in A + B$  then  $t = x + y$  with  $(x, y) \in A \times B$ . Then  $t = x + y \leq \sup A + \sup B$ , and so  $\sup A + \sup B$  is an upper bound for  $A + B$ . By the Completeness Axiom,  $A + B$  is bounded. Thus  $\sup(A + B) \leq \sup A + \sup B$ .

We now prove that  $\sup A + \sup B \leq \sup(A + B)$ . By the approximation property,  $\forall \varepsilon > 0 \exists a \in A$  and  $b \in B$  such that  $\sup A - \frac{\varepsilon}{2} < a$  and  $\sup B - \frac{\varepsilon}{2} < b$ . Observe that  $a + b \in A + B$  and so  $a + b \leq \sup(A + B)$ . Then

$$\sup A + \sup B - \varepsilon < a + b \leq \sup(A + B),$$

and by Lemma 95 we must have

$$\sup A + \sup B \leq \sup(A + B).$$

This completes the proof.  $\square$

**97 THEOREM (Archimedean Property of the Real Numbers)** If  $(x, y) \in \mathbb{R}^2$  with  $x > 0$ , then there exists a natural number  $n$  such that  $nx > y$ .

**Proof:** Consider the set

$$A = \{nx : n \in \mathbb{N}\}.$$

Since  $1 \cdot x \in A$ ,  $A$  is non-empty. If  $\forall n \in \mathbb{N}$  we had  $nx \leq y$ , then  $A$  would be bounded above by  $y$ . By the Completeness Axiom,  $A$  would have a supremum  $\sup A$ . Thus  $\forall n \in \mathbb{N}$ ,  $nx \leq \sup A$ . Since  $(n + 1)x \in A$ , we would also have

$$(n + 1)x \leq \sup A \implies nx \leq \sup A - x.$$

This means that  $\sup A - x$  is an upper bound for  $A$  which is smaller than its supremum, a contradiction. Thus there must be an  $n$  for which  $nx > y$ .  $\square$

**98 COROLLARY**  $\mathbb{N}$  is unbounded above.

**Proof:** This follows by taking  $x = 1$  in Theorem 97.  $\square$

The Completeness Axioms tells us, essentially, that there are no “holes” in the real numbers. We will see that this property distinguishes the reals from the rational numbers.

**99 LEMMA** [Hipassos of Metapontum]  $\sqrt{2}$  is irrational.

**Proof:** Assume there is  $s \in \mathbb{Q}$  such that  $s^2 = 2$ . We can find integers  $m, n \neq 0$  such that  $s = \frac{m}{n}$ . The crucial part of the argument is that we can choose  $m, n$  such that this fraction be in least terms, and hence,  $m, n$  must not be both even. Now,  $m^2 s^2 = n^2$ , that is  $2m^2 = n^2$ . This means that  $n^2$  is even. But then  $n$  itself must be even, since the product of two odd numbers is odd. Thus  $n = 2a$  for some non-zero integer  $a$  (since  $n \neq 0$ ). This means that  $2m^2 = (2a)^2 = 4a^2 \implies m^2 = 2a^2$ . This means once again that  $m$  is even. But then we have a contradiction, since  $m$  and  $n$  were not both even.  $\square$

**100 THEOREM**  $\mathbb{Q}$  is not complete.

**Proof:** We must shew that there is a non-empty set of rational numbers which is bounded above but that does not have a supremum in  $\mathbb{Q}$ . Consider the set  $A = \{r \in \mathbb{Q} : r^2 \leq 2\}$  of rational numbers. This set is bounded above by  $u = 2$ . For assume that there were a rogue element of  $A$ , say  $r_0$  such that  $r_0 > 2$ . Then  $r_0^2 > 4$  and so  $r_0$  would not belong to  $A$ , a contradiction. Thus  $r \leq 2$  for every  $r \in A$  and so  $A$  is bounded above. Suppose that  $A$  had a supremum  $s$ , which must satisfy  $s \leq 2$ . Now, by Lemma 99 we cannot have  $s^2 = 2$  and thus  $s^2 < 2$ . By Theorem 97 there is an integer  $n$  such that  $2 - s^2 > \frac{1}{10^n}$ . Put  $t = s + \frac{1}{10^{n-1}}$ , a rational number and observe that since  $s \leq 2$  one has

$$t^2 = s^2 + \frac{2s}{10^{n-1}} + \frac{1}{10^{2n-2}} < s^2 + \frac{2s}{10^{n-1}} + \frac{1}{10^{n-1}} \leq s^2 + \frac{5}{10^{n-1}} < s^2 + \frac{1}{10^n} < 2.$$

Thus  $t \in A$  and  $t > s$ , that is  $t$  is an element of  $A$  larger than its least upper bound, a contradiction. Hence  $A$  does not have a least upper bound.  $\square$

### 1.8.1 Greatest Integer Function

**101 THEOREM** Given  $y \in \mathbb{R}$  there exists a unique integer  $n$  such that

$$n \leq y < n + 1.$$

**Proof:** By Theorem 97, the set  $\{n \in \mathbb{Z} : n \leq y\}$  is non-empty and bounded above. We put

$$\lfloor y \rfloor = \sup\{n \in \mathbb{Z} : n \leq y\}.$$

□



$$\forall x \in \mathbb{R}, \quad \lfloor x \rfloor \leq x < \lfloor x \rfloor + 1.$$

**102 Definition** The unique integer in Theorem 101 is called the *floor* of  $x$  and is denoted by  $\lfloor x \rfloor$ .

The greatest integer function enjoys the following properties:

**103 THEOREM** Let  $\alpha, \beta \in \mathbb{R}, a \in \mathbb{Z}, n \in \mathbb{N}$ . Then

1.  $\lfloor \alpha + a \rfloor = \lfloor \alpha \rfloor + a$
2.  $\lfloor \frac{\alpha}{n} \rfloor = \lfloor \frac{\lfloor \alpha \rfloor}{n} \rfloor$
3.  $\lfloor \alpha \rfloor + \lfloor \beta \rfloor \leq \lfloor \alpha + \beta \rfloor \leq \lfloor \alpha \rfloor + \lfloor \beta \rfloor + 1$

**Proof:**

1. Let  $m = \lfloor \alpha + a \rfloor$ . Then  $m \leq \alpha + a < m + 1$ . Hence  $m - a \leq \alpha < m - a + 1$ . This means that  $m - a = \lfloor \alpha \rfloor$ , which is what we wanted.
2. Write  $\alpha/n$  as  $\alpha/n = \lfloor \alpha/n \rfloor + \theta, 0 \leq \theta < 1$ . Since  $n\lfloor \alpha/n \rfloor$  is an integer, we deduce by (1) that

$$\lfloor \alpha \rfloor = \lfloor n\lfloor \alpha/n \rfloor + n\theta \rfloor = n\lfloor \alpha/n \rfloor + \lfloor n\theta \rfloor.$$

Now,  $0 \leq \lfloor n\theta \rfloor \leq n\theta < n$ , and so  $0 \leq \lfloor n\theta \rfloor/n < 1$ . If we let  $\Theta = \lfloor n\theta \rfloor/n$ , we obtain

$$\frac{\lfloor \alpha \rfloor}{n} = \lfloor \frac{\alpha}{n} \rfloor + \Theta, \quad 0 \leq \Theta < 1.$$

This yields the required result.

3. From the inequalities  $\alpha - 1 < \lfloor \alpha \rfloor \leq \alpha, \beta - 1 < \lfloor \beta \rfloor \leq \beta$  we get  $\alpha + \beta - 2 < \lfloor \alpha \rfloor + \lfloor \beta \rfloor \leq \alpha + \beta$ . Since  $\lfloor \alpha \rfloor + \lfloor \beta \rfloor$  is an integer less than or equal to  $\alpha + \beta$ , it must be less than or equal to the integral part of  $\alpha + \beta$ , i.e.  $\lfloor \alpha + \beta \rfloor$ . We obtain thus  $\lfloor \alpha \rfloor + \lfloor \beta \rfloor \leq \lfloor \alpha + \beta \rfloor$ . Also,  $\alpha + \beta$  is less than the integer  $\lfloor \alpha \rfloor + \lfloor \beta \rfloor + 2$ , so its integer part  $\lfloor \alpha + \beta \rfloor$  must be less than  $\lfloor \alpha \rfloor + \lfloor \beta \rfloor + 2$ , but  $\lfloor \alpha + \beta \rfloor < \lfloor \alpha \rfloor + \lfloor \beta \rfloor + 2$  yields  $\lfloor \alpha + \beta \rfloor \leq \lfloor \alpha \rfloor + \lfloor \beta \rfloor + 1$ . This proves the inequalities.

□

**104 Definition** The *ceiling* of a real number  $x$  is the unique integer  $\lceil x \rceil$  satisfying the inequalities

$$\lceil x \rceil - 1 < x \leq \lceil x \rceil.$$

**105 Definition** The *fractional part* of a real number  $x$  is defined and denoted by

$$\{x\} = x - \lfloor x \rfloor.$$

Observe that  $0 \leq \{x\} < 1$ .

### Homework



**Problem 1.8.1** Let  $A$  and  $B$  be non-empty sets of real numbers. Put

$$-A = \{-x : x \in A\}, \quad A - B = \{a - b : (a, b) \in A \times B\}.$$

Prove that

1. If  $A$  is bounded above, then  $-A$  is bounded below and  $\sup A = -\inf(-A)$ .
2. If  $A$  and  $B$  are bounded above then  $A \cup B$  is also bounded above and  $\sup(A \cup B) = \max(\sup A, \sup B)$ .
3. If  $A$  is bounded above and  $B$  is bounded below, then  $A - B$  is bounded above and  $\sup(A - B) = \sup A - \inf B$ .

**Problem 1.8.2** Assume that  $A$  is a subset of the strictly positive real numbers. Prove that if  $A$  is bounded above, then the set  $A^{-1} = \{\frac{1}{x} : x \in A\}$  is bounded below and that  $\sup A = \frac{1}{\inf A^{-1}}$ .

**Problem 1.8.3** Let  $n \geq 2$  be an integer. Prove that

$$\max_{0 \leq x_1 \leq x_2 \leq \dots \leq x_n \leq 1} \left( \sum_{1 \leq i < j \leq n} (x_j - x_i) \right) = \left\lfloor \frac{n^2}{4} \right\rfloor.$$

**Problem 1.8.4** Find a non-zero polynomial  $P(x, y)$  such that

$$P(\lfloor 2t \rfloor, \lfloor 3t \rfloor) = 0$$

for all real  $t$ .

**Problem 1.8.5** Prove that the integers

$$\left\lfloor (1 + \sqrt{2})^n \right\rfloor$$

with  $n$  a positive integer, are alternately even or odd.

**Problem 1.8.6** Let  $x \in \mathbb{R}$  and let  $n$  be a strictly positive integer. Prove that

$$\left\lfloor nx \right\rfloor = \sum_{k=1}^{n-1} \left\lfloor x + \frac{k}{n} \right\rfloor.$$

**Problem 1.8.7 (Putnam 1948)** If  $n$  is a positive integer, demonstrate that

$$\left\lfloor \sqrt{n} + \sqrt{n+1} \right\rfloor = \left\lfloor \sqrt{4n+2} \right\rfloor.$$

**Problem 1.8.8** Find a formula for the  $n$ -th non-square.

**Problem 1.8.9** Prove that if  $a, b$  are strictly positive integers then

$$\frac{a^2}{b^2} < 2 \implies \frac{(a+2b)^2}{(a+b)^2} < 2.$$

Prove, moreover, that

$$\frac{(a+2b)^2}{(a+b)^2} - 2 < 2 - \frac{a^2}{b^2}.$$

This means that  $\frac{(a+2b)^2}{(a+b)^2}$  is closer to 2 than  $\frac{a^2}{b^2}$  is.

**Problem 1.8.10** Show that  $\forall x > 0$ ,  $x$  is farther from  $\sqrt{5}$  than  $\frac{2x+5}{x+2}$  is.

**Problem 1.8.11 (Existence of  $n$ -th Roots)** Let  $a > 0$  and let  $n \in \mathbb{R}$ ,  $n \geq 2$ . Prove that there is a unique  $b \in \mathbb{R}$  such that  $b^n = a$ .


# Chapter 2

## Topology of $\mathbb{R}$

### 2.1 Intervals

**Why bother?** In this section we give a more precise definition of what an interval is, and establish the interesting property that between any two real numbers there is always a rational number.

**106 Definition** An *interval*  $I$  is a subset of the real numbers with the following property: if  $s \in I$  and  $t \in I$ , and if  $s < x < t$ , then  $x \in I$ . In other words, intervals are those subsets of real numbers with the property that every number between two elements is also contained in the set. Since there are infinitely many decimals between two different real numbers, intervals with distinct endpoints contain infinitely many members.

 The empty set  $\emptyset$  is trivially an interval.

We will now establish that there are nine types of intervals.











Interval Notation	Set Notation	Graphical Representation
$[a; b]$	$\{x \in \mathbb{R} : a \leq x \leq b\}$ <sup>1</sup>	
$]a; b[$	$\{x \in \mathbb{R} : a < x < b\}$	
$[a; b[$	$\{x \in \mathbb{R} : a \leq x < b\}$	
$]a; b]$	$\{x \in \mathbb{R} : a < x \leq b\}$	
$]a; +\infty[$	$\{x \in \mathbb{R} : x > a\}$	
$[a; +\infty[$	$\{x \in \mathbb{R} : x \geq a\}$	
$] -\infty; b[$	$\{x \in \mathbb{R} : x < b\}$	
$] -\infty; b]$	$\{x \in \mathbb{R} : x \leq b\}$	
$] -\infty; +\infty[$	$\mathbb{R}$	

Table 2.1: Types of Intervals. Observe that we indicate that the endpoints are included by means of shading the dots at the endpoints and that the endpoints are excluded by not shading the dots at the endpoints.

 If  $x \in \mathbb{R}$ , then  $\{x\} = [x; x]$ .

**107 THEOREM** The only kinds of intervals are those sets shewn in Table 2.1, and conversely, all sets shewn in this table are intervals.

**Proof:** The converse is easily established, so assume that  $I \subseteq \mathbb{R}$  possesses the property that  $\forall (a, b) \in I^2$ ,  $[a; b] \subseteq I$ . Since  $\emptyset$  is an interval one may assume that  $I \neq \emptyset$ . Let  $a \in I$  be a fixed element of  $I$  and put  $M_a = \{x \in I : x \leq a\} = ]-\infty; a] \cap I$  and  $N_a = \{x \in I : x \geq a\} = [a; +\infty[ \cap I$ .

If  $N_a$  is not bounded above, then  $\forall b \in [a; +\infty[$ ,  $\exists c \in N_a$  such that  $b \leq c$ . Since  $a \leq b \leq c$ , this entails that  $b \in N_a$ . Thus  $N_a = [a; +\infty[$ .

If  $N_a$  is bounded above, then it has supremum  $s = \sup(N_a)$  and  $N_a \subseteq [a; s[$ . By Theorem 93,  $\forall b \in [a; s[$ ,  $c \in N_a$  such that  $b \leq c$ , and since  $a \leq b \leq c$ , this entails that  $b \in N_a$ . Thus

$$[a; s[ \subseteq N_a \subseteq [a; s[ ,$$

and so  $N_a = [a; s[$  or  $N_a = [a; s]$ .

Thus  $N_a$  is one among three possible forms:  $[a; +\infty[$ ,  $[a; s]$ , or  $[a; s[$ . Applying a similar reasoning, one obtains that  $M_a$  is of one of the forms  $]-\infty; a]$ ,  $]l; a]$ , or  $]l; a[$ , where  $l = \inf(M_a)$ . Since  $I = M_a \cup N_a$ , there are 3 choices for  $M_a$  and 3 for  $N_a$ , hence there are  $3 \cdot 3 = 9$  choices for  $I$ . The result is established.  $\square$

**108 Example** Determine  $\bigcap_{k=1}^{\infty} \left[1 - \frac{1}{2^k}; 1 + \frac{1}{k}\right]$ .

**Solution:** Observe that the intervals are, in sequence,

$$\left[\frac{1}{2}; 2\right]; \left[\frac{3}{4}; \frac{3}{2}\right]; \left[\frac{7}{8}; \frac{4}{3}\right]; \dots$$

We claim that  $\bigcap_{k=1}^{\infty} \left[1 - \frac{1}{2^k}; 1 + \frac{1}{k}\right] = 1$ . For we see that

$$\forall k \geq 1, \quad \frac{1}{2} \leq 1 - \frac{1}{2^k} < 1 < 1 + \frac{1}{k} \leq 2,$$

so 1 is in every interval. Could this intersection contain a number smaller than 1? No, for if  $\frac{1}{2} \leq a < 1$ , then we can take  $k$  large enough so that

$$a < 1 - \frac{1}{2^k},$$

for example

$$a < 1 - \frac{1}{2^k} \implies k > -\log_2(1 - a),$$

so taking  $k \geq \lceil -\log_2(1 - a) \rceil + 1$  will work. Could the intersection contain a number  $b$  larger than 1? No, for if  $1 < b < 2$ , then we can take  $k$  large enough so that

$$1 + \frac{1}{k} < b,$$

for example

$$1 + \frac{1}{k} < b \implies k > \frac{1}{b - 1},$$

so taking  $k \geq \lceil \frac{1}{b - 1} \rceil + 1$  will work. Hence the only number in the intersection is 1.

## 2.2 Dense Sets

**109 Definition** A set  $B \subseteq \mathbb{R}$  is *dense in*  $A \subseteq \mathbb{R}$  if  $\forall (a_1, a_2) \in A^2$ ,  $a_1 < a_2$ ,  $\exists b \in B$  such that  $a_1 < b < a_2$ , that is, between any two different elements of  $A$  one can always find an element of  $B$ .

**110 THEOREM**  $\mathbb{Q}$  is dense in  $\mathbb{R}$ .

**Proof:** Let  $x, y$  be real numbers with  $x < y$ . Since there are infinitely many positive integers, there must be a positive integer  $n$  such that  $n > \frac{1}{y-x}$  by the Archimedean Property of  $\mathbb{R}$ . Consider the rational number  $r = \frac{m}{n}$ , where  $m$  is the least natural number with  $m > nx$ . This means that

$$m > nx \geq m - 1.$$

We claim that  $x < \frac{m}{n} < y$ . The first inequality is clear, since by choice  $x < \frac{m}{n}$ . For the second inequality observe that, again

$$nx \geq m - 1 \text{ and } y - x > \frac{1}{n} \implies x > \frac{m}{n} - \frac{1}{n} \text{ and } y > x + \frac{1}{n} \implies y > \frac{m}{n} - \frac{1}{n} + \frac{1}{n} = \frac{m}{n}.$$

Thus  $\frac{m}{n}$  is a rational number between  $x$  and  $y$ .  $\square$

**111 THEOREM**  $\mathbb{R} \setminus \mathbb{Q}$  is dense in  $\mathbb{R}$ .

**Proof:** Let  $a < b$  be two real numbers. By Theorem 110, there is a rational number  $r$  with  $\frac{a}{\sqrt{2}} < r < \frac{b}{\sqrt{2}}$ . But then  $a < \sqrt{2}r < b$ , and the number  $\sqrt{2}r$  is an irrational number.  $\square$

**112 THEOREM (Dirichlet)** For any real number  $\theta$  and any integer  $Q \geq 1$ , there exist integers  $a$  and  $q$ ,  $1 \leq q \leq Q$ , such that

$$\left| \theta - \frac{a}{q} \right| \leq \frac{1}{qQ}.$$

**Proof:** For  $1 \leq n \leq Q$ , let

$$I_n = \left[ \frac{n-1}{Q}; \frac{n}{Q} \right].$$

Thus these  $Q$  intervals partition the interval  $[0; 1[$ . The  $Q+1$  numbers

$$\{0\theta\}, \{1\theta\}, \{2\theta\}, \dots, \{Q\theta\}$$

lie in  $[0; 1[$ . Hence by the pigeonhole principle there is an  $n$  such that  $I_n$  contains at least two of these numbers, say

$$\{q_1\theta\} \in I_n, \quad \{q_2\theta\} \in I_n, \quad 0 \leq q_1 < q_2 \leq Q.$$

Put  $q = q_2 - q_1$ ,  $a = [q_2\theta] - [q_1\theta]$ . Since  $\{q_1\theta\} \in I_n, \{q_2\theta\} \in I_n$  we must have

$$|\{q_2\theta\} - \{q_1\theta\}| < \frac{1}{Q}.$$

But

$$\{q_2\theta\} - \{q_1\theta\} = q_2\theta - [q_2\theta] - q_1\theta + [q_1\theta] = q\theta - a,$$

whence the result.  $\square$

**113 COROLLARY** If  $\theta$  is irrational prove that there exist infinitely many rational numbers  $\frac{a}{q}$ ,  $\gcd(a, q) = 1$ , such that  $\theta$  lies in the open intervals  $\left] \frac{a}{q} - \frac{1}{q^2}; \frac{a}{q} + \frac{1}{q^2} \right[$ .

**Proof:** Suppose that  $\left| \theta - \frac{a_r}{q_r} \right| < \frac{1}{q_r^2}$  for  $1 \leq r \leq R$ . Since the differences  $\theta - \frac{a_r}{q_r}$  are non-zero, we may choose  $Q$  so large in Theorem 112 that none of these rational numbers is a solution of  $\left| \theta - \frac{a}{q} \right| < \frac{1}{qQ}$ . Since this latter inequality does have a solution, the  $R$  given rational approximations do not exhaust the set of solutions of  $\left| \theta - \frac{a}{q} \right| < \frac{1}{q^2}$ .  $\square$

## Homework

**Problem 2.2.1** Determine  $\bigcap_{1 \leq k \leq 500} [k; 1001 - k]$ .

**Problem 2.2.2** Determine  $\bigcup_{k=1}^{\infty} \left[ 1; 1 + \frac{1}{k} \right]$ .

**Problem 2.2.3** Determine  $\bigcup_{k=1}^{\infty} [-k; k]$ .

**Problem 2.2.4** Determine  $\bigcap_{k=1}^{\infty} \left[ 1; 1 + \frac{1}{k} \right]$ .

**Problem 2.2.5** Determine  $\bigcap_{k=1}^{\infty} [k; +\infty[$ .

**Problem 2.2.6** Determine  $\bigcap_{k=1}^{\infty} \left] 1; 1 + \frac{1}{k} \right[$ .

**Problem 2.2.7** Let  $I = [a; b]$ , and  $I' = [a'; b']$  be closed intervals in  $\mathbb{R}$ . Prove that  $I \subseteq I'$  if and only if  $a' \leq a$  and  $b \leq b'$ .

**Problem 2.2.8** Let

$$\mathbb{Q} + \sqrt{2}\mathbb{Q} = \{a + \sqrt{2}b : (a, b) \in \mathbb{Q}^2\}$$

and define addition on this set as

$$(a + \sqrt{2}b) + (c + \sqrt{2}d) = (a + c) + \sqrt{2}(b + d),$$

and multiplication as

$$(a + \sqrt{2}b)(c + \sqrt{2}d) = (ac + 2bd) + \sqrt{2}(ad + bc).$$

Then  $(\mathbb{Q} + \sqrt{2}\mathbb{Q}, \cdot, +)$  is a field.

**Problem 2.2.9** Put  $D = \{x : x = q^2 \text{ or } x = -q^2, q \in \mathbb{Q}\}$ . Prove that  $D$  is dense in  $\mathbb{R}$ .

**Problem 2.2.10** A dyadic rational is a rational number of the form  $\frac{m}{2^n}$ , where  $m \in \mathbb{Z}$ ,  $n \in \mathbb{N}$ . Prove that the set of dyadic rationals is dense in  $\mathbb{R}$ .

## 2.3 Open and Closed Sets

**Why bother?** Many of the properties that we will study in these notes generalise to sets other than  $\mathbb{R}$ . To better understand what is it from the features of  $\mathbb{R}$  that is essential for a generalisation, the language of topology is used.

**114 Definition** The open ball  $\mathcal{B}_{x_0}(r)$  centred at  $x = x_0$  and radius  $\epsilon > 0$  is the set

$$\mathcal{B}_{x_0}(\epsilon) = ]x_0 - \epsilon; x_0 + \epsilon[.$$

**115 Definition** A set  $\mathcal{N}_{x_0} \subseteq \mathbb{R}$  is an open neighbourhood of a point  $x_0$  if  $\exists \epsilon > 0$  such that  $\mathcal{B}_{x_0}(\epsilon) \subseteq \mathcal{N}_{x_0}$ , that is, there is a sufficiently small open ball containing  $x_0$  completely contained in  $\mathcal{N}_{x_0}$ .

**116 Definition** A set  $U \subseteq \mathbb{R}$  is said to be open in  $\mathbb{R}$  if  $\forall x \in U$  there is an open neighbourhood  $\mathcal{N}_{x_0}$  such that  $\mathcal{N}_{x_0} \subseteq U$ . A set  $F \subseteq \mathbb{R}$  is said to be closed in  $\mathbb{R}$  if its complement  $U = \mathbb{R} \setminus F$  is open in  $\mathbb{R}$ .

**117 THEOREM** Every open ball is open.

**Proof:** Let  $\mathcal{B}_{x_0}(r)$  with  $r > 0$  be an open ball and let  $x \in \mathcal{B}_{x_0}(r)$ . We must show that there is a sufficiently small neighbourhood of  $x$  completely within  $\mathcal{B}_{x_0}(r)$ . That is, we search for  $\epsilon > 0$  such that  $y \in \mathcal{B}_x(\epsilon) \implies y \in \mathcal{B}_{x_0}(r)$ .

Now,

$$y \in \mathcal{B}_x(\varepsilon) \implies y \in \mathcal{B}_{x_0}(r) \iff |y - x| < \varepsilon \implies |y - x_0| < r.$$

By the Triangle Inequality

$$|y - x_0| \leq |y - x| + |x - x_0| < \varepsilon + |x - x_0|.$$

So, as long as

$$\varepsilon + |x - x_0| < r,$$

we will be within  $\mathcal{B}_{x_0}(r)$ . One can take

$$\varepsilon = \frac{r - |x - x_0|}{2}.$$

□

**118 Example** The open intervals  $]a; b[$ ,  $]a; +\infty[$ ,  $] -\infty; b[$ ,  $] -\infty; +\infty[$ , are open in  $\mathbb{R}$ .

The closed intervals  $\{a\}$ ,  $[a; b]$ ,  $[a; +\infty[$ ,  $] -\infty; b]$ ,  $] -\infty; +\infty[ = \mathbb{R}$ , are closed in  $\mathbb{R}$ .

The sets  $\emptyset$  and  $\mathbb{R}$  are simultaneously open and closed in  $\mathbb{R}$ .

The intervals  $]a; b]$  and  $[a; b[$  are neither open nor closed in  $\mathbb{R}$ .

**119 THEOREM** The union of any (finite or infinite) number of open sets in  $\mathbb{R}$  is open in  $\mathbb{R}$ . The union of a finite number of closed in  $\mathbb{R}$  sets is closed in  $\mathbb{R}$ .

The intersection of a finite number of open sets in  $\mathbb{R}$  is open in  $\mathbb{R}$ . The intersection of any (finite or infinite) number of closed sets in  $\mathbb{R}$  is closed in  $\mathbb{R}$ .

**Proof:** Let  $U_1, U_2, \dots$ , be a sequence of open sets in  $\mathbb{R}$  (some may be empty) and consider  $x \in \bigcup_{n=1}^{\infty} U_n$ . There is an index  $N$  such that  $x \in U_N$ . Since  $U_N$  is open in  $\mathbb{R}$ , there is an open neighbourhood of  $x$   $]x - \varepsilon; x + \varepsilon[ \subseteq U_N$ , for  $\varepsilon > 0$  small enough. But then

$$]x - \varepsilon; x + \varepsilon[ \subseteq U_N \subseteq \bigcup_{n=1}^{\infty} U_n,$$

and so given an arbitrary point of the union, there is a small enough open neighbourhood enclosing the point and within the union, meaning that the union is open.

If  $\bigcap_{n=1}^{\infty} F_n$  is an arbitrary intersection of closed sets, then there are open sets  $U_n = \mathbb{R} \setminus F_n$ . By the De Morgan Laws,

$$\bigcap_{n=1}^{\infty} F_n = \bigcap_{n=1}^{\infty} (\mathbb{R} \setminus U_n) = \mathbb{R} \setminus \bigcup_{n=1}^{\infty} U_n,$$

and since  $\bigcup_{n=1}^{\infty} U_n$  is open by the above paragraph,  $\bigcap_{n=1}^{\infty} F_n$  is the complement of an open set, that is, it is closed.

Let  $U_1, U_2, \dots, U_L$  be a sequence of open sets in  $\mathbb{R}$  and consider  $x \in \bigcap_{n=1}^L U_n$ . Then  $x$  belongs to each of the  $U_k$  and so there are  $\varepsilon_k > 0$  such that  $x \in ]x - \varepsilon_k; x + \varepsilon_k[ \subseteq U_k$ . Let  $\varepsilon = \min_{1 \leq k \leq L} \varepsilon_k$  be the smallest one of such. But then for all  $k$ ,

$$]x - \varepsilon; x + \varepsilon[ \subseteq ]x - \varepsilon_k; x + \varepsilon_k[ \subseteq U_k, \implies ]x - \varepsilon; x + \varepsilon[ \subseteq \bigcap_{n=1}^L U_n,$$

and so given an arbitrary point of the intersection, there is a small enough open neighbourhood enclosing the point and within the intersection, meaning that the intersection is open.

Using the De Morgan Laws and the preceding paragraph, the remaining statement can be proved. □

**120 Example** The intersection of an infinite number of open sets may not be open. For example

$$\bigcap_{k=1}^{\infty} \left] 1 - \frac{1}{n+1}; 2 - \frac{1}{n+1} \right[ = \left] 1; 2 \right[ ,$$

which is neither open nor closed.

**121 THEOREM (Characterisation of the Open Sets of  $\mathbb{R}$ )** A set  $A \subseteq \mathbb{R}$  is open if and only if it is the countable union of open sets of  $\mathbb{R}$ .

## 2.4 Interior, Boundary, and Closure of a Set

**122 Definition** Let  $A \subseteq \mathbb{R}$ . The *interior* of  $A$  is defined and denoted by


$$\overset{\circ}{A} = \bigcup_{\substack{\Omega \subseteq A \\ \Omega \text{ open}}} \Omega,$$

that is, the largest open set inside  $A$ . The points of  $\overset{\circ}{A}$  are called the *interior points* of  $A$ .

**123 Definition** Let  $A \subseteq \mathbb{R}$ . The *closure* of  $A$  is defined and denoted by

$$\overline{A} = \bigcup_{\substack{\Omega \supseteq A \\ \Omega \text{ closed}}} \Omega,$$

that is, the smallest closed set containing  $A$ . The points of  $\overline{A}$  are called the *adherence points* of  $A$ .

 One always has  $\overset{\circ}{A} \subseteq A \subseteq \overline{A}$ . A set  $U$  is open if and only if  $U = \overset{\circ}{U}$ . A set  $F$  is closed if and only if  $F = \overline{F}$ .

**124 Definition** Let  $A \subseteq \mathbb{R}$ . The *boundary* of  $A$  is defined and denoted by

$$\mathbf{Bdy}(A) = \overline{A} - \overset{\circ}{A}.$$

The elements of  $\mathbf{Bdy}(A)$  are called the *boundary points* of  $A$ .

**125 Example** We have

1.  $\overset{\circ}{\left] 0; 1 \right[} = \left] 0; 1 \right[$ ,  $\overline{\left] 0; 1 \right[} = \left[ 0; 1 \right]$ ,  $\mathbf{Bdy}\left(\left] 0; 1 \right[\right) = \{0, 1\}$
2.  $\overset{\circ}{\{0, 1\}} = \emptyset$ ,  $\overline{\{0, 1\}} = \{0, 1\}$ ,  $\mathbf{Bdy}(\{0, 1\}) = \{0, 1\}$
3.  $\overset{\circ}{\mathbb{Q}} = \emptyset$ ,  $\overline{\mathbb{Q}} = \mathbb{R}$ ,  $\mathbf{Bdy}(\mathbb{Q}) = \mathbb{R}$

**126 THEOREM** Let  $A \subseteq \mathbb{R}$ . Then

$$\mathbb{R} \setminus \overset{\circ}{A} = \overline{\mathbb{R} \setminus A}, \quad \mathbb{R} \setminus \overline{A} = \overset{\circ}{\mathbb{R} \setminus A}.$$

**Proof:** The theorem follows from the De Morgan Laws, as

$$\mathbb{R} \setminus \overset{\circ}{A} = \mathbb{R} \setminus \bigcup_{\substack{\Omega \subseteq A \\ \Omega \text{ open}}} \Omega = \bigcap_{\substack{\Omega \subseteq A \\ \Omega \text{ open}}} (\mathbb{R} \setminus \Omega) = \bigcap_{\substack{\mathbb{R} \setminus A \subseteq \mathbb{R} \setminus \Omega \\ \Omega \text{ open}}} (\mathbb{R} \setminus \Omega) = \bigcap_{\substack{\mathbb{R} \setminus A \subseteq F \\ F \text{ closed}}} F = \overline{\mathbb{R} \setminus A},$$

and

$$\mathbb{R} \setminus \overline{A} = \mathbb{R} \setminus \bigcap_{\substack{F \supseteq A \\ F \text{ closed}}} F = \bigcup_{\substack{F \supseteq A \\ F \text{ closed}}} (\mathbb{R} \setminus F) = \bigcup_{\substack{\mathbb{R} \setminus A \supseteq \mathbb{R} \setminus F \\ F \text{ closed}}} (\mathbb{R} \setminus F) = \bigcap_{\substack{\mathbb{R} \setminus A \supseteq \Omega \\ \Omega \text{ open}}} \Omega = \overset{\circ}{\mathbb{R} \setminus A}.$$

□

**127 THEOREM**  $x \in \bar{A} \iff \forall \mathcal{N}_x, \mathcal{N}_x \cap A \neq \emptyset$ . That is,  $x$  is an adherent point if and only if every neighbourhood of  $x$  has a nonempty intersection with  $A$ .

**Proof:** Assume  $x \in \bar{A}$  and let  $r > 0$ . If  $\mathcal{N}_x \cap A = \emptyset$ , then  $\mathcal{N}_x \subseteq \mathbb{R} \setminus A$ . Since  $\mathcal{N}_x$  is open, we have—in particular— $\mathcal{N}_x \subseteq \overset{\circ}{\mathbb{R} \setminus A} = \mathbb{R} \setminus \bar{A}$  by Theorem 126. This means that  $x \notin \bar{A}$ , a contradiction.

Conversely, assume that for all neighbourhoods  $\mathcal{N}_x$  of  $x$  we have  $\mathcal{N}_x \cap A \neq \emptyset$ . If  $x \notin \bar{A}$  then  $x \in \mathbb{R} \setminus \bar{A} = \overset{\circ}{\mathbb{R} \setminus A}$ . Since  $\overset{\circ}{\mathbb{R} \setminus A}$  is open there is an  $r' > 0$  such that  $\mathcal{N}_{x-r'; x+r'} \subseteq \overset{\circ}{\mathbb{R} \setminus A} \subseteq \mathbb{R} \setminus A$ . But then  $\mathcal{N}_{x-r'; x+r'} \cap A = \emptyset$ , a contradiction.  $\square$

**128 THEOREM** Let  $\emptyset \subsetneq A \subseteq \mathbb{R}$  be bounded above. Then  $\sup A \in \bar{A}$ . If, moreover,  $A$  is closed then  $\sup(A) \in A$ .

**Proof:** Let  $r > 0$ . By Theorem 93, there exists  $a \in A$  such that  $\sup(A) - r < a$ , which gives  $|\sup(A) - a| < r$ . This shows that  $\sup A - r; \sup A + r \cap A \neq \emptyset$  regardless of how small  $r > 0$  might be and, hence,  $\sup(A) \in \bar{A}$  by Theorem 127. If  $A$  is closed, then  $A = \bar{A}$ .  $\square$

**129 Definition** Let  $A \subseteq \mathbb{R}$ . A point  $x \in A$  is called an *isolated point* of  $A$  if there exists an  $r > 0$  such that  $\mathcal{B}_x(r) \cap A = \{x\}$ . The set of isolated points of  $A$  is denoted by  $A^*$ .

A point  $y \in \mathbb{R}$  is called an *accumulation point* of  $A$  in  $\mathbb{R}$  if

$$\forall \mathcal{N}_x, (\mathcal{N}_x \setminus \{x\}) \cap A \neq \emptyset,$$

that is, if any neighbourhood of  $x$  meets  $A$  at a point different than  $x$ . The set of accumulation points of  $A$  is called the *derived set* of  $A$  and is denoted by  $\text{Acc}(A)$ .

**130 Example** We have

1.  $0$  is an isolated point of the set  $A = \{0\} \cup [1; 2]$ .
2. Every point of the set  $A = \left\{1, \frac{1}{2}, \frac{1}{3}, \dots\right\}$  is isolated. This is because we may take  $r = \frac{1}{2^{n+2}}$  in the definition of isolated point, and then  $\left] \frac{1}{n} - \frac{1}{2^{n+2}}; \frac{1}{n} + \frac{1}{2^{n+2}} \right[ \cap A = \left\{ \frac{1}{n} \right\}$ . Observe that  $\frac{1}{n} - \frac{1}{n+1} = \frac{1}{n(n+1)}$  and  $\frac{1}{n-1} - \frac{1}{n} = \frac{1}{n(n-1)}$  and that  $2^{n+2} > \max(n(n+1), n(n-1))$ .
3.  $0$  is an accumulation point of  $A = \left\{1, \frac{1}{2}, \frac{1}{3}, \dots\right\}$ .

**131 THEOREM**  $x$  is an accumulation point of  $A$  if and only if every neighbourhood of  $x$  in  $\mathbb{R}$  has an infinite number of points of  $A$ .

**Proof:** Suppose  $x \in A'$ . Suppose a neighbourhood of  $x$  had only finitely many elements of  $A$ , say  $\{y_1, y_2, \dots, y_n\}$ . Take  $2r = \min_{1 \leq k \leq n} |y_k - x|$ . Then  $\left( \mathcal{N}_x \setminus \{x\} \right) \cap A = \emptyset$  contradicting the fact that every neighbourhood of  $x$  meets  $A$  at a point different from  $x$ .

Conversely if every neighbourhood of  $x$  in  $\mathbb{R}$  has an infinite number of points of  $A$ , then a fortiori, any intersection of such a neighbourhood with  $A$  will contain a point different from  $x$ , and so  $x \in \text{Acc}(A)$ .  $\square$

**132 THEOREM** A set is closed if and only if it contains all its accumulation points.



**Proof:** If  $A$  is closed then  $\mathbb{R} \setminus A$  is open. If  $c \in \mathbb{R} \setminus A$  then there exists  $r > 0$  such that  $c - r ; c + r \left[ \subseteq \mathbb{R} \setminus A$ , a neighbourhood that clearly does not contain any points of  $A$ , which means  $c \notin \text{Acc}(A)$ .

Conversely, suppose a set  $\text{Acc}(A) \subseteq A$ . We will prove that  $\mathbb{R} \setminus A$  is open. If  $x \in \mathbb{R} \setminus A$ , then a fortiori,  $x \notin \text{Acc}(A)$ . This means that there is an  $r > 0$  such that  $x - r ; x + r \left[ \cap A = \emptyset$ . Hence  $x - r ; x + r \left[ \subseteq \mathbb{R} \setminus A$ , and so  $\mathbb{R} \setminus A$  is open.  $\square$

 One has

$$A^* \subseteq A, \quad \overline{A} - A \subseteq \text{Acc}(A), \quad A^* \cap \text{Acc}(A) = \emptyset, \quad A^* \cup \text{Acc}(A) = \overline{A}.$$

## 2.5 Connected Sets

**133 Definition** A set  $X \subseteq \mathbb{R}$  is connected if, given open sets  $U, V$  of  $\mathbb{R}$  with  $U \cup V = X$ ,  $U \cap V = \emptyset$ , either  $U = \emptyset$  or  $V = \emptyset$ .

**134 THEOREM** If  $X \subseteq \mathbb{R}$  is connected, and if  $(a, c) \in X^2$ ,  $b \in \mathbb{R}$ , are such that  $a < b < c$  then  $b \in X$ .

**135 COROLLARY** The only connected sets of  $\mathbb{R}$  are the intervals. In particular,  $\mathbb{R}$  is connected.

## 2.6 Compact Sets

**136 Definition** A sequence of open sets  $U_1, U_2, \dots$  is said to be an *open cover* for  $A \subseteq \mathbb{R}$  if  $A \subseteq \bigcup_{n=1}^{\infty} U_n$ .  $U_1, U_2, \dots$  has a *subcover*  $U_{k_1}, U_{k_2}, \dots$  of  $A$  if  $A \subseteq \bigcup_{n=1}^{\infty} U_{k_n}$ .

**137 Definition** A set of real numbers is said to be *compact in*  $\mathbb{R}$  if every open cover of the set has a finite subcover.<sup>2</sup>

**138 Example** Since  $\mathbb{R} = \bigcup_{n \in \mathbb{Z}} ]n - 1 ; n + 1 \left[$ , the sequence of intervals  $]n - 1 ; n + 1 \left[$ ,  $n \in \mathbb{Z}$  is a cover for  $\mathbb{R}$ .

**139 THEOREM** Let  $a, b$  be real numbers with  $a \leq b$ . The closed interval  $[a ; b]$  is compact in  $\mathbb{R}$ .

**Proof:** Let  $U_1, U_2, \dots$  be an open cover for  $[a ; b]$ . Let  $E$  be the collection of all  $x \in [a ; b]$  such that  $[a ; x]$  has a finite subcover from the  $U_i$ . We will shew that  $b \in E$ .

Since  $a \in \bigcup_{i=1}^{\infty} U_i$ , there exists  $U_r$  such that  $a \in U_r$ . Thus  $\{a\} = [a ; a] \subseteq U_r$  and so  $E \neq \emptyset$ . Clearly,  $b$  is an upper bound for  $E$ . By the Completeness Axiom,  $\sup E$  exists. We will shew that  $b = \sup E$ .

By Theorem 128,  $\sup E \in [a ; b] \subseteq \bigcup_{i=1}^{\infty} U_i$ , hence there exists  $U_s$  such that  $\sup E \in U_s$ . Since  $U_s$  is open, there exists  $\varepsilon > 0$  such that  $\sup E - \varepsilon ; \sup E + \varepsilon \left[ \subseteq U_s$ . By Theorem 93 there is  $x \in E$  such that  $\sup E - \varepsilon < x \leq \sup E$ . Thus there is a finite subcover from the  $U_i$ , say,  $U_{p_1}, U_{p_2}, \dots, U_{p_n}$  such that  $[a ; x] \subseteq \bigcup_{i=1}^n U_{k_i}$ .

We thus have

$$[a ; \sup E] \subseteq [a ; x] \cup ]\sup E - \varepsilon ; \sup E + \varepsilon \left[ \subseteq \left( \bigcup_{i=1}^n U_{k_i} \right) \cup U_s,$$

<sup>2</sup>This definition is appropriate for  $\mathbb{R}$  but it is not valid in general. However, it very handy for one-variable calculus, hence we will retain it.

a finite subcover. This means that  $\sup E \in E$ .

Suppose now that  $\sup E < b$ , and consider  $y = \sup E + \frac{1}{2} \min(b - \sup E, \epsilon)$ . Then

$$\sup E < y, \quad [a; y] = [a; \sup E] \cup [\sup E; y] \subseteq \left( \bigcup_{i=1}^n U_{k_i} \right) \cup U_s,$$

whence  $y \in E$ , contradicting the definition of  $\sup E$ . This proves that  $\sup E = b$  and finishes the proof of the theorem.  $\square$

**140 THEOREM (Heine-Borel)** A set  $A$  of  $\mathbb{R}$  is closed and bounded if and only if it is compact.

**Proof:** Let  $A$  be closed and bounded in  $\mathbb{R}$ , and let  $U_1, U_2, \dots$ , be an open cover for  $A$ . There exist  $(a, b) \in \mathbb{R}^2$ ,  $a \leq b$ , such that  $A \subseteq [a; b]$ . Since

$$[a; b] \subseteq (\mathbb{R} \setminus A) \cup \bigcup_{i=1}^{\infty} U_i,$$

by Theorem 139 there is a finite subcover of the  $U_i$ , say,  $U_{k_i}$  such that

$$[a; b] \subseteq (\mathbb{R} \setminus A) \cup \bigcup_{i=1}^{\infty} U_{k_i}.$$

Therefore

$$A = A \cap [a; b] \subseteq [a; b] \subseteq \bigcup_{i=1}^{\infty} U_{k_i},$$

and so  $A$  admits an open subcover.

Conversely, suppose that every open cover of  $A$  admits a finite subcover. The open cover  $\left] -n; n \right[, n \in \mathbb{R}$  of  $A$  must admit a finite subcover by our assumption, hence there is  $N \in \mathbb{N}$  such that  $A \subseteq \left] -N; N \right[$ , meaning that  $A$  is bounded. Let us show now that  $\mathbb{R} \setminus A$  is open.

Let  $x \in \mathbb{R} \setminus A$ . We have

$$\bigcup_{n \geq 1} \left( \mathbb{R} \setminus \left[ x - \frac{1}{n}; x + \frac{1}{n} \right] \right) = \mathbb{R} \setminus \bigcap_{n \geq 1} \left[ x - \frac{1}{n}; x + \frac{1}{n} \right] = \mathbb{R} \setminus \{x\} \supseteq A,$$

since  $x \notin A$ . By hypothesis there is  $N \in \mathbb{N}$  and  $n_1, n_2, \dots, n_N$  such that

$$A \subseteq \bigcup_{k=1}^N \left( \mathbb{R} \setminus \left[ x - \frac{1}{n_k}; x + \frac{1}{n_k} \right] \right) \subseteq \mathbb{R} \setminus \left[ x - \frac{1}{n_m}; x + \frac{1}{n_m} \right],$$

where  $m = \max(n_1, n_2, \dots, n_N)$ . This gives  $\left[ x - \frac{1}{n_m}; x + \frac{1}{n_m} \right] \subseteq \mathbb{R} \setminus A$ , meaning that  $\mathbb{R} \setminus A$  is open, whence  $A$  is closed.

$\square$

**141 COROLLARY (Cantor's Intersection Theorem)** Let

$$[a_1; b_1] \supseteq [a_2; b_2] \supseteq [a_3; b_3] \supseteq \dots$$

be a sequence of non-empty, bounded, nested closed intervals. Then

$$\bigcap_{j=1}^{\infty} [a_j; b_j] \neq \emptyset.$$

**Proof:** Assume that  $[a_1; b_1] \cap \bigcap_{j=2}^{\infty} [a_j; b_j] = \emptyset$ . Then

$$[a_1; b_1] \subseteq \mathbb{R} \setminus \bigcap_{j=2}^{\infty} [a_j; b_j] = \bigcup_{j=2}^{\infty} (\mathbb{R} \setminus [a_j; b_j]).$$

The  $\mathbb{R} \setminus [a_j; b_j]$  for an open cover for  $[a_1; b_1]$ , which is closed and bounded. By Theorem 7 we have

$$[a_j; a_j] \subseteq [a_i; b_i] \implies \mathbb{R} \setminus [a_i; b_i] \subseteq \mathbb{R} \setminus [a_j; b_j].$$

By the Heine-Borel Theorem 140 there is a finite subcover, say

$$[a_1; b_1] \subseteq \bigcup_{j=1}^N (\mathbb{R} \setminus [a_{n_j}; b_{n_j}]) \subseteq \mathbb{R} \setminus [a_{n_N}; b_{n_N}].$$

But then  $[a_{n_N}; b_{n_N}] \subseteq \mathbb{R} \setminus [a_1; b_1]$ , which contradicts  $[a_{n_N}; b_{n_N}] \subseteq [a_1; b_1]$ , and the proof is complete.  $\square$

**142 THEOREM (Bolzano-Weierstrass)** Every bounded infinite set of  $\mathbb{R}$  has at least one accumulation point.

**Proof:** Let  $A$  be a bounded set of  $\mathbb{R}$  with  $\text{Acc}(A) = \emptyset$ . Then  $A^* = A = \overline{A}$ . Notice that then every element of  $A$  is an isolated point of  $A$ , and hence,

$$\forall a \in A, \exists r_a > 0, \text{ such that } ]a - r_a; a + r_a[ \cap A = \{a\}.$$

Observe that

$$A \subseteq \bigcup_{a \in A} ]a - r_a; a + r_a[ ,$$

and so the  $]a - r_a; a + r_a[$  form an open cover for  $A$ . Since  $A = \overline{A}$ ,  $A$  is closed. By the Heine-Borel Theorem 140  $A$  has a finite subcover from among the  $]a - r_a; a + r_a[$  and so there exists an integer  $N > 0$  and  $a_i$  such that

$$A \subseteq \bigcup_{i=1}^N ]a_i - r_{a_i}; a_i + r_{a_i}[ .$$

Since

$$A = A \cap \bigcup_{i=1}^N ]a_i - r_{a_i}; a_i + r_{a_i}[ = \bigcup_{i=1}^N \{a_i\},$$

$A$  has only  $N$  elements and thus it is finite.  $\square$

**143 THEOREM** Let  $X \subseteq \mathbb{R}$ . Then the following are equivalent.

1.  $X$  is compact.
2.  $X$  is closed and bounded.
3. every infinite set of  $X$  has an accumulation point.
4. every infinite sequence of  $X$  has a converging subsequence in  $X$ .

## Homework

**Problem 2.6.1** Give an example shewing that the union of an infinite number of closed sets is not necessarily closed.

**Problem 2.6.2** Prove that a set  $A \subseteq \mathbb{R}$  is dense if and only if  $\overline{A} = \mathbb{R}$ .

**Problem 2.6.3** For any set  $A \subseteq \mathbb{R}$  prove that  $\text{Bdy}(A) = \text{Bdy}(\mathbb{R} \setminus A)$ .

**Problem 2.6.4** Let  $A \neq \emptyset$  be a subset of  $\mathbb{R}$ . Assume that  $A$  is bounded above. Prove that  $\sup(A) = \sup(\overline{A})$ .

**Problem 2.6.5** Demonstrate that the only subsets of  $\mathbb{R}$  which are simultaneously open and closed in  $\mathbb{R}$  are  $\emptyset$  and  $\mathbb{R}$ . One codifies this by saying that  $\mathbb{R}$  is connected.

**Problem 2.6.6** Prove that the closed additive subgroups of the real numbers are (i) just zero; or (ii) all integral multiples of a fixed non-

zero number (which may be assumed positive); or (iii) all reals.

**Problem 2.6.7** Let  $A \in \mathbb{R}$ . Prove the following

- |   |   |
|---|---|
| 1. $\overline{\overline{A}} = \overline{A}$                                 | 6. $\overline{A \cap B} \subseteq \overline{A} \cap \overline{B}$                   |
| 2. $\overset{\circ}{A} = A$   | 7. $\overset{\circ}{A} \cup \overset{\circ}{B} \subseteq \overset{\circ}{A \cup B}$ |
| 3. $A \subseteq B \implies \overline{A} \subseteq \overline{B}$             | 8. $\overset{\circ}{A \cap B} = \overset{\circ}{A} \cap \overset{\circ}{B}$         |
| 4. $A \subseteq B \implies \overset{\circ}{A} \subseteq \overset{\circ}{B}$ |   |
| 5. $\overline{A \cup B} = \overline{A} \cup \overline{B}$                   |   |

## 2.7 $\overline{\mathbb{R}}$

**Why bother?** The algebraic rules introduced here will simplify some computations and statements in subsequent chapters.

Geometrically, each real number can be viewed as a point on a straight line. We make the convention that we orient the real line with  $0$  as the origin, the positive numbers increasing towards the right from  $0$  and the negative numbers decreasing towards the left of  $0$ , as in figure 2.1.




Figure 2.1: The Real Line.

We append the object  $+\infty$ , which is larger than any real number, and the object  $-\infty$ , which is smaller than any real number. Letting  $x \in \mathbb{R}$ , we make the following conventions.

$$(+\infty) + (+\infty) = +\infty \quad (2.1)$$

$$(-\infty) + (-\infty) = -\infty \quad (2.2)$$

$$x + (+\infty) = +\infty \quad (2.3)$$

$$x + (-\infty) = -\infty \quad (2.4)$$

$$x(+\infty) = +\infty \quad \text{if } x > 0 \quad (2.5)$$

$$x(+\infty) = -\infty \quad \text{if } x < 0 \quad (2.6)$$

$$x(-\infty) = -\infty \quad \text{if } x > 0 \quad (2.7)$$

$$x(-\infty) = +\infty \quad \text{if } x < 0 \quad (2.8)$$

$$\frac{x}{\pm\infty} = 0 \quad (2.9)$$

Observe that we leave the following undefined:

$$\frac{\pm\infty}{\pm\infty}, \quad (+\infty) + (-\infty), \quad 0(\pm\infty).$$

**144 Definition** We denote by  $\overline{\mathbb{R}} = [-\infty; +\infty]$  the set of real numbers such with the two symbols  $-\infty$  and  $+\infty$  appended, obeying the algebraic rules above. Observe that then every set in  $\overline{\mathbb{R}}$  has a supremum (it may as well be  $+\infty$  if the set is unbounded by finite numbers) and an infimum (which may be  $-\infty$ ).

## 2.8 Lebesgue Measure

**145 Definition** Let  $(a, b) \in \mathbb{R}^2$ . The *measure* of the open interval  $]a; b[$  is  $b - a$ . We denote this by  $\mu(]a; b[) = b - a$ . If  $G = \bigcup_{k=1}^{\infty} ]a_k; b_k[$  is a union of disjoint, bounded, open intervals, then  $\mu(G) = \sum_{k=1}^{\infty} (b_k - a_k)$ .

**146 Definition** Let  $E \subseteq \mathbb{R}$  be a bounded set. The *outer measure* of  $E$  is defined and denoted by

$$\overline{\mu}(E) = \inf_{\substack{E \subseteq O \\ O \text{ open}}} \mu(O).$$

**147 Definition** A set  $E \subseteq \mathbb{R}$  is said to be *Lebesgue measurable* if  $\forall \varepsilon > 0, \exists G \supseteq E$  open such that  $\overline{\mu}(G \setminus E) < \varepsilon$ . In this case  $\mu(E) = \overline{\mu}(E)$ .

## 2.9 The Cantor Set

**148 Definition (The Cantor Set)** The Cantor set  $C$  is the canonical example of an uncountable set of measure zero. We construct  $C$  as follows.

Begin with the unit interval  $C_0 = [0; 1]$ , and remove the middle third open segment  $R_1 := ]\frac{1}{3}; \frac{2}{3}[$ . Define  $C_1$  as

$$C_1 := C_0 \setminus R_1 = [0; \frac{1}{3}] \cup [\frac{2}{3}; 1] \quad (2.10)$$

Iterate this process on each remaining segment, removing the open set

$$R_2 := ]\frac{1}{9}; \frac{2}{9}[ \cup ]\frac{7}{9}; \frac{8}{9}[ \quad (2.11)$$

to form the four-interval set

$$C_2 := C_1 \setminus R_2 = [0; \frac{1}{9}] \cup [\frac{2}{9}; \frac{1}{3}] \cup [\frac{2}{3}; \frac{7}{9}] \cup [\frac{8}{9}; 1] \quad (2.12)$$

Continue the process, forming  $C_3, C_4, \dots$ . Note that  $C_k$  has  $2^k$  pieces.

At each step, the endpoints of each closed segment will remain in the set. See figure 2.2.

The *Cantor set* is defined as

$$C := \bigcap_{k=1}^{\infty} C_k = C_0 \setminus \bigcup_{n=1}^{\infty} R_n \quad (2.13)$$

**149 THEOREM (Cardinality of the Cantor Set)** The Cantor Set is uncountable.

**Proof:** Starting with the two pieces of  $C_1$ , we mark the sinistral segment “0” and the dextral segment “1”. We then continue to  $C_2$ , and consider only the leftmost pair. Again, mark the segments “0” and “1”, and do the same for the rightmost pair. Successively then, mark the  $2^{k-1}$  leftmost segments of  $C_k$  “0” and the  $2^{k-1}$  rightmost segments “1.” The elements of the Cantor Set are those with infinite binary expansions. Since there uncountable many such expansions, the Cantor Set is uncountable.  $\square$

**150 THEOREM (Measure of the Cantor Set)** The Cantor Set has (Lebesgue) measure 0.




Figure 2.2: Construction of the Cantor Set.

**Proof:** Using the notation of Definition 148, observe that

$$\mu(R_1) = \frac{2}{3} - \frac{1}{3} = \frac{1}{3} \tag{2.14}$$

$$\mu(R_2) = \left(\frac{2}{9} - \frac{1}{9}\right) + \left(\frac{8}{9} - \frac{7}{9}\right) = \frac{2}{9} \tag{2.15}$$

$$\vdots \tag{2.16}$$

$$\mu(R_k) = \sum_{n=1}^k \frac{2^{n-1}}{3^n} \tag{2.17}$$

Since the  $R$ 's are disjoint, the measure of their union is the sum of their measures. Taking the limit as  $k \rightarrow \infty$ ,

$$\mu\left(\bigcup_{n=1}^{\infty} R_n\right) = \sum_{n=1}^{\infty} \frac{2^{n-1}}{3^n} = 1. \tag{2.18}$$

Since clearly  $\mu(C_0) = 1$ , we then have

$$\mu(C) = \mu\left(C_0 \setminus \bigcup_{n=1}^{\infty} R_n\right) = \mu(C_0) - \sum_{n=1}^{\infty} \frac{1}{2^n} = 1 - 1 = 0. \tag{2.19}$$

□

**151 THEOREM** The Cantor set is closed and its interior is empty.

**Proof:** Each of  $C_0, C_1, C_2, \dots$ , is closed, being the union of a finite number of closed intervals. Thus the Cantor Set is closed, as it is the intersection of closed sets.

Now, let  $I$  be an open interval. Since the numbers of the form  $\frac{m}{3^n}$ ,  $(m, n) \in \mathbb{Z}$  are dense in the reals, there is exists a rational number  $\frac{m}{3^n} \in I$ . Expressed in ternary, this rational number has a finite expansion. If this expansion contains the digit “1”, then this number does not belong to Cantor Set, and we are done. If not, since  $I$  is open, there must exist a number  $k > n$  such that  $\frac{m}{3^n} + \frac{1}{3^k} \in I$ . By construction, the last digit of the ternary expansion of this number is also “1”, and hence this number does not belong to the Cantor Set either. □


# Chapter 3

## Sequences

### 3.1 Limit of a Sequence

**Why bother?** The *limit* concept is at the centre of calculus. We deal with discrete quantities first, that is, with limits of sequences.

**152 Definition** A (numerical) sequence is a function  $\mathbf{a} : \mathbb{N} \rightarrow \mathbb{R}$ . We usually denote  $\mathbf{a}(n)$  by  $\mathbf{a}_n$ .<sup>1</sup>

 We will use the notation  $\{\mathbf{a}_n\}_{n=k}^l$  to denote the sequence  $\mathbf{a}_k, \mathbf{a}_{k+1}, \dots, \mathbf{a}_l$ . For example

$$\{\mathbf{a}_n\}_{n=0}^{10} = \{\mathbf{a}_0, \mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_{10}\},$$

$$\{\mathbf{b}_n\}_{n=4}^6 = \{\mathbf{b}_4, \mathbf{b}_5, \mathbf{b}_6\},$$

$$\left\{ \left(1 + \frac{1}{n}\right)^n \right\}_{n=1}^{+\infty} = \left\{ 2, \frac{9}{4}, \frac{64}{27}, \dots \right\},$$

etc.

**153 Example** The *Harmonic sequence* is

$$1, \quad \frac{1}{2}, \quad \frac{1}{3}, \quad \dots,$$

or  $\mathbf{a}_n = \frac{1}{n}$  for  $n \geq 1$ .

**154 Definition** A sequence  $\{\mathbf{a}_n\}_{n=1}^{+\infty}$  is *bounded* if there exists a constant  $K > 0$  such that  $\forall n, |\mathbf{a}_n| \leq K$ . It is *increasing* if for all  $n > 0$ ,  $\mathbf{a}_n \leq \mathbf{a}_{n+1}$  and *decreasing* if for all  $n \geq 0$ ,  $\mathbf{a}_n \geq \mathbf{a}_{n+1}$ .

### 3.2 Convergence of Sequences

**155 Definition** A sequence  $\{\mathbf{a}_n\}_{n=1}^{+\infty}$  is said to *converge* if

$$\exists L \in \mathbb{R}, \forall \varepsilon > 0, \exists N > 0 \text{ such that } \forall n \in \mathbb{N}, n \geq N \implies |\mathbf{a}_n - L| < \varepsilon.$$

In other words, eventually<sup>2</sup> the differences

$$|\mathbf{a}_n - L|, |\mathbf{a}_{n+1} - L|, |\mathbf{a}_{n+2} - L|, \dots$$

remain smaller than an arbitrarily prescribed small quantity. We denote the fact that the sequence  $\{\mathbf{a}_n\}_{n=1}^{+\infty}$  converges to  $L$  as  $n \rightarrow +\infty$  by


$$\lim_{n \rightarrow +\infty} \mathbf{a}_n = L, \text{ or by } \mathbf{a}_n \rightarrow L \text{ as } n \rightarrow +\infty.$$

<sup>1</sup>It is customary to start at  $n = 1$  rather than  $n = 0$ . We won't be too fuzzy about such complications, but we will be careful to write sense.

<sup>2</sup>A good word to use in informal speech "eventually" will mean "for large enough values" or in the case at hand  $\forall n \geq N$  for some strictly positive integer  $N$ .

A sequence that does not converge is said to *diverge*. Thus a sequence diverges if

$$\forall L \in \mathbb{R}, \exists \varepsilon > 0, \forall N \in \mathbb{N}, \exists n \in \mathbb{N} \text{ such that } n > N \text{ and } |a_n - L| \geq \varepsilon.$$

 Given a sequence  $\{a_n\}_{n=1}^{+\infty}$  and  $L \in \mathbb{R}$ ,

$$a_n \rightarrow L \text{ as } n \rightarrow +\infty \text{ if and only if } \liminf a_n = \limsup a_n = \lim a_n = L.$$

**156 Definition** A sequence  $\{b_n\}_{n=1}^{+\infty}$  *diverges to plus infinity* if  $\forall M > 0, \exists N > 0$  such that  $\forall n \geq N, b_n > M$ . A sequence  $\{c_n\}_{n=1}^{+\infty}$  *diverges to minus infinity* if  $\forall M > 0, \exists N > 0$  such that  $\forall n \geq N, c_n < -M$ . A sequence that diverges to plus or minus infinity is said to *properly diverge*. Otherwise it is said to *oscillate*.

**157 Definition** Given a sequence  $\{a_n\}_{n=1}^{+\infty}$ , we say that  $\lim_{n \rightarrow +\infty} a_n$  *exists* if it is either convergent or properly divergent.

**158 Example** The constant sequence

$$1, 1, 1, 1, \dots$$

converges to **1**. Similarly, if a sequence is eventually stationary, that is, constant, it converges to that constant.

**159 Example** Consider the sequence

$$1, \frac{1}{2}, \frac{1}{3}, \dots, \frac{1}{n}, \dots,$$

We claim that  $\frac{1}{n} \rightarrow 0$  as  $n \rightarrow +\infty$ . Suppose we wanted terms that get closer to **0** by at least  $.00001 = \frac{1}{10^5}$ . We only need to look at the **100000**-term of the sequence:  $\frac{1}{100000} = \frac{1}{10^5}$ . Since the terms of the sequence get smaller and smaller, any term after this one will be within **.00001** of **0**. We had to wait a long time—till after the **100000**-th term—but the sequence eventually did get closer than **.00001** to **0**. The same argument works for any distance, no matter how small, so we can eventually get arbitrarily close to **0**. A rigorous proof is as follows. If  $\varepsilon > 0$  is no matter how small, we need only to look at the terms after  $N = \lfloor \frac{1}{\varepsilon} + 1 \rfloor$  to see that, indeed, if  $n > N$ , then

$$s_n = \frac{1}{n} < \frac{1}{N} = \frac{1}{\lfloor \frac{1}{\varepsilon} + 1 \rfloor} < \varepsilon.$$

Here we have used the inequality

$$t - 1 < \lfloor t \rfloor \leq t, \quad \forall t \in \mathbb{R}.$$

**160 Example** The sequence

$$0, 1, 4, 9, 16, \dots, n^2, \dots$$

diverges to  $+\infty$ , as the sequence gets arbitrarily large. A rigorous proof is as follows. If  $M > 0$  is no matter how large, then the terms after  $N = \lfloor \sqrt{M} \rfloor + 1$  satisfy ( $n > N$ )

$$t_n = n^2 > N^2 = (\lfloor \sqrt{M} \rfloor + 1)^2 > M.$$

**161 Example** The sequence

$$1, -1, 1, -1, 1, -1, \dots, (-1)^n, \dots$$

has no limit (diverges), as it bounces back and forth from  $-1$  to  $+1$  infinitely many times.

**162 Example** The sequence

$$0, -1, 2, -3, 4, -5, \dots, (-1)^n n, \dots,$$

has no limit (diverges), as it is unbounded and alternates back and forth positive and negative values..

We will now see some properties of limits of sequences.



**163 THEOREM (Uniqueness of Limits)** If  $a_n \rightarrow L$  and  $a_n \rightarrow L'$  as  $n \rightarrow +\infty$  then  $L = L'$ .

**Proof:** The statement only makes sense if both  $L$  and  $L'$  are finite, so assume so. If  $L \neq L'$ , take  $\varepsilon = \frac{|L - L'|}{2} > 0$  in the definition of convergence. Now

$$\lim_{n \rightarrow +\infty} a_n = L \implies \exists N_1 > 0, \quad \forall n \geq N_1 |a_n - L| < \varepsilon,$$

$$\lim_{n \rightarrow +\infty} a_n = L' \implies \exists N_2 > 0, \quad \forall n \geq N_2 |a_n - L'| < \varepsilon.$$

If  $n > \max(N_1, N_2)$  then by the Triangle Inequality (Theorem 76) then

$$|L - L'| \leq |L - a_n| + |a_n - L'| < 2\varepsilon = |L - L'|,$$

a contradiction, so  $L = L'$ .  $\square$

**164 THEOREM** Every convergent sequence is bounded.

**Proof:** Let  $\{a_n\}_{n=1}^{+\infty}$  converge to  $L$ . Using  $\varepsilon = 1$  in the definition of convergence,  $\exists N > 0$  such that

$$n \geq N \implies |a_n - L| < 1 \implies L - 1 < a_n < L + 1,$$

hence, eventually,  $a_n$  does not exceed  $L + 1$ .  $\square$




Figure 3.1: Theorem ??.

When is it guaranteed that a sequence of real numbers has a limit? We have the following result.

**165 THEOREM** Every bounded increasing sequence  $\{a_n\}_{n=0}^{+\infty}$  of real numbers converges to its supremum. Similarly, every bounded decreasing sequence of real numbers converges to its infimum.

**Proof:** The idea of the proof is sketched in figure 3.1. By virtue of Axiom 92, the sequence has a supremum  $s$ . Every term of the sequence satisfies  $a_n \leq s$ . We claim that eventually all the terms of the sequence are closer to  $s$  than a preassigned small distance  $\varepsilon > 0$ . Since  $s - \varepsilon$  is not an upper bound for the sequence, there must be a term of the sequence, say  $a_{n_0}$  with  $s - \varepsilon \leq a_{n_0}$  by virtue of the Approximation Property Theorem 93. Since the sequence is increasing, we then have

$$s - \varepsilon \leq a_{n_0} \leq a_{n_0+1} \leq a_{n_0+2} \leq a_{n_0+2} \leq \dots \leq s,$$

which means that after the  $n_0$ -th term, we get to within  $\varepsilon$  of  $s$ .

To obtain the second half of the theorem, we simply apply the first half to the sequence  $\{-a_n\}_{n=0}^{+\infty}$ .  $\square$

**166 THEOREM (Order Properties of Sequences)** Let  $\{a_n\}_{n=1}^{+\infty}$  be a sequence of real numbers converging to the real number  $L$ . Then

1. If  $a < L$  then eventually  $a < a_n$ .
2. If  $L < b$  then eventually  $a_n < b$ .
3. If  $a < L < b$  then eventually  $a < a_n < b$ .
4. If eventually  $a_n \geq a$  then  $L \geq a$ .

5. If eventually  $\mathbf{a}_n \leq \mathbf{b}$  then  $L \leq \mathbf{b}$ .
6. If eventually  $\mathbf{a} \leq \mathbf{a}_n \leq \mathbf{b}$  then  $\mathbf{a} \leq L \leq \mathbf{b}$ .

**Proof:** We apply the definition of convergence repeatedly.

1. Taking  $\varepsilon = L - \mathbf{a}$  in the definition of convergence,  $\exists N_1 > \mathbf{0}$  such that

$$\forall n \geq N_1, \quad |\mathbf{a}_n - L| < L - \mathbf{a} \implies \forall n \geq N_1, \quad \mathbf{a} - L < \mathbf{a}_n - L < L - \mathbf{a} \implies \forall n \geq N_1, \quad \mathbf{a} < \mathbf{a}_n,$$

that is, eventually  $\mathbf{a} < \mathbf{a}_n$ .

2. Taking  $\varepsilon = \mathbf{b} - L$  in the definition of convergence,  $\exists N_2 > \mathbf{0}$  such that

$$\forall n \geq N_2, \quad |\mathbf{a}_n - L| < \mathbf{b} - L \implies \forall n \geq N_2, \quad L - \mathbf{b} < \mathbf{a}_n - L < \mathbf{b} - L \implies \forall n \geq N_2, \quad \mathbf{a}_n < \mathbf{b},$$

that is, eventually  $\mathbf{a}_n < \mathbf{b}$ .

3. It suffices to take  $N = \max(N_1, N_2)$  above.
4. If, to the contrary,  $L > \mathbf{a}$ , then by part (1) we will eventually have  $\mathbf{a}_n > \mathbf{a}$ , a contradiction.
5. If, to the contrary,  $L < \mathbf{b}$ , then by part (2) we will eventually have  $\mathbf{a}_n < \mathbf{b}$ , a contradiction.
6. If either  $L < \mathbf{a}$  or  $\mathbf{b} < L$  we would obtain a contradiction to parts (4) or (5).

□

**167 THEOREM (Sandwich Theorem)** Let  $\{\mathbf{a}_n\}_{n=1}^{+\infty}$ ,  $\{\mathbf{u}_n\}_{n=1}^{+\infty}$ ,  $\{\mathbf{v}_n\}_{n=1}^{+\infty}$  be sequences of real numbers such that eventually

$$\mathbf{u}_n \leq \mathbf{a}_n \leq \mathbf{v}_n.$$

If for  $L \in \mathbb{R}$ ,  $\mathbf{u}_n \rightarrow L$  and  $\mathbf{v}_n \rightarrow L$  then  $\mathbf{a}_n \rightarrow L$ .

**Proof:** For all  $\varepsilon > \mathbf{0}$  there are  $N_1 > \mathbf{0}$ ,  $N_2 > \mathbf{0}$  such that

$$\forall n \geq \max(N_1, N_2), \quad |\mathbf{u}_n - L| < \varepsilon, \quad |\mathbf{v}_n - L| < \varepsilon \implies -\varepsilon < \mathbf{u}_n - L < \varepsilon, \quad -\varepsilon < \mathbf{v}_n - L < \varepsilon.$$

Thus for such  $n$ ,

$$-\varepsilon < \mathbf{u}_n - L \leq \mathbf{a}_n - L \leq \mathbf{v}_n - L < \varepsilon, \implies -\varepsilon < \mathbf{a}_n - L < \varepsilon \implies |\mathbf{a}_n - L| < \varepsilon,$$

from where  $\{\mathbf{a}_n\}_{n=1}^{+\infty}$  converges to  $L$ . □

**168 THEOREM** Let  $\{\mathbf{a}_n\}_{n=1}^{+\infty}$  be a sequence of real numbers such that  $\mathbf{a}_n \rightarrow L$ . Then  $|\mathbf{a}_n| \rightarrow |L|$ .

**Proof:** From Corollary 77, we have the inequality  $||\mathbf{a}_n| - |L|| \leq |\mathbf{a}_n - L|$  from where the result follows. □

**169 THEOREM** Let  $\{\mathbf{a}_n\}_{n=1}^{+\infty}$  be a sequence of real numbers such that  $\mathbf{a}_n \rightarrow \mathbf{0}$ , and let  $\{\mathbf{b}_n\}_{n=1}^{+\infty}$  be a bounded sequence. Then  $\mathbf{a}_n \mathbf{b}_n \rightarrow \mathbf{0}$ .

**Proof:** Eventually  $|\mathbf{a}_n| < \varepsilon$ . Assume that eventually  $|\mathbf{b}_n| \leq U$ . Then

$$|\mathbf{a}_n \mathbf{b}_n| \leq U |\mathbf{a}_n| < U \varepsilon,$$

from where the result follows. □

**170 THEOREM** If  $\mathbf{b}_n \rightarrow l \neq \mathbf{0}$  then  $\mathbf{b}_n$  is eventually different from  $\mathbf{0}$  and  $\frac{1}{\mathbf{b}_n} \rightarrow \frac{1}{l}$ .

**Proof:** By Theorem 169,  $|\mathbf{b}_n| \rightarrow |\mathbf{l}|$ . Using  $\varepsilon = \frac{|\mathbf{l}|}{2} > 0$  in the definition of convergence, we have that eventually

$$||\mathbf{b}_n| - |\mathbf{l}|| < \frac{|\mathbf{l}|}{2} \implies |\mathbf{l}| - \frac{|\mathbf{l}|}{2} < |\mathbf{b}_n| < |\mathbf{l}| + \frac{|\mathbf{l}|}{2} \implies \frac{|\mathbf{l}|}{2} < |\mathbf{b}_n|,$$

That is, eventually  $|\mathbf{b}_n|$  is strictly positive and so  $\frac{1}{\mathbf{b}_n}$  makes sense. Also, eventually,  $\frac{1}{|\mathbf{b}_n|} < \frac{2}{|\mathbf{l}|}$ . Now, for sufficiently large  $n$ ,

$$\left| \frac{1}{\mathbf{b}_n} - \frac{1}{\mathbf{l}} \right| = \left| \frac{\mathbf{l} - \mathbf{b}_n}{|\mathbf{b}_n| |\mathbf{l}|} \right| = \frac{|\mathbf{b}_n - \mathbf{l}|}{|\mathbf{b}_n| |\mathbf{l}|} < \frac{2\varepsilon}{|\mathbf{l}| |\mathbf{l}|},$$

from where the result follows.  $\square$

**171 THEOREM (Algebraic Properties of Sequences)** Let  $k \in \mathbb{R}$ . If  $\{a_n\}_{n=1}^{+\infty}$  converges to  $L$  and  $\{b_n\}_{n=1}^{+\infty}$  converges to  $L'$  then

$$\lim_{n \rightarrow +\infty} (ka_n + b_n) = kL + L', \quad \lim_{n \rightarrow +\infty} (a_n b_n) = LL'.$$

Moreover, if  $L' \neq 0$  then

$$\lim_{n \rightarrow +\infty} \left( \frac{a_n}{b_n} \right) = \frac{L}{L'}.$$

**Proof:** The trick in all these proofs is the following observation: If one multiplies a bounded quantity by an arbitrarily small quantity, one gets an arbitrarily small quantity. Hence one considers the absolute value of the desired terms of the sequence from the expected limit.

Given  $\varepsilon > 0$  there are  $N_1 > 0$  and  $N_2 > 0$  such that  $|a_n - L| < \varepsilon$  and  $|b_n - L'| < \varepsilon$ . Then

$$|(ka_n + b_n) - (kL + L')| = |(ka_n - kL) + (b_n - L')| \leq |k| |a_n - L| + |b_n - L'| < \varepsilon(k + 1),$$

and so the sinistral side is arbitrarily close to  $0$ , establishing the first assertion.

For the product, observe that by Theorem 164 there exists a constant  $K > 0$  such that  $|b_n| < K$ . Hence

$$|a_n b_n - LL'| = |(a_n - L)b_n + L(b_n - L')| \leq |a_n - L| |b_n| + |L| |b_n - L'| < \varepsilon K + |L| \varepsilon = \varepsilon(K + |L|),$$

and again, the sinistral side is made arbitrarily close to  $0$ .

Finally, if  $L' \neq 0$  then by Theorem 170,  $b_n$  is eventually  $\neq 0$  and  $\frac{1}{b_n} \rightarrow \frac{1}{L'}$ . We now simply apply the result we obtained for products, giving

$$a_n b_n \rightarrow L \left( \frac{1}{L'} \right) = \frac{L}{L'}.$$

$\square$

## Homework

**Problem 3.2.1** If  $\forall n > 0$ ,  $a_n > 0$  and  $\{a_n\}_{n=1}^{+\infty}$  converges to  $L$  must it be the case that  $L > 0$ ?

**Problem 3.2.2** Prove that if  $a_n \rightarrow +\infty$  and if  $\{b_n\}_{n=1}^{+\infty}$  is bounded, then  $a_n + b_n \rightarrow +\infty$ .

**Problem 3.2.3** Prove that if  $a_n \rightarrow +\infty$  and  $b_n \rightarrow +\infty$  is bounded, then  $a_n + b_n \rightarrow +\infty$ .

**Problem 3.2.4** Prove that if  $a_n \rightarrow +\infty$  and if there exists  $K > 0$  such that eventually  $b_n \geq K$ , then  $a_n b_n \rightarrow +\infty$ .

**Problem 3.2.5** Prove that if  $a_n \rightarrow +\infty$  and  $b_n \rightarrow +\infty$  is bounded, then  $a_n b_n \rightarrow +\infty$ .

**Problem 3.2.6** Prove that if  $a_n \rightarrow +\infty$  and if  $\{b_n\}_{n=1}^{+\infty}$  is bounded, then  $a_n + b_n \rightarrow +\infty$ .

**Problem 3.2.7** Prove that if  $a_n \rightarrow +\infty$  then  $\frac{1}{a_n} \rightarrow 0$ .

**Problem 3.2.8** Prove that if  $a_n \rightarrow 0$  and if eventually  $a_n > 0$ , then  $\frac{1}{a_n} \rightarrow +\infty$ .

**Problem 3.2.9** Prove that  $\sum_{i=1}^n \frac{n}{n^2+i} \rightarrow 1$  as  $n \rightarrow +\infty$ .

**Problem 3.2.10** Prove that  $\frac{1}{(n!)^{1/n}} \rightarrow 0$ .

**Problem 3.2.11** Prove that  $\frac{2^n}{n!} \rightarrow 0$ .

**Problem 3.2.12** Let  $x_1, x_2, \dots$  be a bounded sequence of real numbers, and put  $s_n = x_1 + x_2 + \dots + x_n$ . Suppose that  $\frac{s_n^2}{n^2} \rightarrow 0$ . Prove that  $\frac{s_n}{n} \rightarrow 0$ .

**Problem 3.2.13** Prove rigorously that the sequence  $\{\sin n\}_{n=0}^{+\infty}$  is divergent.

**Problem 3.2.14** Prove that  $(n!)^{1/n} \rightarrow +\infty$  as  $n \rightarrow +\infty$ .

**Problem 3.2.15** A sequence of real numbers  $a_1, a_2, \dots$  satisfies, for all  $m, n$ , the inequality

$$|a_m + a_n - a_{m+n}| \leq \frac{1}{m+n}.$$

Prove that this sequence is an arithmetic progression.

**Problem 3.2.16** Prove rigorously that  $\sqrt{n+1} - \sqrt{n} \rightarrow 0$  as  $n \rightarrow +\infty$ .

**Problem 3.2.17** Prove that the sequence  $H_n = 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n}$  diverges to  $+\infty$ .

**Problem 3.2.18** Find

$$\lim_{K \rightarrow +\infty} \sum_{n=1}^K \frac{\sqrt{(n-1)!}}{(1+\sqrt{1})(1+\sqrt{2})(1+\sqrt{3}) \cdots (1+\sqrt{n})}.$$

**Problem 3.2.19** What reasonable meaning can be given to

$$\sqrt{1 + \sqrt{1 + \sqrt{1 + \sqrt{\dots}}}} \quad ?$$

**Problem 3.2.20** Prove that

$$\frac{1+2+\dots+n}{n^2} \rightarrow \frac{1}{2}, \text{ as } n \rightarrow +\infty.$$

**Problem 3.2.21** Calculate the following limits:

$$1. \lim_{n \rightarrow +\infty} \left( \frac{1^2}{n^2} + \frac{2^2}{n^2} + \dots + \frac{(n-1)^2}{n^2} \right),$$

$$2. \lim_{n \rightarrow +\infty} \left( \frac{1}{1 \cdot 2} + \frac{1}{2 \cdot 3} + \dots + \frac{1}{n(n+1)} \right),$$

$$3. \lim_{n \rightarrow +\infty} \left( \frac{1}{1 \cdot 2 \cdot 3} + \frac{1}{2 \cdot 3 \cdot 4} + \dots + \frac{1}{n(n+1)(n+2)} \right),$$

**Problem 3.2.22** What reasonable meaning can be given to

$$\frac{1}{1 + \frac{1}{1 + \frac{1}{1 + \frac{1}{\ddots}}}} \quad ?$$

**Problem 3.2.23** Let  $K \in \mathbb{N} \setminus \{0\}$ , and let  $a_1, \dots, a_K, \lambda_1, \dots, \lambda_K$  be strictly positive real numbers. Prove that

$$\lim_{n \rightarrow +\infty} \left( \sum_{k=1}^K \lambda_k a_k^n \right)^{1/n} = \max_{1 \leq k \leq K} a_k, \quad \lim_{n \rightarrow +\infty} \left( \sum_{k=1}^K \lambda_k a_k^{-n} \right)^{-1/n} = \min_{1 \leq k \leq K} a_k.$$

**Problem 3.2.24** Prove that if  $\left\{ \frac{a_n}{b_n} \right\}_{n=1}^{+\infty}$  is a monotonic sequence,

then the  $\left\{ \frac{a_1 + a_2 + \dots + a_n}{b_1 + b_2 + \dots + b_n} \right\}_{n=1}^{+\infty}$  is also monotonic in the same sense.

**Problem 3.2.25** Let  $a, b, c$  be real numbers such that  $b^2 - 4ac < 0$ . Let  $\{X_n\}_{n=1}^{+\infty}, \{Y_n\}_{n=1}^{+\infty}$  be sequences of real numbers such that

$$aX_n^2 + bX_n Y_n + cY_n^2 \rightarrow 0, \text{ as } n \rightarrow +\infty.$$

Prove that  $X_n \rightarrow 0$  and  $Y_n \rightarrow 0$  as  $n \rightarrow +\infty$ .

**Problem 3.2.26 (Gram's Product)** Prove that

$$\lim_{n \rightarrow +\infty} \prod_{k=2}^n \frac{k^3 - 1}{k^3 + 1} = \frac{2}{3}.$$

**Problem 3.2.27** Prove that the sequence  $\{x_n\}_{n=1}^{+\infty}$  with  $x_n = 1 + \frac{1}{2^2} + \frac{1}{3^2} + \dots + \frac{1}{n^2}$  satisfies  $x_n \leq 2 - \frac{1}{n}$  for  $n \geq 1$ . Hence deduce that it converges.

**Problem 3.2.28** Prove the convergence of the sequence  $x_n = \sum_{k=1}^n \frac{1}{n+k}, n \geq 1$ .

**Problem 3.2.29** Prove the convergence of the sequence,  $x_1 = a, x_2 = b, x_{n+1} = \frac{x_n + x_{n-1}}{2}, n \geq 2$  and  $(a, b) \in \mathbb{R}^2, a \neq b$ . Also, find its limit.

**Problem 3.2.30** Prove the convergence of the sequence,  $x_1 = a, x_{n+1} = \frac{1}{2} \left( x_n + \frac{b}{x_n} \right), n \geq 1$  and  $(a, b) \in \mathbb{R}^2, a > 0, b > 0$ . Also, find its limit.

**Problem 3.2.31** Prove the convergence of the sequence,  $x_1 = a, x_{n+1} = \frac{1}{2} \left( x_n + \frac{b}{x_n} \right), n \geq 1$  and  $(a, b) \in \mathbb{R}^2, a < 0, b > 0$ . Also, find its limit.

**Problem 3.2.32** Let  $(a, b) \in \mathbb{R}^2$ ,  $a > b > 0$ . Set  $a_1 = \frac{a+b}{2}$ ,  $b_1 = \sqrt{ab}$ . If for  $n > 1$ ,

$$a_{n+1} = \frac{a_n + b_n}{2}, \quad b_{n+1} = \sqrt{a_n b_n},$$

Prove that

1.  $\{a_n\}_{n=1}^{+\infty}$  is monotonically decreasing,
2.  $\{b_n\}_{n=1}^{+\infty}$  is monotonically increasing,
3. both sequences converge,
4. their limits are equal.

### 3.3 Classical Limits of Sequences

**Why bother?** In this section we will obtain various classical limits. In particular, we define the constant  $e$  and obtain a few interesting results about it.

**172 THEOREM** Let  $r \in \mathbb{R}$  be fixed. If  $|r| < 1$  then  $r^n \rightarrow 0$  as  $n \rightarrow +\infty$ . If  $|r| > 1$  then  $r^n \rightarrow +\infty$  as  $n \rightarrow +\infty$ .

**Proof:** Taking  $x = \left| \frac{1}{r} \right| - 1$  in Bernoulli's Inequality (Theorem 81), we find

$$\left| \frac{1}{r} \right|^n > 1 + n \left( \left| \frac{1}{r} \right| - 1 \right) > n \left( \left| \frac{1}{r} \right| - 1 \right).$$

Therefore

$$|r|^n < \frac{|r|}{n(1-|r|)} \rightarrow 0,$$

as  $n \rightarrow +\infty$ , since  $\frac{1}{n} \rightarrow 0$  as  $n \rightarrow +\infty$ .

If  $|r| > 1$ , again by Bernoulli's Inequality

$$|r|^n = (1 + |r| - 1)^n > 1 + n(|r| - 1),$$

and the dextral side can be made arbitrarily large.  $\square$


**173 THEOREM** Let  $|r| < 1$ . Then

$$1 + r + r^2 + \cdots + r^n \rightarrow \frac{1}{1-r}, \quad \text{as } n \rightarrow +\infty.$$

**Proof:** If  $S_n = 1 + r + r^2 + \cdots + r^n$  then  $rS_n = r + r^2 + r^3 + \cdots + r^{n+1}$  and

$$S_n - rS_n = 1 - r^{n+1} \implies S_n = \frac{1 - r^{n+1}}{1 - r}.$$

Then apply Theorem 172.  $\square$

 An estimating trick that we will use often is the following. If  $0 < r < 1$  then the truncated sum is smaller than the infinite sum and so, for all positive integers  $k$ :

$$1 + r + r^2 + \cdots + r^k < 1 + r + r^2 + \cdots = \frac{1}{1-r}.$$

**174 THEOREM** Let  $a \in \mathbb{R}$ ,  $a > 0$ , be fixed. Then  $a^{1/n} \rightarrow 1$  as  $n \rightarrow +\infty$ .

**Proof:** If  $a > 1$  then  $a^{1/n} > 1$  and by Bernoulli's Inequality,

$$a = (1 + (a^{1/n} - 1))^n > 1 + n(a^{1/n} - 1) \implies 0 \leq a^{1/n} - 1 < \frac{a-1}{n},$$

whence  $a^{1/n} - 1 \rightarrow 0$  as  $n \rightarrow +\infty$ .

If  $0 < a < 1$  then  $b = \frac{1}{a} > 1$  and so by what we just proved,

$$b^{1/n} \rightarrow 1 \implies \frac{1}{a^{1/n}} \rightarrow 1 \implies a^{1/n} \rightarrow 1,$$

proving the theorem.  $\square$

**175 THEOREM** Let  $a \in \mathbb{R}$ ,  $a > 1$ ,  $k \in \mathbb{N} \setminus \{0\}$ , be fixed. Then  $\frac{a^n}{n^k} \rightarrow +\infty$  as  $n \rightarrow +\infty$ .


**Proof:** Observe that  $a^{1/k} > 1$ . We have, using the Binomial Theorem,

$$(a^{1/k})^n = (1 + (a^{1/k} - 1))^n = \sum_{i=0}^n \binom{n}{i} (a^{1/k} - 1)^i.$$

Since each term of the above expansion is  $\geq 0$ , we gather that

$$(a^{1/k})^n \geq \frac{n(n-1)}{2} (a^{1/k} - 1)^2 \implies \frac{(a^{1/k})^n}{n} \geq \frac{(n-1)}{2} (a^{1/k} - 1)^2 \implies \frac{(a^{1/k})^n}{n} \rightarrow +\infty \implies \frac{a^n}{n^k} \rightarrow +\infty,$$

as desired.  $\square$

 In particular  $\frac{2^n}{n} \rightarrow +\infty$  as  $n \rightarrow +\infty$ .

**176 THEOREM** Let  $a \in \mathbb{R}$ , , be fixed. Then  $\frac{a^n}{n!} \rightarrow 0$  as  $n \rightarrow +\infty$ .

**Proof:** Put  $N = \lfloor |a| \rfloor + 1$  and let  $n \geq N$ . Then

$$\left| \frac{a^n}{n!} \right| = \left( \frac{|a|}{1} \cdot \frac{|a|}{2} \cdots \frac{|a|}{N} \right) \left( \frac{|a|}{N+1} \cdot \frac{|a|}{N+2} \cdots \frac{|a|}{n} \right) \leq \left( \frac{|a|^N}{N!} \right) \left( 1 \cdot 1 \cdots 1 \cdot \frac{|a|}{n} \right) \rightarrow 0,$$

as  $n \rightarrow +\infty$ .  $\square$

**177 THEOREM** The sequence

$$e_n = \left( 1 + \frac{1}{n} \right)^n, n = 1, 2, \dots$$

is a bounded increasing sequence, and hence it converges to a limit, which we call  $e$ . Also, for all strictly positive integers  $n$ ,  $\left( 1 + \frac{1}{n} \right)^n < e$ .

**Proof:** By Theorem 80

$$\frac{b^{n+1} - a^{n+1}}{b - a} \leq (n+1)b^n \implies b^n [(n+1)a - nb] < a^{n+1}.$$


Putting  $a = 1 + \frac{1}{n+1}$ ,  $b = 1 + \frac{1}{n}$  we obtain

$$e_n = \left( 1 + \frac{1}{n} \right)^n < \left( 1 + \frac{1}{n+1} \right)^{n+1} = e_{n+1},$$

whence the sequence  $e_n$ ,  $n = 1, 2, \dots$  increases. Again, by putting  $a = 1$ ,  $b = 1 + \frac{1}{2n}$  we obtain

$$\left( 1 + \frac{1}{2n} \right)^n < 2 \implies \left( 1 + \frac{1}{2n} \right)^{2n} < 4 \implies e_{2n} < 4.$$

Since  $e_n < e_{2n} < 4$  for all  $n$ , the sequence is bounded above. In view of Theorem 165 the sequence converges to a limit. We call this limit  $e$ . It follows also from this proof and from Theorem 166 that for all strictly positive integers  $n$ ,  $\left( 1 + \frac{1}{n} \right)^n < e$ .  $\square$

 Another way of obtaining  $\left(1 + \frac{1}{n}\right)^n < \left(1 + \frac{1}{n+1}\right)^{n+1}$  is as follows. Using the AM-GM Inequality with  $x_1 = 1, x_2 = \dots = x_{n+1} = 1 + \frac{1}{n}$  we have

$$\left(1 + \frac{1}{n}\right)^{n/(n+1)} < \frac{1+n\left(1 + \frac{1}{n}\right)}{n+1} \Rightarrow \left(1 + \frac{1}{n}\right)^{n/(n+1)} < \frac{n+2}{n+1} = \left(1 + \frac{1}{n+1}\right)$$

from where the desired inequality is obtained.

**178 THEOREM** The sequence  $\left\{\left(1 + \frac{1}{n}\right)^{n+1}\right\}_{n=1}^{+\infty}$  is strictly decreasing and  $\left(1 + \frac{1}{n}\right)^{n+1} \rightarrow e$ . Also, for all strictly positive integers  $n$ ,  $\left(1 + \frac{1}{n}\right)^{n+1} > e$ .

**Proof:** By Theorem 80

$$\frac{b^{n+1} - a^{n+1}}{b - a} \geq (n+1)a^n.$$

Putting  $a = 1 + \frac{1}{n+1}$ ,  $b = 1 + \frac{1}{n}$  we obtain

$$\left(1 + \frac{1}{n}\right)^{n+1} > \left(1 + \frac{1}{n+1}\right)^{n+2} \left(\frac{n^3 + 4n^2 + 4n + 1}{n(n+2)^2}\right).$$

The result will follow as long as  $\left(\frac{n^3 + 4n^2 + 4n + 1}{n(n+2)^2}\right) > 1$ . But

$$n(n+2)^2 = n(n^2 + 4n + 4) = n^3 + 4n^2 + 4n < n^3 + 4n^2 + 4n + 1 \Rightarrow \frac{n^3 + 4n^2 + 4n + 1}{n(n+2)^2} > 1.$$


Thus the sequence is a sequence of strictly decreasing sequence of real numbers. Putting  $a = 1$ ,  $b = 1 + \frac{1}{n}$  in  $\frac{b^{n+1} - a^{n+1}}{b - a} \geq (n+1)a^n$  we get

$$\left(1 + \frac{1}{n}\right)^{n+1} > 1 + n(n+1) > 2,$$

so the sequence is bounded below. In view of Theorem 165 the sequence converges to a limit  $L$ . To see that  $L = e$  observe that

$$\left(1 + \frac{1}{n}\right)^{n+1} = \left(1 + \frac{1}{n}\right)^n \left(1 + \frac{1}{n}\right) \rightarrow e \cdot 1 = e.$$

It follows also from this proof and from Theorem 166 that for all strictly positive integers  $n$ ,  $\left(1 + \frac{1}{n}\right)^{n+1} > e$ .  $\square$

 The inequality  $\left(1 + \frac{1}{n+1}\right)^{n+2} < \left(1 + \frac{1}{n}\right)^{n+1}$  can be obtained by the Harmonic Mean-Geometric Mean Inequality by putting  $x_1 = 1, x_2 = x_2 = \dots = x_{n+2} = 1 + \frac{1}{n}$

$$\frac{n+2}{\frac{1}{x_1} + \frac{1}{x_2} + \dots + \frac{1}{x_{n+2}}} \leq (x_1 x_2 \dots x_{n+2})^{1/(n+2)} \Rightarrow \frac{n+2}{1 + (n+1)\left(\frac{1}{n+1}\right)} < \left(1 + \frac{1}{n}\right)^{(n+1)/(n+2)}.$$

**179 THEOREM**  $2 < e < 3$ .

**Proof:** By the Binomial Theorem

$$\left(1 + \frac{1}{n}\right)^n = \sum_{k=0}^n \binom{n}{k} \cdot \frac{1}{n^k}.$$

Now, for  $2 \leq k \leq n$ ,

$$\binom{n}{k} \cdot \frac{1}{n^k} = \frac{1}{k!} \cdot \frac{n(n-1)(n-2)\cdots(n-k+1)}{n^k} = \frac{1}{k!} \cdot (1) \cdot \left(1 - \frac{1}{n}\right) \left(1 - \frac{2}{n}\right) \cdots \left(1 - \frac{k-1}{n}\right) \leq \frac{1}{2 \cdot 3 \cdots k} \leq \frac{1}{2^{k-1}}.$$

Thus

$$\left(1 + \frac{1}{n}\right)^n = \sum_{k=0}^n \binom{n}{k} \cdot \frac{1}{n^k} \leq 1 + 1 + \frac{1}{2} + \frac{1}{4} + \cdots + \frac{1}{2^{n-1}} < 1 + 1 + \frac{1}{2} + \frac{1}{4} + \cdots + \frac{1}{2^{n-1}} + \cdots < 1 + 2 = 3,$$

by Theorem 173 (with  $r = \frac{1}{2}$ ), and so the dextral inequality is proved. The sinistral inequality follows from Theorem 177.  $\square$



$$e = 2.718281828459045235360287471352\dots$$

**180 THEOREM**  $e = \lim_{n \rightarrow +\infty} \left(1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \cdots + \frac{1}{n!}\right).$

**Proof:** Put  $y_k = 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \cdots + \frac{1}{k!}$ . Clearly  $y_{k+1} > y_k$  so that  $\{y_k\}_{k=1}^{+\infty}$  is an increasing sequence. We will prove that it is bounded above with supremum  $e$ . By the Binomial Theorem

$$\left(1 + \frac{1}{n}\right)^n = \sum_{j=0}^n \binom{n}{j} \cdot \frac{1}{n^j} = 1 + \binom{n}{1} \frac{1}{n} + \cdots + \binom{n}{k} \frac{1}{n^k} + \cdots + \binom{n}{n} \frac{1}{n^n} \geq 1 + \binom{n}{1} \frac{1}{n} + \cdots + \binom{n}{k} \frac{1}{n^k},$$

for  $0 < k < n$ . Now let  $j$  be fixed,  $0 < j < n$ . Taking limits as  $n \rightarrow +\infty$ ,

$$\binom{n}{j} \cdot \frac{1}{n^j} = \frac{1}{j!} \cdot \frac{n(n-1)(n-2)\cdots(n-k+1)}{n^j} = \frac{1}{j!} \cdot (1) \cdot \left(1 - \frac{1}{n}\right) \left(1 - \frac{2}{n}\right) \cdots \left(1 - \frac{j-1}{n}\right) \Rightarrow \lim_{n \rightarrow +\infty} \binom{n}{j} \cdot \frac{1}{n^j} = \frac{1}{j!}.$$

Hence, taking limits as  $n \rightarrow +\infty$ ,

$$\left(1 + \frac{1}{n}\right)^n \geq 1 + \binom{n}{1} \frac{1}{n} + \cdots + \binom{n}{k} \frac{1}{n^k} \Rightarrow e \geq 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \cdots + \frac{1}{k!} = y_k,$$

or renaming,

$$e \geq 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \cdots + \frac{1}{n!} = y_n. \quad (3.1)$$

Moreover, since  $\binom{n}{k} \cdot \frac{1}{n^k} = \frac{1}{k!} \cdot (1) \cdot \left(1 - \frac{1}{n}\right) \left(1 - \frac{2}{n}\right) \cdots \left(1 - \frac{k-1}{n}\right) \leq \frac{1}{2 \cdot 3 \cdots k} \leq \frac{1}{k!}$ , we have

$$\begin{aligned} \left(1 + \frac{1}{n}\right)^n &= 1 + \binom{n}{1} \frac{1}{n} + \cdots + \binom{n}{k} \frac{1}{n^k} + \cdots + \binom{n}{n} \frac{1}{n^n} \\ &\leq 1 + \frac{1}{1!} + \cdots + \frac{1}{k!} + \cdots + \frac{1}{n!} \\ &= y_n. \end{aligned} \quad (3.2)$$

In conclusion, from 3.1 and 3.2 we get

$$\left(1 + \frac{1}{n}\right)^n \leq y_n \leq e,$$

and by taking limits and using the Sandwich Theorem, we get that  $y_n \rightarrow e$  as  $n \rightarrow +\infty$ .  $\square$

**181 LEMMA** Let  $n, m$  be strictly positive integers and let  $1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \cdots + \frac{1}{n!} = y_n$ . Then  $y_{m+n} - y_n < \frac{1}{n \cdot n!}$ .



**Proof:** We have

$$\begin{aligned}
 y_{m+n} - y_n &= \frac{1}{(n+1)!} + \frac{1}{(n+2)!} + \frac{1}{(n+3)!} + \cdots + \frac{1}{(n+m)!} \\
 &< \frac{1}{(n+1)!} \left( 1 + \frac{1}{n+2} + \frac{1}{(n+2)^2} + \cdots + \frac{1}{(n+2)^{m-1}} \right) \\
 &< \frac{1}{(n+1)!} \left( 1 + \frac{1}{n+2} + \frac{1}{(n+2)^2} + \cdots + \cdots \right) \\
 &= \frac{1}{(n+1)!} \left( \frac{1}{1 - \frac{1}{n+2}} \right) \\
 &= \frac{1}{(n+1)!} \cdot \frac{n+2}{n+1}.
 \end{aligned}$$

Here the second inequality follows by using the estimating trick deduced from Theorem 173. Observe that this bound is independent of  $m$ .  $\square$

**182 LEMMA** Let  $1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \cdots + \frac{1}{n!} = y_n$ . Then  $0 < e - y_n < \frac{1}{n!n}$ .

**Proof:** From Lemma 181,

$$0 < y_{m+n} - y_n < \frac{1}{(n+1)!} \cdot \frac{n+2}{n+1}.$$

Taking the limit as  $m \rightarrow +\infty$  we deduce

$$0 < e - y_n \leq \frac{1}{(n+1)!} \cdot \frac{n+2}{n+1}.$$

(The first inequality is strict by Theorem 180.) We only need to shew that for integer  $n \geq 1$

$$\frac{1}{(n+1)!} \cdot \frac{n+2}{n+1} < \frac{1}{n!n}.$$

But working backwards (which we are allowed to do, as all quantities are strictly positive),

$$\begin{aligned}
 \frac{1}{(n+1)!} \cdot \frac{n+2}{n+1} < \frac{1}{n!n} &\Leftrightarrow n!n(n+2) < (n+1)!(n+1) \\
 &\Leftrightarrow n(n+2) < (n+1)(n+1) \\
 &\Leftrightarrow n^2 + 2n < n^2 + 2n + 1 \\
 &\Leftrightarrow 0 < 1,
 \end{aligned}$$

and the theorem is proved.  $\square$

**183 THEOREM**  $e$  is irrational.

**Proof:** Assume  $e$  is rational, with  $e = \frac{p}{q}$ , where  $p$  and  $q$  are positive integers and the fraction is in lowest terms. Since  $qe = p$ , an integer,  $q!e$  must also be an integer. Also  $q!y_q$  must be an integer, since

$$q!y_q = q! \left( 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \cdots + \frac{1}{q!} \right).$$

But by Lemma 182,

$$0 < e - y_q < \frac{1}{q!q} \implies 0 < q!(e - y_q) < \frac{1}{q} \leq 1.$$

That is, the integer  $q!(e - y_q)$  is strictly between 0 and 1, a contradiction.  $\square$

**184 THEOREM** The sequence  $\{n^{1/n}\}_{n=1}^{+\infty}$  is decreasing for  $n \geq 3$ . Also,  $n^{1/n} \rightarrow 1$  as  $n \rightarrow +\infty$ .

**Proof:** Consider the ratio

$$\frac{(n+1)^n}{n^{n+1}} = \left(1 + \frac{1}{n}\right)^n \cdot \frac{1}{n} < \frac{e}{n}.$$

Thus for  $n \geq 3$ ,

$$\frac{(n+1)^n}{n^{n+1}} < 1 \implies (n+1)^{1/(n+1)} < n^{1/n}.$$

Hence we have

$$3^{1/3} > 4^{1/4} > 5^{1/5} > \dots.$$


Clearly, if  $n > 1$  then  $n^{1/n} > 1^{1/n} = 1$ . Also, by the Binomial Theorem, again, if  $n > 1$ ,

$$\left(1 + \sqrt{\frac{2}{n}}\right)^n = 1^n + \binom{n}{1}\left(\sqrt{\frac{2}{n}}\right)^1 + \binom{n}{2}\left(\sqrt{\frac{2}{n}}\right)^2 + \dots > 1 + \binom{n}{2}\left(\sqrt{\frac{2}{n}}\right)^2 = 1 + \frac{n(n-1)}{2}\left(\frac{2}{n}\right) = n.$$

We then conclude that

$$1 < n^{1/n} < 1 + \sqrt{\frac{2}{n}},$$

and that  $n^{1/n} \rightarrow 1$  follows from the Sandwich Theorem.  $\square$

  $2^{1/2} = 4^{1/4}$ .

### Homework

**Problem 3.3.1** What's wrong with the following? Since the product of the limits is the limit of the product,

$$e = \lim_{n \rightarrow +\infty} \left(1 + \frac{1}{n}\right)^n = \underbrace{\left(\lim_{n \rightarrow +\infty} 1 + \frac{1}{n}\right) \cdot \left(\lim_{n \rightarrow +\infty} 1 + \frac{1}{n}\right) \cdots \left(\lim_{n \rightarrow +\infty} 1 + \frac{1}{n}\right)}_{n \text{ times}} = 1 \cdot 1 \cdots 1 = 1.$$

**Problem 3.3.2** Demonstrate that for all strictly positive integers  $n$ :

$$\cos \frac{\pi}{2^{n+1}} = \frac{1}{2} \sqrt{\underbrace{2 + \sqrt{2 + \sqrt{2 + \cdots + \sqrt{2}}}}_{n \text{ radicands}}},$$

$$\sin \frac{\pi}{2^{n+1}} = \frac{1}{2} \sqrt{\underbrace{2 - \sqrt{2 + \sqrt{2 + \cdots + \sqrt{2}}}}_{n \text{ radicands}}}.$$

Hence deduce Viète's Formula for  $\pi$ :

$$\pi = \lim_{n \rightarrow +\infty} 2^n \sqrt{\underbrace{2 - \sqrt{2 + \sqrt{2 + \cdots + \sqrt{2}}}}_{n \text{ radicands}}}.$$

**Problem 3.3.3** Prove that the sequence  $\left\{\sum_{k=n}^{2n} \frac{1}{k}\right\}_{n=1}^{+\infty}$  converges to  $\log 2$ .

**Problem 3.3.4** Prove that the sequence  $\left\{1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \cdots + \frac{1}{2n-1} - \frac{1}{2n}\right\}_{n=1}^{+\infty}$  converges to  $\log 2$ .

**Problem 3.3.5** Let  $n$  be a strictly positive integer and let  $x_n$  denote the unique real solution of the equation  $x^n + x + 1 = 0$ . Prove that  $x_n \rightarrow 1$  as  $n \rightarrow +\infty$ .

**Problem 3.3.6** Let

$$a_n = \sqrt{\underbrace{n + \sqrt{(n-1) + \sqrt{(n-2) + \cdots + \sqrt{2 + \sqrt{1}}}}}_{n \text{ radicands}}},$$

for  $n \geq 1$ . Prove that  $a_n - \sqrt{n} \rightarrow \frac{1}{2}$ .

**Problem 3.3.7** Prove that  $e$  is not a quadratic irrational.

**Problem 3.3.8** Find  $\lim_{n \rightarrow +\infty} \prod_{k=1}^n \left(1 + \frac{k}{n}\right)$ .

**Problem 3.3.9** A quadratic integer is any number  $x$  that satisfies an equation

$$x^2 + mx + n = 0, \quad (m, n) \in \mathbb{Z}^2.$$

Prove that the real quadratic integers are dense in the reals.

### 3.4 Averages of Sequences

**Why bother?** In this section we will examine some classical results that allow us to compute more complicated limits. Had we the language of matrices, most results here could be deduced from a classical result of Toeplitz. Since we don't, we will develop ad hoc methods which are interesting by themselves.

We start with the following discrete analogues of L'Hôpital's Rule.

**185 THEOREM** Let  $\{x_n\}_{n=1}^{+\infty}$ ,  $\{y_n\}_{n=1}^{+\infty}$  be two sequences of real numbers such that  $x_n \rightarrow 0$ ,  $y_n \rightarrow 0$ . Suppose, moreover, that the  $x_n$  are eventually strictly decreasing. Then

$$\lim_{n \rightarrow +\infty} \frac{x_n - x_{n-1}}{y_n - y_{n-1}} = \lim_{n \rightarrow +\infty} \frac{x_n}{y_n},$$

provided the sinistral limit exists (be it finite or  $+\infty$ ).

**Proof:** Assume first that  $\frac{x_{n-1} - x_n}{y_{n-1} - y_n} = \frac{x_n - x_{n-1}}{y_n - y_{n-1}} \rightarrow L$ , a finite real number. Then, given  $\varepsilon > 0$  we can find  $N > 0$  such that for  $n > N$ ,

$$L - \varepsilon < \frac{x_{n-1} - x_n}{y_{n-1} - y_n} < L + \varepsilon, \quad y_n < y_{n-1}.$$

Thus  $(L - \varepsilon)(y_{n-1} - y_n) < x_{n-1} - x_n < (L + \varepsilon)(y_{n-1} - y_n)$ , and repeating this inequality for  $n+1, n+2, \dots, n+m$ ,

$$\begin{aligned} (L - \varepsilon)(y_n - y_{n+1}) &< x_n - x_{n+1} < (L + \varepsilon)(y_n - y_{n+1}), \\ (L - \varepsilon)(y_{n+1} - y_{n+2}) &< x_{n+1} - x_{n+2} < (L + \varepsilon)(y_{n+1} - y_{n+2}), \\ &\vdots \\ (L - \varepsilon)(y_{m+n-1} - y_{m+n}) &< x_{m+n-1} - x_{m+n} < (L + \varepsilon)(y_{m+n-1} - y_{m+n}). \end{aligned}$$

Adding columnwise,

$$(L - \varepsilon)(y_n - y_{m+n}) < x_n - x_{m+n} < (L + \varepsilon)(y_n - y_{m+n}).$$

Letting  $m \rightarrow +\infty$ , and since the  $y_n$  are strictly positive,

$$(L - \varepsilon)y_n < x_n < (L + \varepsilon)y_n \implies L - \varepsilon < \frac{x_n}{y_n} < L + \varepsilon \implies \frac{x_n}{y_n} \rightarrow L$$

as  $n \rightarrow +\infty$ .

If  $\frac{x_{n-1} - x_n}{y_{n-1} - y_n}$  diverges to  $+\infty$  then for all  $M > 0$  we can find  $N' > 0$  such that for all  $n \geq N'$ ,

$$\frac{x_{n-1} - x_n}{y_{n-1} - y_n} > M \implies x_{n-1} - x_n > M(y_{n-1} - y_n).$$

Reasoning as above, for positive integers  $m \geq 0$ ,

$$x_n - x_{m+n} > M(y_n - y_{m+n}).$$

Taking the limit as  $m \rightarrow +\infty$ ,

$$x_n \geq M y_n \implies \frac{x_n}{y_n} \geq M \implies \frac{x_n}{y_n} \rightarrow +\infty.$$

□

**186 THEOREM (Stolz's Theorem)** Let  $\{a_n\}_{n=1}^{+\infty}$ ,  $\{b_n\}_{n=1}^{+\infty}$  be two sequences of real numbers. Suppose that  $\{b_n\}_{n=1}^{+\infty}$  is strictly increasing for sufficiently large  $n$  and that  $b_n \rightarrow +\infty$  as  $n \rightarrow +\infty$ . Then

$$\lim_{n \rightarrow +\infty} \frac{a_n - a_{n-1}}{b_n - b_{n-1}} = \lim_{n \rightarrow +\infty} \frac{a_n}{b_n},$$

provided the sinistral side exists (be it finite or infinite).

**Proof:** Assume first that  $\frac{a_n - a_{n-1}}{b_n - b_{n-1}} \rightarrow L$ , finite. Then for every  $\varepsilon > 0$  there is  $N > 0$  such that  $(\forall)n \geq N$ ,

$$L - \varepsilon < \frac{a_{n+1} - a_n}{b_{n+1} - b_n} < L + \varepsilon, \quad b_{n+1} > b_n.$$

This means that

$$(L - \varepsilon)(b_{n+1} - b_n) < a_{n+1} - a_n < (L + \varepsilon)(b_{n+1} - b_n)$$

By iterating the above relation for  $N+1, N+2, \dots, m+N$  we obtain

$$\begin{aligned} (L - \varepsilon)(b_{N+1} - b_N) &< a_{N+1} - a_N < (L + \varepsilon)(b_{N+1} - b_N), \\ (L - \varepsilon)(b_{N+2} - b_{N+1}) &< a_{N+2} - a_{N+1} < (L + \varepsilon)(b_{N+2} - b_{N+1}), \\ &\vdots \\ (L - \varepsilon)(b_{m+N} - b_{m+N-1}) &< a_{m+N} - a_{m+N-1} < (L + \varepsilon)(b_{m+N} - b_{m+N-1}). \end{aligned}$$

Adding columnwise,

$$(L - \varepsilon)(b_{m+N} - b_N) < a_{m+N} - a_N < (L + \varepsilon)(b_{m+N} - b_N) \implies \left| \frac{a_{m+N} - a_N}{b_{m+N} - b_N} - L \right| < \varepsilon.$$

Now,

$$\frac{a_{m+N}}{b_{m+N}} - L = \frac{a_N - Lb_N}{b_{m+N}} + \left(1 - \frac{b_N}{b_{m+N}}\right) \left(\frac{a_{m+N} - a_N}{b_{m+N} - b_N} - L\right),$$

so by the Triangle Inequality

$$\left| \frac{a_{m+N}}{b_{m+N}} - L \right| \leq \left| \frac{a_N - Lb_N}{b_{m+N}} \right| + \left| 1 - \frac{b_N}{b_{m+N}} \right| \left| \frac{a_{m+N} - a_N}{b_{m+N} - b_N} - L \right|.$$

Since  $N$  is fixed,  $\frac{a_N - Lb_N}{b_{m+N}} \rightarrow 0$  and  $\frac{b_N}{b_{m+N}} \rightarrow 0$  as  $m \rightarrow +\infty$ . Thus the dextral side is arbitrarily small, proving that  $\frac{a_m}{b_m} \rightarrow L$  as  $m \rightarrow +\infty$ .

Assume now that  $\frac{a_n - a_{n-1}}{b_n - b_{n-1}} \rightarrow +\infty$ . For sufficiently large  $n$  thus  $a_n - a_{n-1} > b_n - b_{n-1}$ . Thus the  $a_n$  are eventually increasing and  $a_n \rightarrow +\infty$  as  $n \rightarrow +\infty$ . Applying the results above to the  $\frac{b_n}{a_n}$  we obtain

$$\lim_{n \rightarrow +\infty} \frac{b_n}{a_n} = \lim_{n \rightarrow +\infty} \frac{b_n - b_{n-1}}{a_n - a_{n-1}} = 0$$

and so  $\lim_{n \rightarrow +\infty} \frac{a_n}{b_n} = +\infty$  too.  $\square$

**187 THEOREM (Cèsaro)** If a sequence of real numbers converges to a number, then its sequence of arithmetic means converges to the same number, that is, if  $x_n \rightarrow a$  then  $\frac{x_1 + x_2 + \dots + x_n}{n} \rightarrow a$ .

**First Proof:** Let  $a_n = x_1 + x_2 + \dots + x_n$  and  $b_n = n$  in Stolz's Theorem.  $\square$

**Second Proof:** It is instructive to give an ad hoc proof of this result. Given  $\varepsilon > 0$  there exists  $N > 0$  such that if  $n \geq N$  then  $|x_n - a|$ . Then

$$\left| \frac{x_1 + x_2 + \dots + x_n}{n} - a \right| = \left| \frac{(x_1 - a) + (x_2 - a) + \dots + (x_n - a)}{n} \right| \leq \frac{|(x_1 - a)| + |(x_2 - a)| + \dots + |(x_n - a)|}{n}.$$

Now we run into a slight problem. We can control the differences  $|x_k - a|$  after a certain point, but the early differences need to be taken care of. To this end we consider the differences  $x_k - a$  with  $k \leq \lfloor \sqrt{n} \rfloor$  or  $k > \lfloor n \rfloor$  where  $n$  is so large that  $\lfloor \sqrt{n} \rfloor \geq N$ . We have

$$\begin{aligned} \frac{|(x_1 - a)| + |(x_2 - a)| + \cdots + |(x_n - a)|}{n} &= \frac{|(x_1 - a)| + |(x_2 - a)| + \cdots + |(x_{\lfloor \sqrt{n} \rfloor} - a)|}{n} \\ &\quad + \frac{|(x_{\lfloor \sqrt{n} \rfloor + 1} - a)| + |(x_2 - a)| + \cdots + |(x_n - a)|}{n} \\ &< \frac{\lfloor \sqrt{n} \rfloor \max_{1 \leq k \leq \lfloor \sqrt{n} \rfloor} |x_k - a|}{n} + \frac{(n - \lfloor \sqrt{n} \rfloor)\varepsilon}{n}. \end{aligned}$$

These two last quantities tend to 0 as  $n \rightarrow +\infty$ , from where the result follows.  $\square$

**188 Example** Since  $n^{1/n} \rightarrow 1$ ,  $\frac{1 + 2^{1/2} + 3^{1/3} + \cdots + n^{1/n}}{n} \rightarrow 1$ .

**189 Example** Since  $\frac{1}{n} \rightarrow 0$ ,  $\frac{1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n}}{n} \rightarrow 0$ .

**190 Example** Since  $\left(1 + \frac{1}{n}\right)^n \rightarrow e$ ,  $\frac{\left(1 + \frac{1}{1}\right)^1 + \left(1 + \frac{1}{2}\right)^2 + \left(1 + \frac{1}{3}\right)^3 + \cdots + \left(1 + \frac{1}{n}\right)^n}{n} \rightarrow e$ .

**191 Example** The converse of Cèsaro's Theorem is false. For, the sequence  $a_n = (-1)^n$  oscillates and does not converge. But its sequence of averages is  $b_n = \frac{1 - 1 + 1 - 1 + \cdots + (-1)^n}{n} \rightarrow 0$  as  $n \rightarrow +\infty$  since the numerator is either 0 or -1.

**192 THEOREM** If a sequence of positive real numbers converges to a number, then its sequence of geometric means converges to the same number, that is, if  $\forall n > 0$ ,  $x_n \geq 0$  and  $x_n \rightarrow a$  then  $(x_1 x_2 \cdots x_n)^{1/n} \rightarrow a$ .

**Proof:** The proof mimics Cèsaro's Theorem 187. Since  $x_n \rightarrow a$ , for all  $\varepsilon > 0$  there is  $N > 0$  such that for all  $n \geq N$ ,

$$|x_n - a| < \varepsilon \implies a - \varepsilon < x_n < a + \varepsilon.$$

Then

$$\left( \min_{1 \leq k \leq \lfloor \sqrt{n} \rfloor} x_k \right)^{\lfloor \sqrt{n} \rfloor / n} (x_{\lfloor \sqrt{n} \rfloor + 1} \cdots x_n)^{1/n} \leq (x_1 x_2 \cdots x_{\lfloor \sqrt{n} \rfloor} x_{\lfloor \sqrt{n} \rfloor + 1} \cdots x_n)^{1/n} \leq \left( \max_{1 \leq k \leq \lfloor \sqrt{n} \rfloor} x_k \right)^{\lfloor \sqrt{n} \rfloor / n} (x_{\lfloor \sqrt{n} \rfloor + 1} \cdots x_n)^{1/n}.$$

This gives, for  $\lfloor \sqrt{n} \rfloor \geq N$ ,

$$\left( \min_{1 \leq k \leq \lfloor \sqrt{n} \rfloor} x_k \right)^{\lfloor \sqrt{n} \rfloor / n} (a - \varepsilon)^{(n - \lfloor \sqrt{n} \rfloor) / n} \leq (x_1 x_2 \cdots x_{\lfloor \sqrt{n} \rfloor} x_{\lfloor \sqrt{n} \rfloor + 1} \cdots x_n)^{1/n} \leq \left( \max_{1 \leq k \leq \lfloor \sqrt{n} \rfloor} x_k \right)^{\lfloor \sqrt{n} \rfloor / n} (a + \varepsilon)^{(n - \lfloor \sqrt{n} \rfloor) / n}.$$

Now, both  $\left( \min_{1 \leq k \leq \lfloor \sqrt{n} \rfloor} x_k \right)^{\lfloor \sqrt{n} \rfloor / n}$  and  $\left( \max_{1 \leq k \leq \lfloor \sqrt{n} \rfloor} x_k \right)^{\lfloor \sqrt{n} \rfloor / n}$  converge to 1 as  $n \rightarrow +\infty$  by virtue of Theorem 174, and again by the same theorem,

$$(a - \varepsilon)^{(n - \lfloor \sqrt{n} \rfloor) / n} = (a - \varepsilon)(a - \varepsilon)^{\lfloor \sqrt{n} \rfloor / n} \rightarrow a - \varepsilon, \quad (a + \varepsilon)^{(n - \lfloor \sqrt{n} \rfloor) / n} = (a + \varepsilon)(a + \varepsilon)^{\lfloor \sqrt{n} \rfloor / n} \rightarrow a + \varepsilon$$

as  $n \rightarrow +\infty$ . This gives the result.  $\square$

**193 Example** Since  $e_n = \left(\frac{n+1}{n}\right)^n \rightarrow e$ , then by the Theorem 192

$$(e_1 e_2 \cdots e_n)^{1/n} = \left(\left(\frac{2}{1}\right)^1 \left(\frac{3}{2}\right)^2 \left(\frac{4}{3}\right)^3 \cdots \left(\frac{n+1}{n}\right)^n\right)^{1/n} = \left(\frac{(n+1)^n}{n!}\right)^{1/n} \rightarrow e.$$

This gives  $\frac{n}{(n!)^{1/n}} = \frac{n}{n+1} \cdot \frac{n+1}{(n!)^{1/n}} \rightarrow 1 \cdot e = e$ .

### Homework

**Problem 3.4.1** If  $\{a_n\}_{n=1}^{+\infty}$  is a sequence of strictly positive real numbers such that  $\frac{a_n}{a_{n-1}} \rightarrow a > 0$ . Prove that

$$\lim_{n \rightarrow +\infty} \frac{a_n}{a_{n-1}} = \lim_{n \rightarrow +\infty} \sqrt[n]{a_n}.$$

**Problem 3.4.2** Let  $x_n \rightarrow a$  and  $y_n \rightarrow b$ . Prove that  $\frac{x_1 y_n + x_2 y_{n-1} + \cdots + x_n y_1}{n} \rightarrow ab$ .

**Problem 3.4.3** Prove that  $\lim_{n \rightarrow +\infty} \left(\frac{(2n)}{n}\right)^{1/n} = 4$ .

**Problem 3.4.4** Prove that  $\lim_{n \rightarrow +\infty} \frac{1}{n} (n(n+1) \cdots (n+n))^{1/n} = 4e$ .


**Problem 3.4.5** Prove that  $\lim_{n \rightarrow +\infty} \frac{1}{n} (1 \cdot 3 \cdot 5 \cdots (2n-1))^{1/n} = \frac{2}{e}$ .

**Problem 3.4.6** Prove that  $\lim_{n \rightarrow +\infty} \frac{1}{n^2} \left(\frac{(3n)!}{n!}\right)^{1/n} = \frac{2}{e}$ .

## 3.5 Orders of Infinity

**Why bother?** It is clear that the sequences  $\{n\}_{n=1}^{+\infty}$  and  $\{n^2\}_{n=1}^{+\infty}$  both tend to  $+\infty$  as  $n \rightarrow +\infty$ . We would like now to refine this statement and compare one with the other. In other words, we will examine their speed towards  $+\infty$ .


**194 Definition** We write  $a_n = O(b_n)$  if  $\exists C > 0, \exists N > 0$  such that  $\forall n \geq N$  we have  $|a_n| \leq C|b_n|$ . We then say that  $a_n$  is *Big Oh* of  $b_n$ , or that  $a_n$  is of order at most  $b_n$  as  $n \rightarrow +\infty$ . Observe that this means that  $\left|\frac{a_n}{b_n}\right|$  is bounded for sufficiently large  $n$ . The notation  $a_n \ll b_n$ , due to Vinogradov, is often used as a synonym of  $a_n = O(b_n)$ .

 A sequence  $\{a_n\}_{n=1}^{+\infty}$  is bounded if and only if  $a_n \ll 1$ .

An easy criterion to identify Big Oh relations is the following.

**195 THEOREM** If  $\lim_{n \rightarrow +\infty} \frac{a_n}{b_n} = c \in \mathbb{R}$ , then  $a_n \ll b_n$ .

**Proof:** By Theorem 164, a convergent sequence is bounded, hence the sequence  $\left\{\frac{a_n}{b_n}\right\}_{n=1}^{+\infty}$  is bounded: so for sufficiently large  $n$ ,  $\left|\frac{a_n}{b_n}\right| < C$  for some constant  $C > 0$ . This proves the theorem.  $\square$

 The  $=$  in the relation  $a_n = O(b_n)$  is not a true equal sign. For example  $n^2 = O(n^3)$  since  $\lim_{n \rightarrow +\infty} \frac{n^2}{n^3} = 0$  and so  $n^2 \ll n^3$  by Theorem 195. On the other hand,  $\lim_{n \rightarrow +\infty} \frac{n^3}{n^2} = +\infty$  so that for sufficiently large  $n$ , and for all  $M > 0$ ,  $n^3 > Mn^2$ , meaning that  $n^3 \neq O(n^2)$ . Thus the Big Oh relation is not symmetric.<sup>3</sup>

<sup>3</sup>One should more properly write  $a_n \in O(b_n)$ , as  $O(b_n)$  is the set of sequences growing to infinity no faster than  $b_n$ , but one keeps the  $=$  sign for historical reasons.

**196 THEOREM (Lexicographic Order of Powers)** Let  $(\alpha, \beta) \in \mathbb{R}$  and consider the sequences  $\{n^\alpha\}_{n=1}^{+\infty}$  and  $\{n^\beta\}_{n=1}^{+\infty}$ . Then  $n^\alpha \ll n^\beta \iff \alpha \leq \beta$ .

**Proof:** If  $\alpha \leq \beta$  then  $\lim_{n \rightarrow +\infty} \frac{n^\alpha}{n^\beta}$  is either 1 (when  $\alpha = \beta$ ) or 0 (when  $\alpha < \beta$ ), hence  $n^\alpha \ll n^\beta$  by Theorem 195.

If  $n^\alpha \ll n^\beta$  then for sufficiently large  $n$ ,  $n^\alpha \leq Cn^\beta$  for some constant  $C > 0$ . If  $\alpha > \beta$  then this would mean that for all large  $n$  we would have  $n^{\alpha-\beta} \leq C$ , which is absurd, since for a strictly positive exponent  $\alpha - \beta$ ,  $n^{\alpha-\beta} \rightarrow +\infty$  as  $n \rightarrow +\infty$ .  $\square$

**197 Example** As  $n \rightarrow +\infty$ ,

$$n^{1/10} \ll n^{1/3} \ll n \ll n^{10/9} \ll n^2,$$

for example.

**198 THEOREM** If  $c \in \mathbb{R} \setminus \{0\}$  then  $O(ca_n) = O(a_n)$ , that is, the set of sequences of order at most  $ca_n$  is the same set at those of order at most  $a_n$ .

**Proof:** We prove that  $b_n = O(ca_n) \iff b_n = O(a_n)$ . If  $b_n = O(ca_n)$  then there are constants  $C > 0$  and  $N > 0$  such that  $|b_n| \leq C|ca_n|$  whenever  $n \geq N$ . Therefore,  $|b_n| \leq C'|a_n|$  whenever  $n \geq N$ , where  $C' = C|c|$ , meaning that  $b_n = O(a_n)$ . Similarly, if  $b_n = O(a_n)$  then there are constants  $C_1 > 0$  and  $N_1 > 0$  such that  $|b_n| \leq C_1|a_n|$  whenever  $n \geq N_1$ . Since  $c \neq 0$  this is equivalent to  $|b_n| \leq \frac{C_1}{c}(c|a_n|) = C''(c|a_n|)$  whenever  $n \geq N_1$ , where  $C'' = \frac{C_1}{c}$ , meaning that  $b_n = O(ca_n)$ . Therefore,  $O(a_n) = O(ca_n)$ .  $\square$

**199 Example** As  $n \rightarrow +\infty$ ,

$$O(n^3) = O\left(\frac{n^3}{3}\right) = O(4n^3).$$

**200 THEOREM (Sum Rule)** Let  $a_n = O(x_n)$  and  $b_n = O(y_n)$ . Then  $a_n + b_n = O(\max(|x_n|, |y_n|))$ .

**Proof:** There exist strictly positive constants  $C_1, N_1, C_2, N_2$  such that

$$n \geq N_1, \implies |a_n| \leq C_1|x_n| \quad \text{and} \quad n \geq N_2, \implies |b_n| \leq C_2|y_n|.$$

Let  $N' = \max(N_1, N_2)$ . Then for  $n \geq N'$ , by the Triangle inequality

$$|a_n + b_n| \leq |a_n| + |b_n| \leq C_1|x_n| + C_2|y_n|.$$

Let  $C' = \max(C_1, C_2)$ . Then

$$|a_n + b_n| \leq C'(|x_n| + |y_n|) \leq 2C' \max(|x_n|, |y_n|),$$

whence the theorem follows.  $\square$

**201 COROLLARY** Let  $a_n = k_0n^m + k_1n^{m-1} + k_2n^{m-2} + \dots + k_{m-1}n + k_m$  be a polynomial of degree  $m$  in  $n$  with real number coefficients. The  $a_n = O(n^m)$ , that is,  $a_n$  is of order at most its leading term.

**Proof:** By the Sum Rule Theorem 200 the leading term dominates.  $\square$

**202 THEOREM (Transitivity Rule)** If  $a_n = O(b_n)$  and  $b_n = O(c_n)$ , then  $a_n = O(c_n)$ .

**Proof:** There are strictly positive constants  $C_1, C_2, N_1, N_2$  such that

$$n \geq N_1, \implies |a_n| \leq C_1|b_n| \quad \text{and} \quad n \geq N_2, \implies |b_n| \leq C_2|c_n|.$$

If  $n \geq \max(N_1, N_2)$ , then  $|a_n| \leq C_1|b_n| \leq C_1C_2|c_n| = C|c_n|$ , with  $C = C_1C_2$ . This gives  $a_n = O(c_n)$ .  $\square$

**203 Example** By Corollary 201,  $5n^4 - 2n^2 + 100n - 8 = O(5n^4)$ . By Theorem 198,  $O(5n^4) = O(n^4)$ . Hence

$$5n^4 - 2n^2 + 100n - 8 = O(n^4).$$

**204 THEOREM (Multiplication Rule)** If  $a_n = O(x_n)$  and  $b_n = O(y_n)$ , then  $a_n b_n = O(x_n y_n)$ .

**Proof:** There are strictly positive constants  $C_1, C_2, N_1, N_2$  such that

$$n \geq N_1, \implies |a_n| \leq C_1 |x_n| \quad \text{and} \quad n \geq N_2, \implies |b_n| \leq C_2 |y_n|.$$

If  $n \geq \max(N_1, N_2)$ , then  $|a_n b_n| \leq C_1 C_2 |x_n y_n| = C |x_n y_n|$ , with  $C = C_1 C_2$ . This gives  $a_n b_n = O(x_n y_n)$ .  $\square$

**205 THEOREM (Lexicographic Order of Exponentials)** Let  $(a, b) \in \mathbb{R}$ ,  $a > 1$ ,  $b > 1$ , and consider the sequences  $\{a^n\}_{n=1}^{+\infty}$  and  $\{b^n\}_{n=1}^{+\infty}$ . Then  $a^n \ll b^n \iff a \leq b$ .

**Proof:** Put  $r = \frac{a}{b}$ , and use Theorems 172 and 195.  $\square$

**206 Example**  $\frac{1}{2^n} \ll 1 \ll 2^n \ll e^n \ll 3^n$ .

**207 LEMMA** Let  $a \in \mathbb{R}$ ,  $a > 1$ ,  $k \in \mathbb{N} \setminus \{0\}$ . Then  $n^k \ll a^n$ .

**Proof:** By Theorem 175,  $\lim_{n \rightarrow +\infty} \frac{n^k}{a^n} = 0$ . Now apply Theorem 195.  $\square$

**208 THEOREM ("Exponentials are faster than powers")** Let  $a \in \mathbb{R}$ ,  $a > 1$ ,  $\alpha \in \mathbb{R}$ . Then  $n^\alpha \ll a^n$ .

**Proof:** Put  $k = \max(1, \lfloor \alpha \rfloor + 1)$ . Then by Theorem 196,  $n^\alpha \ll n^k$ . By Lemma 207,  $n^k \ll a^n$ , and by the Transitivity of Big Oh (Theorem 202),  $n^\alpha \ll n^k \ll a^n$ .  $\square$

**209 Example**

$$n^{100} \ll e^n.$$

**210 THEOREM ("Logarithms are slower than powers")** Let  $(\alpha, \beta) \in \mathbb{R}^2$ ,  $\alpha > 0$ . Then  $(\log n)^\beta \ll n^\alpha$ .

**Proof:** If  $\beta \leq 0$ , then  $(\log n)^\beta \ll 1$  and the assertion is evident, so assume  $\beta > 0$ . For  $x > 0$ , then  $\log x < x$ . Putting  $x = n^{\alpha/\beta}$ , we get

$$\log n^{\alpha/\beta} < n^{\alpha/\beta} \implies \log n < \frac{\beta n^{\alpha/\beta}}{\alpha} \implies (\log n)^\beta < \frac{\beta^\beta n^\alpha}{\alpha^\beta},$$

whence  $(\log n)^\beta \ll n^\alpha$ .  $\square$

By the Multiplication Rule (Theorem 204) and Theorems 196, 208, 210, in order to compare two expressions of the type  $a^n n^b (\log)^c$  and  $u^n n^v (\log)^w$  we simply look at the lexicographic order of the exponents, keeping in mind that logarithms are slower than powers, which are slower than exponentials.

**211 Example** In increasing order of growth we have

$$\frac{1}{e^n} \ll \frac{1}{2^n} \ll \frac{1}{n^2} = \frac{1}{\log n} \ll 1 \ll (\log \log n)^{10} \ll \sqrt{\log n} \ll \frac{n}{\log n} \ll n \ll n \log n \ll e^n.$$

**212 Example** Decide which one grows faster as  $n \rightarrow +\infty$ :  $n^{\log n}$  or  $(\log n)^n$ .




**Solution:** Since  $n^{\log n} = e^{(\log n)^2}$  and  $(\log n)^n = e^{n \log \log n}$ , and since  $(\log n)^2 \ll n \log \log n$ , we conclude that  $n^{\log n} \ll (\log n)^n$ .

We now define two more fairly common symbols in asymptotic analysis.

**213 Definition** We write  $a_n = o(b_n)$  if  $\frac{a_n}{b_n} \rightarrow 0$  as  $n \rightarrow +\infty$ , and say that  $a_n$  is *small oh* of  $b_n$ , or that  $a_n$  grows slower than  $b_n$  as  $n \rightarrow +\infty$ .

**214 Definition** A sequence  $\{a_n\}_{n=1}^{+\infty}$  is said to be *infinitesimal* if  $a_n = o(1)$ , that is, if  $a_n \rightarrow 0$  as  $n \rightarrow +\infty$ .

 We know from above that for  $a > 1$   $\lim_{n \rightarrow +\infty} \frac{n^a}{a^n} = 0$ , and so  $n^a = o(a^n)$ . Also, for  $\gamma > 0$ ,  $\lim_{n \rightarrow +\infty} \frac{(\log n)^\beta}{n^\gamma} = 0$ , and so  $(\log n)^\beta = o(n^\gamma)$ .

**215 Definition** We write  $a_n \sim b_n$  if  $\frac{a_n}{b_n} \rightarrow 1$  as  $n \rightarrow +\infty$ , and say that  $a_n$  is *asymptotic* to  $b_n$ .

Asymptotic sequences are thus those that grow at the same rate as the index increases.




Figure 3.2: Diagram of  $O$  relations.

**216 Example** The sequences  $\{n^2 - n \sin n\}_{n=1}^{+\infty}$ ,  $\{n^2 + n - 1\}_{n=1}^{+\infty}$  are asymptotic since

$$\frac{n^2 - n \sin n}{n^2 + n - 1} = \frac{1 - \frac{\sin n}{n}}{1 + \frac{1}{n} - \frac{1}{n^2}} \rightarrow 1,$$

as  $n \rightarrow +\infty$ .

**217 THEOREM** Let  $\{a_n\}_{n=1}^{+\infty}$  and  $\{b_n\}_{n=1}^{+\infty}$  be two properly diverging sequences. Then  $a_n \sim b_n \iff a_n = b_n(1 + o(1))$ .

**Proof:** Since the limit is  $1 > 0$ , either both diverge to  $+\infty$  or both to  $-\infty$ . Assume the former, and so, eventually,  $b_n$  will be strictly positive. Now,

$$\begin{aligned} \lim_{n \rightarrow +\infty} \frac{a_n}{b_n} = 1 &\iff \forall \varepsilon > 0, \exists N > 0, 1 - \varepsilon < \frac{a_n}{b_n} < 1 + \varepsilon \\ &\iff b_n - b_n \varepsilon < a_n < b_n + b_n \varepsilon \\ &\iff |a_n - b_n| < b_n \varepsilon \\ &\iff a_n - b_n = o(b_n). \end{aligned}$$

□

The relationship between the three symbols is displayed in figure 3.2.

### Homework

**Problem 3.5.1** Prove that  $e^n \ll n!$ .

**Problem 3.5.2** Prove that  $O(O(a_n)) = O(a_n)$ .

**Problem 3.5.3** Let  $k \in \mathbb{R}$  be a constant. Prove that  $k + O(a_n) = O(k + a_n) = O(a_n)$ .

**Problem 3.5.4** Let  $k \in \mathbb{R}$ ,  $k > 0$ , be a constant. Prove that  $(a_n + b_k)^k \ll a_n^k + b_k^k$ .

**Problem 3.5.5** For a sequence of real numbers  $\{a_n\}_{n=1}^{+\infty}$  it is known

that  $a_n = O(n^2)$  and  $a_n = o(n^2)$ . Which of the two statements conveys more information?

**Problem 3.5.6** True or false:  $a_n = O(n) \implies a_n = o(n)$ .

**Problem 3.5.7** True or false:  $a_n = o(n) \implies a_n = O(n)$ .

**Problem 3.5.8** True or false:  $a_n = o(n^2) \implies a_n = O(n)$ .

**Problem 3.5.9** True or false:  $a_n = o(n) \implies a_n = O(n^2)$ .

## 3.6 Cauchy Sequences

**218 Definition** A sequence of real numbers  $\{a_n\}_{n=1}^{+\infty}$  is called a *Cauchy Sequence* if

$$\forall \varepsilon > 0, \exists N > 0, \text{ such that } \forall n, m \geq N \quad |a_n - a_m| < \varepsilon.$$

**219 THEOREM** Cauchy sequences are bounded.

**Proof:** Let  $\{a_n\}_{n=1}^{+\infty}$  be Cauchy. Take  $N > 0$  such that for all  $n \geq N$ ,  $|a_n - a_N| < 1$ . Then  $a_n$  is bounded by

$$\max(|a_1|, |a_2|, \dots, |a_N|) + 1.$$

□

**220 LEMMA** If a Cauchy sequence of real numbers has a convergent subsequence, then the parent sequence converges, and it does so to the same limit as the subsequence.

**Proof:** Let  $\{a_n\}_{n=1}^{+\infty}$  be a Cauchy sequence of real numbers, and suppose that its subsequence  $\{a_{n_k}\}_{k=1}^{+\infty}$  converges to the real number  $a$ . Given  $\varepsilon > 0$ , take  $N > 0$  sufficiently large such that

$$\forall m, n, n_k \geq N, \quad |a_n - a_m| < \varepsilon, \quad \text{and} \quad |a_{n_k} - a| < \varepsilon.$$

By the Triangle Inequality,

$$|a_n - a| \leq |a_n - a_{n_k}| + |a_{n_k} - a| < \varepsilon + \varepsilon = 2\varepsilon,$$

whence  $a_n \rightarrow a$ . □

**221 THEOREM (General Principle of Convergence)** A sequence of real numbers converges if and only if it is Cauchy.

**Proof:**

( $\implies$ ) If  $a_n \rightarrow a$ , given  $\varepsilon > 0$ , choose  $N > 0$  such that  $|a_n - a| < \varepsilon$  for all  $n \geq N$ .

Then if  $m, n \geq N$ ,

$$|a_n - a_m| \leq |a_n - a| + |a_m - a| \leq \varepsilon + \varepsilon = 2\varepsilon.$$

Since  $2\varepsilon > 0$  can be made arbitrarily small,  $a_n$  is Cauchy.

( $\Leftarrow$ ) Suppose  $\mathbf{a}_n$  is Cauchy. By virtue of Theorem 219 it is bounded, say that for all  $n > 0$ ,  $\mathbf{a}_n \in [\alpha; \beta]$ . Put

$$\mathcal{S} = \{s : \mathbf{a}_n \geq s \text{ for infinitely many } n\}.$$

As  $\alpha \in \mathcal{S}$ ,  $\mathcal{S} \neq \emptyset$ .  $\mathcal{S}$  is bounded above by  $\beta$ . By the Completeness Axiom,  $\mathcal{S}$  has a supremum,  $\mathbf{a} = \sup \mathcal{S}$ . Given  $\varepsilon > 0$ ,  $\mathbf{a} - \varepsilon < \mathbf{a}$  and so there is  $s \in \mathcal{S}$  such that  $\mathbf{a} - \varepsilon < s$ . By definition of  $\mathcal{S}$ , there are infinitely many  $n$  with  $\mathbf{a}_n \geq s > \mathbf{a} - \varepsilon$ .  $\mathbf{a} + \varepsilon > \mathbf{a}$ , so that  $\mathbf{a} + \varepsilon \notin \mathcal{S}$  and so there are only finitely many  $n$  for which  $\mathbf{a}_n \geq \mathbf{a} + \varepsilon$ . Thus there are infinitely many  $n$  with  $\mathbf{a}_n \in (\mathbf{a} - \varepsilon, \mathbf{a} + \varepsilon)$ .

Choose  $N > 0$  such that  $|\mathbf{a}_n - \mathbf{a}_m| < \varepsilon$  for all  $m, n \geq N$ . We can find  $m \geq N$  with  $\mathbf{a}_m \in (\mathbf{a} - \varepsilon, \mathbf{a} + \varepsilon)$  ie  $|\mathbf{a}_m - \mathbf{a}| < \varepsilon$ . Then if  $n \geq N$ ,

$$|\mathbf{a}_n - \mathbf{a}| \leq |\mathbf{a}_n - \mathbf{a}_m| + |\mathbf{a}_m - \mathbf{a}| < \varepsilon + \varepsilon = 2\varepsilon$$

As  $2\varepsilon$  can be made arbitrarily small this shews  $\mathbf{a}_n \rightarrow \mathbf{a}$ .

□

## Homework

### 3.7 Topology of sequences. Limit Superior and Limit Inferior

**222 THEOREM** A set  $X \subseteq \mathbb{R}$  is dense in  $\mathbb{R}$  if and only if for every  $x \in \mathbb{R}$  there is a sequence  $\{x_n\}_{n=1}^{+\infty}$  of elements of  $X \setminus \{x\}$  that converges to  $x$ .

**Proof:**

$\Rightarrow$  For each positive integer  $n$ , since  $X$  is dense in  $\mathbb{R}$ , there exists  $x_n \in X \setminus \{x\}$  such that  $|x_n - x| < \frac{1}{2^n}$ . But then  $x_n \rightarrow x$  as  $n \rightarrow +\infty$ .

$\Leftarrow$  Let  $x \in \mathbb{R}$  and let  $\{x_n\}_{n=1}^{+\infty}$  of elements of  $X \setminus \{x\}$  that converges to  $x$ . Then  $\forall \varepsilon > 0$ ,  $\exists N \in \mathbb{N}$  such that  $\forall n \geq N$ ,  $|x_n - x| < \varepsilon$ . But then we have found elements of  $X \setminus \{x\}$  which are arbitrarily close to  $x$ , meaning that  $X$  is dense in  $\mathbb{R}$ .

□

**223 THEOREM** Let  $X \subseteq \mathbb{R}$ . A point  $x \in \mathbb{R}$  is an accumulation point of  $X$  if and only if there exists a sequence of elements of  $X \setminus \{x\}$  converging to  $x$ .

**Proof:**

$\Rightarrow$  If  $x$  is an accumulation point of  $X$ , every closed interval  $I_n := [x - 1/n; x + 1/n]$ ,  $n \in \mathbb{N}$ , satisfies  $I_n \cap (X \setminus \{x\}) \neq \emptyset$ , thus  $\forall n \in \mathbb{N}$ ,  $\exists x_n \in I_n \cap (X \setminus \{x\})$ . Since  $|x_n - x| < \frac{1}{n}$ , we conclude that  $\lim x_n = x$ .

$\Leftarrow$  Suppose now that  $\{x_n\}_{n=1}^{+\infty}$  is an infinite sequence of points of  $X \setminus \{x\}$  converging to  $x$ . If  $x \notin \text{Acc}(X)$ , then  $x \notin \text{Acc}(x_1, x_2, \dots)$ . Thus there is a neighbourhood of  $x$ ,  $\mathcal{N}_x$  such that  $\mathcal{N}_x \cap \{x_1, x_2, \dots\} = \emptyset$ . Thus there is a  $\varepsilon > 0$  such that  $|x - \varepsilon; x + \varepsilon| \subseteq \mathcal{N}_x$ . For this  $\varepsilon$  and for none of the  $x_n$  it is true then that  $|x_n - x| < \varepsilon$ , contradicting the fact that  $\lim_{n \rightarrow +\infty} x_n = x$ .

□

**224 Definition** Given a sequence  $\{a_n\}_{n=1}^{+\infty}$ , the new sequence

$$b_k = \inf_{n \geq k} a_n = \inf\{a_k, a_{k+1}, a_{k+2}, \dots\}, \quad k \geq 1,$$

satisfies  $b_k \leq b_{k+1}$ , that is, it is increasing, and hence it converges to its supremum. We then put

$$\lim_{n \rightarrow +\infty} \inf_{k \geq n} a_k = \sup_{n \geq 1} \inf_{k \geq n} a_k.$$

Similarly, the new sequence

$$c_k = \sup_{n \geq k} a_n = \sup\{a_k, a_{k+1}, a_{k+2}, \dots\}, \quad k \geq 1,$$

satisfies  $c_k \geq c_{k+1}$ , that is, it is decreasing, and hence it converges to its infimum. We then put

$$\limsup_{n \rightarrow +\infty} a_n = \inf_{n \geq 1} \sup_{k \geq n} a_k.$$

We now prove the following theorem for future reference.

**225 THEOREM** For any sequence  $\{a_n\}_{n=0}^{+\infty}$  of strictly positive real numbers

$$\liminf_{n \rightarrow +\infty} \frac{a_{n+1}}{a_n} \leq \liminf_{n \rightarrow +\infty} \sqrt[n]{a_n} \leq \limsup_{n \rightarrow +\infty} \sqrt[n]{a_n} \leq \limsup_{n \rightarrow +\infty} \frac{a_{n+1}}{a_n}.$$

**Proof:** We will prove the last inequality. The first is quite similar, and the two middle ones are obvious.

Put  $r = \limsup_{n \rightarrow +\infty} \frac{a_{n+1}}{a_n}$ . If  $r = +\infty$  then there is nothing to prove. For  $r < +\infty$  choose  $r' > r$ . There is  $N \in \mathbb{N}$  such that

$$\forall n \geq N, \quad \frac{a_{n+1}}{a_n} \leq r'.$$

Hence,

$$a_{N+1} \leq r' a_N, \quad a_{N+2} \leq r' a_{N+1}, \quad a_{N+3} \leq r' a_{N+2}, \dots, \quad a_{N+t} \leq r' a_{N+t-1},$$

and so, upon multiplication and cancelling,

$$a_{N+t} \leq a_N (r')^t,$$

and putting  $n = N + t$ ,

$$a_n \leq a_N (r')^{-N} (r')^n \implies \sqrt[n]{a_n} \leq r' \sqrt[n]{a_N (r')^{-N}} \implies \limsup_{n \rightarrow +\infty} \sqrt[n]{a_n} \leq r',$$

since  $a_N (r')^{-N}$  is a fixed real number (does not depend on  $n$ ), and so,  $\sqrt[n]{a_N (r')^{-N}} \rightarrow 1$  by Theorem 174.

□

The following theorem is an easy exercise left to the reader.

**226 THEOREM** Let  $\{a_n\}_1^{+\infty}$  be a sequence of real numbers. Then

1. if  $\limsup_{n \rightarrow +\infty} a_n = +\infty$ , then  $\{a_n\}_1^{+\infty}$  has a subsequence converging to  $+\infty$ .

2. if  $\limsup_{n \rightarrow +\infty} a_n = -\infty$ , then  $\lim_{n \rightarrow +\infty} a_n = -\infty$ .

3. if  $\limsup_{n \rightarrow +\infty} a_n = a \in \mathbb{R}$ , then

$$\forall \epsilon > 0, \exists n_0 \text{ such that } a_n < a + \epsilon \text{ whenever } n \geq n_0$$

and also, there are infinitely many  $a_n$  such that  $a - \epsilon < a_n$ .

4. if  $\liminf_{n \rightarrow +\infty} a_n = -\infty$ , then  $\{a_n\}_1^{+\infty}$  has a subsequence converging to  $-\infty$ .

5. if  $\liminf_{n \rightarrow +\infty} a_n = +\infty$ , then  $\lim_{n \rightarrow +\infty} a_n = +\infty$ .

6. if  $\liminf_{n \rightarrow +\infty} a_n = a \in \mathbb{R}$ , then

$$\forall \epsilon > 0, \exists n_0 \text{ such that } a - \epsilon < a_n \text{ whenever } n \geq n_0$$

and there are infinitely many  $a_n$  such that  $a_n < a + \epsilon$ .

7.  $\liminf_{n \rightarrow +\infty} a_n \leq \limsup_{n \rightarrow +\infty} a_n$  is always verified, and furthermore,  $\liminf_{n \rightarrow +\infty} a_n = \limsup_{n \rightarrow +\infty} a_n$  if and only if  $\lim_{n \rightarrow +\infty} a_n$  exists, in which case  $\liminf_{n \rightarrow +\infty} a_n = \lim_{n \rightarrow +\infty} a_n = \limsup_{n \rightarrow +\infty} a_n$ .

## Homework

**Problem 3.7.1** Identify the set of accumulation points of the set  $\{\sqrt{a} - \sqrt{b} : (a, b) \in \mathbb{N}^2\}$ .

**Problem 3.7.2** Consider the following enumeration of the proper fractions

$$\frac{0}{1}, \frac{1}{1}, \frac{0}{2}, \frac{1}{2}, \frac{2}{2}, \frac{0}{3}, \frac{1}{3}, \frac{2}{3}, \frac{3}{3}, \dots$$

Clearly, the fraction  $\frac{a}{b}$  in this enumeration occupies the  $a + \frac{b(b+1)}{2}$ -th place. For each integer  $k \geq 1$ , cover the  $k$ -th fraction  $\frac{a}{b}$  by an interval of length  $2^{-k}$  centred at  $\frac{a}{b}$ . Show that the point  $\frac{\sqrt{2}}{2}$  does not belong to any interval in the cover.

# Chapter 4

## Series

### 4.1 Convergence and Divergence of Series

**227 Definition** Let  $\{a_n\}_{n=1}^{+\infty}$  be a sequence of real numbers. A *series* is the sum of a sequence. We write

$$s_n = a_1 + a_2 + \cdots + a_n = \sum_{k=1}^n a_k.$$

Here  $s_n$  is the  $n$ -th *partial sum*. Observe in particular that

$$a_n = s_n - s_{n-1}.$$

**228 Definition** If the sequence  $\{s_n\}_{n=1}^{+\infty}$  has a finite limit  $S$ , we say that the series converges to  $S$  and write

$$\sum_{k=1}^{+\infty} a_k = \lim_{n \rightarrow +\infty} s_n = S.$$

Otherwise we say that the series *diverges*.

Observe that  $\sum_{n=1}^{+\infty} a_n$  converges to  $S$  if  $\forall \epsilon > 0, \exists N$  such that  $\forall n \geq N$ ,

$$\left| \left( \sum_{k \leq n} a_k \right) - S \right| = |s_n - S| < \epsilon.$$

Now, since

$$\left( \sum_{k \leq n} a_k \right) - S = \left( \sum_{k \leq n} a_k \right) - \left( \sum_{k \geq 1} a_k \right) = \sum_{k > n} a_k,$$

we see that a series converges if and only if its “tail” can be made arbitrarily small. Hence, the reader should notice that adding or deleting a finite amount of terms to a series does not affect its convergence or divergence. Furthermore, since the sequence of partial sums of a convergent series must be a Cauchy sequence we deduce that a series is convergent if and only if  $\forall \epsilon > 0, \exists N > 0$  such that  $\forall m \geq N, n \geq N, m \leq n$ ,

$$|s_n - s_m| = \left| \sum_{k=m}^n a_k \right| < \epsilon. \tag{4.1}$$

**229 THEOREM ( $n$ -th Term Test for Divergence)** If  $\sum_{n=1}^{\infty} a_n$  converges, then  $a_n \rightarrow 0$  as  $n \rightarrow +\infty$ .

**Proof:** Put  $s_n = \sum_{k=1}^n a_k$ . Then

$$\lim_{n \rightarrow +\infty} s_n = S \implies a_n = s_n - s_{n-1} \rightarrow S - S = 0.$$

□

In general, the problem of determining whether a series converges or diverges requires some work and it will be dealt with in the subsequent sections. We continue here with some other examples.

**230 Example** The series  $\sum_{n=1}^{+\infty} \left(1 + \frac{2}{n}\right)^n$  diverges, since its  $n$ -th term  $\left(1 + \frac{2}{n}\right)^n \rightarrow e^2$ .

**231 Example** We will prove that the *harmonic series*  $\sum_{n=1}^{+\infty} \frac{1}{n}$  diverges, even though  $\frac{1}{n} \rightarrow 0$  as  $n \rightarrow +\infty$ . Thus the condition in Theorem 229 though necessary for convergence is not sufficient. The divergence of the harmonic series was first demonstrated by Nicole d'Oresme (ca. 1323-1382), but his proof was mislaid for several centuries. The result was proved again by Pietro Mengoli in 1647, by Johann Bernoulli in 1687, and by Jakob Bernoulli shortly thereafter. Write the partial sums in dyadic blocks,

$$\sum_{n=1}^{2^M} \frac{1}{n} = \sum_{m=1}^M \sum_{n=2^{m-1}+1}^{2^m} \frac{1}{n}.$$

As  $1/n \geq 1/N$  when  $n \leq N$ , we deduce that

$$\sum_{n=2^{m-1}+1}^{2^m} \frac{1}{n} \geq \sum_{n=2^{m-1}+1}^{2^m} 2^{-m} = (2^m - 2^{m-1})2^{-m} = \frac{1}{2}$$

Hence,

$$\sum_{n=1}^{2^M} \frac{1}{n} \geq \frac{M}{2}$$

so the series diverges in the limit  $M \rightarrow +\infty$ .

The following theorem says that linear combinations of convergent series converge.

**232 THEOREM** Let  $\sum_{n=1}^{+\infty} a_n = A$  and  $\sum_{n=1}^{+\infty} b_n = B$  be convergent series and let  $\gamma \in \mathbb{R}$  be a real number. Then the series  $\sum_{n=1}^{+\infty} (a_n + \gamma b_n)$  converges to  $A + \gamma B$ .

**Proof:** For all  $\varepsilon > 0$  there exist  $N, N'$  such that for all  $n \geq \max(N, N')$ ,

$$\left| \sum_{k \leq n} a_k - A \right| < \frac{\varepsilon}{2}, \quad \left| \sum_{k \leq n} b_k - B \right| < \frac{\varepsilon}{2(|\gamma| + 1)}.$$

Hence, by the triangle inequality and by the obvious inequality  $\frac{|\gamma|}{|\gamma| + 1} \leq 1$ , we have

$$\left| \left( \sum_{k \leq n} a_k + \gamma b_k \right) - (A + \gamma B) \right| \leq \left| \sum_{k \leq n} a_k - A \right| + |\gamma| \left| \sum_{k \leq n} b_k - B \right| \leq \frac{\varepsilon}{2} + |\gamma| \frac{\varepsilon}{2(|\gamma| + 1)} < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon.$$

□

**233 Definition** A *geometric series* with common ratio  $r$  and first term  $a$  is one of the form

$$a + ar + ar^2 + ar^3 + \dots = \sum_{n=0}^{+\infty} ar^n.$$

By Theorem 173, if  $|r| < 1$  then the series converges and we have

$$a + ar + ar^2 + ar^3 + \dots = \sum_{n=0}^{+\infty} ar^n = \frac{a}{1-r}.$$

**234 Example** A fly starts at the origin and goes 1 unit up, 1/2 unit right, 1/4 unit down, 1/8 unit left, 1/16 unit up, etc., *ad infinitum*. In what coordinates does it end up?

**Solution:** Its *x* coordinate is

$$\frac{1}{2} - \frac{1}{8} + \frac{1}{32} - \dots = \frac{\frac{1}{2}}{1 - \frac{-1}{4}} = \frac{2}{5}.$$

Its *y* coordinate is

$$1 - \frac{1}{4} + \frac{1}{16} - \dots = \frac{1}{1 - \frac{-1}{4}} = \frac{4}{5}.$$

Therefore, the fly ends up in

$$\left(\frac{2}{5}, \frac{4}{5}\right).$$

Here we have used the fact the sum of an infinite geometric progression with common ratio *r*, with  $|r| < 1$  and first term *a* is

$$a + ar + ar^2 + ar^3 + \dots = \frac{a}{1 - r}.$$

**235 Definition** A *telescoping sum* is a sum where adjacent terms cancel out. That is,  $\sum_{n=0}^N a_n$  is a telescoping sum if we can write  $a_n = b_{n+1} - b_n$  and then

$$\sum_{n=0}^N a_n = a_0 + a_1 + \dots + a_N = (b_1 - b_0) + (b_2 - b_1) + \dots + (b_{N+1} - b_N) = b_{N+1} - b_0.$$

**236 Example** We have

$$\sum_{n=1}^N \frac{1}{n(n+1)} = \sum_{n=1}^N \left(\frac{1}{n} - \frac{1}{n+1}\right) = \left(\frac{1}{1} - \frac{1}{2}\right) + \left(\frac{1}{2} - \frac{1}{3}\right) + \dots + \left(\frac{1}{N} - \frac{1}{N+1}\right) = 1 - \frac{1}{N+1}.$$

Thus

$$\sum_{n=1}^{+\infty} \frac{1}{n(n+1)} = \lim_{N \rightarrow +\infty} \sum_{n=1}^N \frac{1}{n(n+1)} = \lim_{N \rightarrow +\infty} \left(1 - \frac{1}{N+1}\right) = 1.$$

**237 Example** We have

$$\begin{aligned} \sum_{n=1}^N \frac{1}{n(n+1)(n+2)} &= \frac{1}{2} \sum_{n=1}^N \left(\frac{1}{n(n+1)} - \frac{1}{(n+1)(n+2)}\right) = \frac{1}{2} \left(\left(\frac{1}{1 \cdot 2} - \frac{1}{2 \cdot 3}\right) + \left(\frac{1}{2 \cdot 3} - \frac{1}{3 \cdot 4}\right) + \dots + \left(\frac{1}{N(N+1)} - \frac{1}{(N+1)(N+2)}\right)\right) \\ &= \frac{1}{2} \left(\frac{1}{2} - \frac{1}{(N+1)(N+2)}\right). \end{aligned}$$

Thus

$$\sum_{n=1}^{+\infty} \frac{1}{n(n+1)(n+2)} = \lim_{N \rightarrow +\infty} \sum_{n=1}^N \frac{1}{n(n+1)(n+2)} = \lim_{N \rightarrow +\infty} \frac{1}{2} \left(\frac{1}{2} - \frac{1}{(N+1)(N+2)}\right) = \frac{1}{4}.$$

## Homework

**Problem 4.1.1** Find the sum of  $\sum_{n=3}^{\infty} \frac{2^n}{e^{n+1}}$ .

$$\sum_{n=0}^{+\infty} \arctan \frac{1}{n^2 + n + 1}.$$

**Problem 4.1.2** Find the sum of the series  $\sum_{n=2}^{+\infty} \frac{1}{4n^2 - 1}$ .

**Problem 4.1.4** Find the exact numerical value of the infinite sum

$$\sum_{n=1}^{+\infty} \frac{\sqrt{(n-1)!}}{(1 + \sqrt{1}) \cdots (1 + \sqrt{n})}.$$

**Problem 4.1.3** Find the exact numerical value of the sum



**Problem 4.1.5** Show that

$$\sum_{k=1}^n \frac{k}{k^4 + k^2 + 1} = \frac{1}{2} \cdot \frac{n^2 + n}{n^2 + n + 1},$$

and thus prove that  $\sum_{k=1}^n \frac{k}{k^4 + k^2 + 1}$  converges.

**Problem 4.1.6** Let  $b(n)$  denote the number of ones in the binary expansion of the positive integer  $n$ , for example  $b(3) = b(11_2) = 2$ .

Prove that  $\sum_{n=1}^{\infty} \frac{b(n)}{n(n+1)} = \log 4$ .

**Problem 4.1.7** Find

$$1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{6} + \frac{1}{8} + \frac{1}{9} + \frac{1}{12} + \frac{1}{16} + \frac{1}{18} + \dots,$$

which is the sum of the reciprocals of all positive integers of the form  $2^n 3^m$  for integers  $n \geq 0, m \geq 0$ .

**Problem 4.1.8** The Fibonacci Numbers  $f_n$  are defined recursively as follows:

$$f_0 = 1, \quad f_1 = 1, \quad f_{n+2} = f_n + f_{n+1}, \quad n \geq 0.$$

Prove that  $\sum_{n=1}^{+\infty} \frac{f_n}{3^n} = \frac{3}{5}$ .

**Problem 4.1.9** Let  $\sum_{n \geq 0} a_n$  be a convergent series and let  $\sum_{n \geq 0} b_n$  be a divergent series. Prove that  $\sum_{n \geq 0} (a_n + b_n)$  diverges.

**Problem 4.1.10** Prove that if  $\sum_{n \geq 1} a_n$  is a series of positive terms and that its partial sums are bounded, then  $\sum_{n \geq 1} a_n$  converges. Show that this is not necessarily true if  $\sum_{n \geq 1} a_n$  is not a series of positive terms.

## 4.2 Convergence and Divergence of Series of Positive Terms

We have several tools to establish convergence and divergence of series of positive terms. We will start with some simple comparison tests.


**238 THEOREM (Direct Comparison Test)** Let  $\{a_n\}_{n=0}^{+\infty}, \{b_n\}_{n=0}^{+\infty}, \{c_n\}_{n=0}^{+\infty}$  be sequences of positive real numbers. Suppose that eventually  $a_n \leq b_n$ , that is, that  $\exists N \geq 0$  such that  $\forall n \geq N$  there holds  $a_n \leq b_n$ . If  $\sum_{n \geq 0} b_n$  converges, then  $\sum_{n \geq 0} a_n$  converges.

If eventually  $a_n \geq c_n$ , and  $\sum_{n \geq 0} c_n$  diverges, then  $\sum_{n \geq 0} a_n$  also must diverge.

**Proof:** The theorem is clear from the inequalities

$$\sum_{n \geq N} a_n \leq \sum_{n \geq N} b_n, \quad \sum_{n \geq N} a_n \geq \sum_{n \geq N} c_n.$$

If  $\sum_{n \geq 0} b_n$  converges, then its tail can be made as small as we please, and so the tail of  $\sum_{n \geq 0} a_n$  can be made as small as we please. Similarly if  $\sum_{n \geq 0} c_n$  diverges, because it is a series of positive terms, its tail grows without bound and so the tail of  $\sum_{n \geq 0} a_n$  grows without bound.  $\square$

 Call a divergent series of positive terms a “giant” and a converging series of positive terms a “midget.” The comparison tests say that if a series is bigger than a giant it must be a giant, and if a series is smaller than a midget, it must be a midget.

**239 Example** From example 236,  $\sum_{n \geq 1} \frac{1}{n(n+1)}$  converges. Since for  $n \geq 1$ ,

$$n(n+1) < (n+1)^2 \implies \frac{1}{(n+1)^2} < \frac{1}{n(n+1)},$$

we deduce that the series

$$\sum_{n \geq 1} \frac{1}{(n+1)^2} = \sum_{n \geq 2} \frac{1}{n^2}$$

converges. Since adding a finite amount of terms to a series does not affect convergence, we deduce that  $1 + \sum_{n \geq 2} \frac{1}{n^2} = \sum_{n \geq 1} \frac{1}{n^2}$  converges.

**240 Example**  $\sum_{n=1}^{+\infty} \frac{1}{n^n}$  converges. For  $n \geq 2$  we have  $\frac{1}{n^n} \leq \frac{1}{n^2}$  and the series converges by direct comparison with  $\sum_{n=1}^{+\infty} \frac{1}{n^2}$ .

**241 Example** From example 230,  $\sum_{n \geq 1} \frac{1}{n}$  diverges. Since for  $n \geq 1$ ,  $\log n < n$ , we deduce that  $\sum_{n \geq 2} \frac{1}{\log n}$  diverges. Notice that here we start the sum at  $n = 2$  since the logarithm vanishes at  $n = 1$ .

**242 Example** Prove that

$$\sum_{\substack{p \\ \text{prime}}} \frac{1}{p}$$

diverges.

**Solution:** We will prove this by contradiction. Let  $p_1 = 2, p_2 = 3, p_3 = 5, \dots$  be the sequence of primes in ascending order and assume that the series converges. Then there exists an integer  $K$  such that

$$\sum_{m \geq K+1} \frac{1}{p_m} < \frac{1}{2}.$$

Let  $P = p_1 p_2 \cdots p_K$  and consider the numbers  $1 + nP$  for  $n = 1, 2, 3, \dots$ . None of these numbers has a prime divisor in the set  $\{p_1, p_2, \dots, p_K\}$  and hence all the prime divisors of the  $1 + nP$  belong to the set  $\{p_{K+1}, p_{K+2}, \dots\}$ . This means that for each  $t \geq 1$ ,

$$\sum_{n=1}^t \frac{1}{1+nP} \leq \sum_{s=1}^t \left( \sum_{m \geq K+1} \frac{1}{p_m} \right)^s \leq \sum_{s=1}^t \frac{1}{2^s} = 1,$$

that is,  $\sum_{n=1}^t \frac{1}{1+nP}$ , a series of positive terms, has bounded partial sums and so it converges. But since  $1+nP \sim nP$  as  $n \rightarrow +\infty$  and  $\frac{1}{P} \sum_{n=1}^t \frac{1}{n}$  diverges, we obtain a contradiction.

Since the convergent behaviour of a series depends of its tail, the following asymptotic comparison tests should be clear, and its proof follows the same line of reasoning as Theorem 238.

**243 THEOREM (Asymptotic Comparison Test)** Let  $\{a_n\}_{n=0}^{+\infty}, \{b_n\}_{n=0}^{+\infty}, \{c_n\}_{n=0}^{+\infty}$  be sequences of real numbers which are eventually positive. Suppose that  $a_n \ll b_n$ , and that  $c_n \ll a_n$ . Then both  $\sum_{n=0}^{\infty} a_n$  and  $\sum_{n=0}^{\infty} b_n$  converge together, and both  $\sum_{n=0}^{\infty} a_n$  and  $\sum_{n=0}^{\infty} c_n$  diverge together. Moreover, if  $\{b_n\}_{n=0}^{+\infty}$  is eventually a strictly positive sequence and  $a_n \sim b_n$ , then  $\sum_{n=0}^{\infty} a_n$  and  $\sum_{n=0}^{\infty} b_n$  converge or diverge together.

In order to effectively use the comparison tests we must have a ready catalogue of series whose convergence or divergence we know. In the subsequent lines we will develop such a catalogue. We start with the following consequence of the comparison tests.

**244 THEOREM (Cauchy Condensation Test)** Let  $\{a_n\}_{n=0}^{+\infty}$  be a sequence of positive real numbers which is monotonically decreasing. Then  $\sum_{n=0}^{\infty} a_n$  converges if and only if the sum  $\sum_{n=0}^{\infty} 2^n a_{2^n}$  converges.

**Proof:** Since the sequence  $\{a_n\}_{n=0}^{+\infty}$  is monotonically decreasing and positive,

$$\sum_{k=2^n}^{2^{n+1}-1} a_{2^{n+1}-1} \leq \sum_{k=2^n}^{2^{n+1}-1} a_k \leq \sum_{k=2^n}^{2^{n+1}-1} a_{2^n} \implies 2^n a_{2^{n+1}-1} \leq \sum_{k=2^n}^{2^{n+1}-1} a_k \leq 2^n a_{2^n}.$$

The second inequality yields

$$\sum_{n=0}^{2^{N+1}-1} a_n = \sum_{n=0}^N \sum_{k=2^n}^{2^{n+1}-1} a_k \leq \sum_{n=0}^N 2^n a_{2^n} \implies \lim_{N \rightarrow +\infty} \sum_{n=0}^{2^{N+1}-1} a_n \leq \lim_{N \rightarrow +\infty} \sum_{n=0}^N 2^n a_{2^n}.$$

Thus if  $\sum_{n=0}^{+\infty} 2^n a_{2^n}$  converges so does  $\sum_{n=0}^{+\infty} a_n$ .

The first inequality yields

$$\begin{aligned} 2^n a_{2^{n+1}-1} &\leq \sum_{k=2^n}^{2^{n+1}-1} a_k \implies (2^{n+1}-1) a_{2^{n+1}-1} \leq 2 \sum_{k=2^n}^{2^{n+1}-1} a_k - a_{2^{n+1}-1} \\ &\implies \sum_{n=0}^N (2^{n+1}-1) a_{2^{n+1}-1} \leq 2 \sum_{n=0}^N \sum_{k=2^n}^{2^{n+1}-1} a_k - \sum_{n=0}^N a_{2^{n+1}-1} = 2 \sum_{n=0}^{2^{N+1}-1} a_n - \sum_{n=0}^N a_{2^{n+1}-1}. \end{aligned}$$

□

As an application of Cauchy's Test, we obtain the following important result.

**245 THEOREM (p-series Test)** If  $p > 1$  then  $\zeta(p) = \sum_{n=1}^{+\infty} \frac{1}{n^p}$  converges, but diverges when  $p \leq 1$ .

**Proof:** If  $p \leq 0$ , divergence follows from Theorem 229. If  $p > 0$ , then using the fact that  $x \mapsto x^p$  is monotonically increasing, we may use Theorem 244. Since

$$\sum_{k \geq 0} \frac{2^k}{2^{pk}} = \sum_{k \geq 0} \left(2^{(1-p)}\right)^k$$

is a geometric series with ratio  $2^{1-p}$ , it converges by Theorem 173 when

$$2^{1-p} < 1 \implies (1-p)\log_2 2 < \log_2 1 \implies 1-p < 0 \implies p > 1,$$

and diverges for  $p > 1$ . The case  $p = 1$  has been shown to diverge in example 230. □

**246 Example** Since  $\sqrt{2} > 1$ , the series  $\sum_{n=1}^{+\infty} \frac{1}{n^{\sqrt{2}}}$  converges.

**247 Example** Since

$$\frac{n^{\sqrt{2}} + (\log \log n)^{2007}}{n^3 + n(\log n)^5 + 1} \sim \frac{n^{\sqrt{2}}}{n^3} = \frac{1}{n^{3-\sqrt{2}}}$$

and  $3 - \sqrt{2} > 1$ , the series  $\sum_{n \geq 1} \frac{n^{\sqrt{2}} + (\log \log n)^{2007}}{n^3 + n(\log n)^5 + 1}$  converges.

**248 COROLLARY (De Morgan's Logarithmic Scale)** If  $p > 1$  then all of

$$\sum_{n=1}^{+\infty} \frac{1}{n^p}; \quad \sum_{n \geq e} \frac{1}{n(\log n)^p}; \quad \sum_{n \geq e^e} \frac{1}{n(\log n)(\log \log n)^p}; \quad \sum_{n \geq e^{e^e}} \frac{1}{n(\log n)(\log \log n)(\log \log \log n)^p}; \quad \dots$$

converge, but diverge when  $p \leq 1$ .

**Proof:** The theorem is proved inductively by successive applications of Cauchy's Condensation Test. We will prove how the case for  $\sum_{n \geq e} \frac{1}{n(\log n)^p}$  follows from the case  $\sum_{n=1}^{+\infty} \frac{1}{n^p}$  and leave the rest to the reader. We see that

$$\sum_{k \geq 1} \frac{2^k}{2^k (\log 2^k)^p} = \frac{1}{(\log 2)^p} \sum_{k \geq 1} \frac{1}{k^p},$$

and so this case follows from Theorem 245. □

**249 Example** Determine whether  $\sum_{n=4}^{+\infty} \frac{(\log n)^{100}}{n^{3/2} \log \log n}$  converges.

**Solution:** Since  $(\log n)^{100} \ll n^{1/4}$ , eventually  $\frac{(\log n)^{100}}{n^{1/4}} \ll 1$ . We have  $\frac{(\log n)^{100}}{n^{3/2} \log \log n} \ll \frac{(\log n)^{100}}{n^{1/4}} \cdot \frac{1}{n^{5/4} \log \log n}$  and since  $\sum_{n=4}^{+\infty} \frac{1}{n^{5/4} \log \log n} < +\infty$ , we have  $\sum_{n=4}^{+\infty} \frac{(\log n)^{100}}{n^{3/2} \log \log n} < +\infty$ , that is, the series converges.

The reader should be aware that the value of the exponent in Theorems 245 and 248 is fixed. The following examples should dissuade him that “having an exponent higher than 1” implies convergence.

**250 Example** Test  $\sum_{n=1}^{\infty} \frac{1}{n^{1+1/n}}$  for convergence by comparing it to a suitable  $p$ -series. Use the direct comparison test.

**Solution:** By induction  $n < 2^n \implies n^{1/n} < 2$  and so  $n^{1+1/n} < 2n \implies \frac{1}{2n} < \frac{1}{n^{1+1/n}}$ . So the series diverges by direct comparison to  $\sum_{n=1}^{\infty} \frac{1}{2n}$ .

**251 Example** Test  $\sum_{n=2}^{\infty} \frac{1}{n^{1+1/\log n}}$  for convergence by comparing it to a suitable  $p$ -series. Use the direct comparison test

**Solution:** We have  $n = e^{\log n} \implies n^{\frac{1}{\log n}} = e$  and so  $n^{1+1/\log n} = en$ ,  $n > 1$ . So the series diverges by direct comparison to  $\sum_{n=2}^{\infty} \frac{1}{en}$ .

**252 Example** Test  $\sum_{n=2}^{\infty} \frac{1}{n^{1+1/\log \log n}}$  for convergence by comparing it to a suitable  $p$ -series. Use the direct comparison test.

**Solution:** By considering the monotonicity of  $f(x) = e^x - \frac{x^2}{2}$  (see Theorem 385) or otherwise, we can prove that  $e^x > \frac{x^2}{2}$  for  $x > 0$ . Now,

$$n^{1/\log \log n} = e^{\log n^{1/\log \log n}} = e^{\frac{\log n}{\log \log n}} > \frac{(\log n)^2}{2(\log \log n)^2}.$$

This gives

$$\frac{2(\log \log n)^2}{n(\log n)^2} > \frac{1}{n^{1+\frac{1}{\log \log n}}}.$$

Now,

$$\sum_{n=2}^{+\infty} \frac{2(\log \log n)^2}{n(\log n)^2}$$

can be shown to converge by comparing to a series in the De Morgan logarithmic scale.

**253 Example** Prove that the series  $\sum_{n=1}^{+\infty} \frac{1}{n^{1.8+\sin n}}$  diverges.

**Solution:** For  $k \in \mathbb{Z}$ , the interval  $I_k = \left[ (2k + \frac{4}{3})\pi ; (2k + \frac{5}{3})\pi \right]$  has length  $\frac{\pi}{3} > 1$  and  $x \in I_k \implies \sin x \leq -\frac{\sqrt{3}}{2}$ .

The gap between  $I_k$  and  $I_{k+1}$  is  $< \frac{5\pi}{3} < 6$ . Hence, among any seven consecutive integers, at least one must fall

into  $I_k$  and for this value of  $n$  we must have  $1.8 + \sin n < 1 - \frac{\sqrt{3}}{2} < 1$ . This means that

$$\sum_{n=1}^{+\infty} \frac{1}{n^{1.8+\sin n}} = \sum_{m=0}^{+\infty} \sum_{n=7m+1}^{n=7m+7} \frac{1}{n^{1.8+\sin n}} \geq \sum_{m=0}^{+\infty} \frac{1}{7m+7},$$

and since the rightmost series diverges, the original series diverges by the direct comparison test.

The following result puts the harmonic series at the “frontier” of convergence and divergence for series with monotonically decreasing positive terms.

**254 THEOREM (Pringsheim’s Theorem)** Let  $\sum_{n \geq 1} a_n$  be a converging series of positive terms of monotonically decreasing terms. Then  $a_n = o\left(\frac{1}{n}\right)$ .

**Problem 4.2.1** Since the series converges, its sequence of partial sums is a Cauchy sequence and by 4.1, given  $\varepsilon > 0$ ,  $\exists m > 0$ , such that  $\forall n \geq m$ ,

$$\sum_{k=m+1}^n a_k < \varepsilon.$$

Because the series decreases monotonically, each of  $a_{m+1}, a_{m+2}, \dots, a_n$  is at least  $a_n$  and thus

$$(n-m)a_n \leq \sum_{k=m+1}^n a_k < \varepsilon.$$

Again, since the series converges,  $a_n \rightarrow 0$  as  $n \rightarrow +\infty$  we may choose  $n$  large enough so that  $a_n < \frac{\varepsilon}{m}$ . In this case

$$(n-m)a_n < \varepsilon \implies na_n < \varepsilon + ma_n < 2\varepsilon \implies a_n < \frac{2\varepsilon}{n},$$

which proves the theorem.

The disadvantage of the comparison tests is that in order test for convergence, we must appeal to the behaviour of an auxiliary series. The next few tests provide a way of testing the series against its own terms.

**255 THEOREM (Root Test)** Let  $\sum_{n=1}^{+\infty} a_n$  be a series of positive terms. Put  $r = \limsup (a_n)^{1/n}$ . Then the series converges if  $r < 1$  and diverges if  $r > 1$ . The test is inconclusive if  $r = 1$ .

**Proof:** If  $r < 1$  choose  $r'$  with  $r < r' < 1$ . Then there exists  $N \in \mathbb{N}$  such that

$$\forall n \geq N, \quad \sqrt[n]{a_n} \leq r' \implies a_n \leq (r')^n.$$

But then  $\sum_{n=0}^{+\infty} a_n$  converges by direct comparison to the converging geometric series  $\sum_{n=0}^{+\infty} (r')^n$ .

If  $r > 1$  then there is a sequence  $\{n_k\}_{k=1}^{+\infty}$  of positive integers such that

$$\sqrt[n_k]{a_{n_k}} \rightarrow r.$$

This means that  $a_n$  will be  $> 1$  for infinitely many values of  $n$ , and so, the condition  $a_n \rightarrow 0$  necessary for convergence, does not hold.

By considering  $\sum_{n=1}^{+\infty} \frac{1}{n}$ , which diverges, and  $\sum_{n=1}^{+\infty} \frac{1}{n^2}$ , which converges, one sees that  $r = 1$  may appear in series of different conditions.  $\square$

**256 THEOREM (Ratio Test)** Let  $\sum_{n=1}^{+\infty} a_n$  be a series of strictly positive terms. Put  $r = \limsup \frac{a_{n+1}}{a_n}$ . Then the series converges if  $r < 1$  and diverges if  $r > 1$ . The test is inconclusive if  $r = 1$ .

**Proof:** If  $r < 1$ , then there exists  $N \in \mathbb{N}$  such that

$$a_{N+1} \leq r a_N, \quad a_{N+2} \leq r a_{N+1}, \quad a_{N+3} \leq r a_{N+2}, \dots, \quad a_{N+t} \leq r a_{N+t-1}.$$

Multiplying all these inequalities together,

$$a_{N+t} \leq a_N r^t.$$

Putting  $N + t = n$  we deduce that

$$a_n \leq a_N r^{-N} r^n.$$

Since  $a_N r^{-N}$  is a constant, we may use direct comparison between  $\sum_{n=1}^{+\infty} a_n$  and the converging geometric series

$$a_N r^{-N} \sum_{n=1}^{+\infty} r^n, \text{ concluding that } \sum_{n=1}^{+\infty} a_n \text{ converges.}$$

If  $r > 1$  then  $a_{n+1} \geq a_n \geq a_N$  for all  $n \geq N$ . This means that the condition  $a_n \rightarrow 0$ , necessary for convergence, does not hold.

By considering  $\sum_{n=1}^{+\infty} \frac{1}{n}$ , which diverges, and  $\sum_{n=1}^{+\infty} \frac{1}{n^2}$ , which converges, one sees that  $r = 1$  may appear in series of different conditions.  $\square$



The root test is more general than the ratio test, as can be seen from Theorem 225.

**257 Example** Since

$$\frac{\frac{(n+1)!}{(n+1)^{n+1}}}{\frac{n!}{n^n}} = \frac{1}{\left(1 + \frac{1}{n}\right)^n} \rightarrow \frac{1}{e} < 1$$

the series  $\sum_{n=1}^{+\infty} \frac{n!}{n^n}$  converges.

**258 Example** Since

$$\left(\frac{(n!)^n}{n^{n^2}}\right)^{1/n} = \frac{n!}{n^n} \rightarrow 0$$

the series  $\sum_{n=1}^{+\infty} \frac{(n!)^n}{n^{n^2}}$  converges.

## Homework

**Problem 4.2.2** True or False: If the infinite series  $\sum_{n=1}^{+\infty} a_n$  of strictly positive terms, converges, then  $\sum_{n=1}^{+\infty} a_n^2$  must necessarily converge.

**Problem 4.2.3** True or False: If the infinite series  $\sum_{n=1}^{+\infty} a_n$  of strictly positive terms converges, then  $\sum_{n=1}^{+\infty} \sin(a_n)$  must necessarily converge.

**Problem 4.2.4** True or False: If the infinite series  $\sum_{n=1}^{+\infty} a_n$  of strictly positive terms converges, then  $\sum_{n=1}^{+\infty} \tan(a_n)$  must necessarily converge.

**Problem 4.2.5** True or False: If the infinite series  $\sum_{n=1}^{+\infty} a_n$  converges, then  $\sum_{n=1}^{+\infty} \cos(a_n)$  must necessarily converge.

**Problem 4.2.6** Use the comparison tests to show that if  $a_n > 0$  and  $\sum_{n=1}^{\infty} a_n$  converges, then  $\sum_{n=1}^{\infty} \frac{a_n}{n}$  converges.

**Problem 4.2.7** Give an example of a series converging to 1 with  $n$ -th term  $a_n > 0$  satisfying  $a_n \ll \frac{1}{n^2}$ . (That is, the  $n$ -th term goes to zero faster than the reciprocal of a square.)

**Problem 4.2.8** Give an example of a converging series of strictly positive terms  $\sum_{n=1}^{+\infty} a_n$  such that  $\sum_{n=1}^{+\infty} (a_n)^{1/n}$  also converges.

**Problem 4.2.9** Give an example of a converging series of strictly positive terms  $\sum_{n=1}^{+\infty} a_n$  such that  $\sum_{n=1}^{+\infty} (a_n)^{1/n}$  diverges.

**Problem 4.2.10** Give an example of a converging series of strictly positive terms  $a_n$  such that  $\lim_{n \rightarrow +\infty} (a_n)^{1/n}$  does not exist.

**Problem 4.2.11** Test  $\sum_{n=1}^{\infty} \frac{3^n}{n^{2n}}$  using both direct comparison and

the root test.

**Problem 4.2.12** Let  $\mathcal{S}$  be the set of positive integers none of whose digits in its decimal representation is a 0:

$$\mathcal{S} = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 11, 12, 13, 14, 15, 16, 17, 18, 19, 21, \dots\}.$$

Prove that the series  $\sum_{n \in \mathcal{S}} \frac{1}{n}$  converges.

**Problem 4.2.13** Let  $d(n)$  be the number of strictly positive divisors of the integer  $n$ . Prove that  $d(n) \leq 2\sqrt{n}$ . Use this to prove that

$$\sum_{n \geq 1} \frac{d(n)}{n^2}$$

converges.

**Problem 4.2.14** Let  $p_n$  be the  $n$ -th prime. Thus  $p_1 = 2$ ,  $p_2 = 3$ ,  $p_3 = 5$ , etc. Put  $a_1 = p_1$  and  $a_{n+1} = p_1 p_2 \cdots p_n + 1$  for  $n \geq 1$ . Find

$$\sum_{n=1}^{+\infty} \frac{1}{a_n}.$$

**Problem 4.2.15** Determine whether  $\sum_{n \geq 2} a_n$  converges, when  $a_n$  is given as below.

- |  |   |
|--|---|
| 1. $\left(1 + \frac{1}{n}\right)^n - e.$         | 7. $\frac{2 \cdot 4 \cdot 6 \cdots (2n)}{n^n}.$ |
| 2. $\cosh^\alpha n - \sinh^\alpha n.$            | 8. $\frac{1! + 2! + \cdots + n!}{(n+2)!}.$      |
| 3. $\log \frac{(n^3+1)^2}{(n^2+1)^3}.$           | 9. $\frac{1! - 2! + \cdots \pm n!}{(n+1)!}.$    |
| 4. $\sqrt[n]{n+1} - \sqrt[n]{n}.$                | 10. $\frac{(\log n)^n}{n^{\log n}}.$            |
| 5. $\arccos \left( \frac{n^3+1}{n^3+2} \right).$ | 11. $\frac{1}{(\log n)^{\log n}}.$              |
| 6. $\frac{a^n}{1+a^{2n}}.$                       |   |

### 4.3 Summation by Parts

In this section we consider series whose terms have arbitrary signs. We first need the following result.


**259 THEOREM (Summation by Parts)** Let  $A_n = \sum_{0 \leq k \leq n} a_k$ ,  $A_{-1} = 0$ . Then for  $p \leq q$ ,

$$\sum_{p \leq k \leq q} a_k b_k = \sum_{p \leq k \leq q-1} A_k (b_k - b_{k+1}) + A_q b_q - A_{p-1} b_p.$$

**Proof:** Changing a subindex,

$$\begin{aligned} \sum_{p \leq k \leq q} a_k b_k &= \sum_{p \leq k \leq q} (A_k - A_{k-1}) b_k \\ &= \sum_{p \leq k \leq q} A_k b_k - \sum_{p \leq k \leq q} A_{k-1} b_k \\ &= \sum_{p \leq k \leq q-1} A_k b_k - \sum_{p \leq k \leq q-1} A_k b_{k+1} + A_q b_q - A_{p-1} b_p. \end{aligned}$$

giving the result.  $\square$

 An alternative and more symmetric formulation will be given once we introduce Riemann-Stieltjes integration.

We now obtain a convergence test.

**260 THEOREM (Dirichlet's Test)** The series  $\sum_{k \geq 0} a_k b_k$  converges if

1. the partial sums  $A_n = \sum_{k=0}^n a_k$  are bounded
2.  $b_n \geq b_{n+1}$
3.  $b_n \rightarrow 0$  as  $n \rightarrow +\infty$

**Proof:**

$\square$

## 4.4 Alternating Series

A series of the form  $\sum_{n=1}^{+\infty} (-1)^n a_n$  where all the  $a_n \geq 0$  is called an *alternating series*.

**261 THEOREM (Leibniz's Alternating Series Test)** The alternating series  $\sum_{n=1}^{+\infty} (-1)^n a_n$  converges if all the following conditions are met

- the  $a_n$  eventually decrease, that is,  $a_{n+1} \leq a_n$  for all  $n \geq N$ .
- $a_n \rightarrow 0$

**262 Example** The series  $\sum_{n=1}^{+\infty} (-1)^{n+1} \frac{1}{n}$  converges by Leibniz's Test. In fact, one can prove that it equals  $\log 2$ .

## 4.5 Absolute Convergence

If  $\sum_{n=1}^{+\infty} |a_n|$  converges then  $\sum_{n=1}^{+\infty} a_n$  converges. The converse is not true.

**263 Example** Since  $\left| \frac{\sin n}{n^2} \right| \leq \frac{1}{n^2}$ ,  $\sum_{n=1}^{+\infty} \left| \frac{\sin n}{n^2} \right|$  converges by the comparison test. Thus  $\sum_{n=1}^{+\infty} \frac{\sin n}{n^2}$  converges absolutely and so it converges.

**264 Example** Determine whether the following two infinite series converge: (I)  $\sum_{n=2}^{\infty} \frac{(-1)^n \sin(3n)}{n^2}$ , (II)  $\sum_{n=1}^{\infty} \frac{(-1)^n n}{n^2 + 2}$ .

**Solution:** We have

$$\left| (-1)^n \frac{\sin 3n}{n^2} \right| \leq \frac{1}{n^2},$$

so (I) converges absolutely. As for number (II),  $f(x) = \frac{x}{x^2 + 2}$  is decreasing (take the first derivative)  $\frac{n}{n^2 + 2} \rightarrow 0$ , so it converges by Leibniz's Test.



# Chapter 5

## Real Functions of One Real Variable

### 5.1 Limits of Functions

**265 DEFINITION-PROPOSITION (Cauchy-Heine, Sinistral Limit)** Let  $f : ]a; b[ \rightarrow \mathbb{R}$  and let  $x_0 \in ]a; b[$ . The following are equivalent.

1.  $\forall \varepsilon > 0, \exists \delta > 0$  such that

$$x_0 - \delta < x < x_0 \implies |f(x) - L| < \varepsilon.$$

2. For each sequence  $\{x_n\}_{n=1}^{+\infty}$  of points in the interval  $]a; b[$  with  $x_n < x_0, x_n \rightarrow x_0 \implies f(x_n) \rightarrow L$ .

If either condition is fulfilled we say that  $f$  has a sinistral limit  $f(x_0-)$  as  $x$  increases towards  $x_0$  and we write

$$f(x_0-) = \lim_{x \rightarrow x_0-} f(x) = \lim_{x/x_0} f(x).$$

**Proof:**

$1 \implies 2$  Suppose that  $\forall \varepsilon > 0, \exists \delta > 0$  such that

$$x_0 - \delta < x < x_0 \implies |f(x) - L| < \varepsilon.$$

Let  $x_n < x_0, x_n \rightarrow x_0$ . Then

$$|x_n - x_0| < \delta \implies x_0 - \delta < x_n < x_0 + \delta$$

for sufficiently large  $n$ . But we are assuming that  $x_n < x_0$ , so in fact we have  $x_0 - \delta < x_n < x_0$ . By our assumption then  $|f(x_n) - L| < \varepsilon$ , and so  $1 \implies 2$ .

$2 \implies 1$  Suppose that for each sequence  $\{x_n\}_{n=1}^{+\infty}$  of points in the interval  $]a; b[$  with  $x_n < x_0, x_n \rightarrow x_0 \implies f(x_n) \rightarrow L$ . If it were not true that  $f(x) \rightarrow L$  as  $x \rightarrow x_0$ , then there exists some  $\varepsilon_0 > 0$  such that for all  $\delta > 0$  we can find  $x$  such that

$$0 < |x - x_0| < \delta \implies |f(x) - L| \geq \varepsilon_0.$$

In particular, for each strictly positive integer  $n$  we can find  $x_n$  satisfying

$$0 < |x_n - x_0| < \frac{1}{n} \implies |f(x_n) - L| \geq \varepsilon_0,$$

a contradiction to the fact that  $f(x_n) \rightarrow L$ .

□

In an analogous manner, we have the following.

**266 DEFINITION-PROPOSITION (Cauchy-Heine, Dextral Limit)** Let  $f : ]a; b[ \rightarrow \mathbb{R}$  and let  $x_0 \in ]a; b[$ . The following are equivalent.

1. For each sequence  $\{x_n\}_{n=1}^{+\infty}$  of points in the interval  $]a; b[$  with  $x_n > x_0$ ,  $x_n \rightarrow x_0 \implies f(x_n) \rightarrow L$ .
2.  $\forall \varepsilon > 0, \exists \delta > 0$  such that

$$x_0 < x < x_0 + \delta \implies |f(x) - L| < \varepsilon.$$

If either condition is fulfilled we say that  $f$  has a dextral limit  $f(x_0+)$  as  $x$  decreases towards  $x_0$  and we write

$$f(x_0+) = \lim_{x \rightarrow x_0+} f(x) = \lim_{x \searrow x_0} f(x).$$

Upon combining Propositions 265 and 266 we obtain the following.

**267 DEFINITION-PROPOSITION (Cauchy-Heine)** Let  $f: ]a; b[ \rightarrow \mathbb{R}$  and let  $x_0 \in ]a; b[$ . The following are equivalent.

1.  $f(x_0-) = f(x_0+)$
2. For each sequence  $\{x_n\}_{n=1}^{+\infty}$  of points in the interval  $]a; b[$  different from  $x_0$ ,  $x_n \rightarrow x_0 \implies f(x_n) \rightarrow L$ .
3.  $\forall \varepsilon > 0, \exists \delta > 0$  such that

$$0 < |x - x_0| < \delta \implies |f(x) - L| < \varepsilon.$$

If either condition is fulfilled we say that  $f$  has a (two-sided) limit  $L$  as  $x$  decreases towards  $x_0$  and we write

$$L = \lim_{x \rightarrow x_0} f(x).$$

We now prove analogues of the theorems that the proved for limits of sequences.

**268 THEOREM (Uniqueness of Limits)** Let  $X \subseteq \mathbb{R}$ ,  $a \in \mathbb{R}$ , and  $f: X \rightarrow \mathbb{R}$ . If  $\lim_{x \rightarrow a} f(x) = L$  and  $\lim_{x \rightarrow a} f(x) = L'$  then  $L = L'$ .

**Proof:** If  $L \neq L'$  then take  $2\varepsilon = |L - L'|$  in the definition of limit. There is  $\delta > 0$  such that

$$0 < |x - a| < \delta \implies |f(x) - L| < \frac{|L - L'|}{2}, \quad |f(x) - L'| < \frac{|L - L'|}{2}.$$

By the Triangle Inequality

$$|L - L'| \leq |L - f(x)| + |f(x) - L'| < \frac{|L - L'|}{2} + \frac{|L - L'|}{2} = |L - L'|,$$

but  $|L - L'| < |L - L'|$  is a contradiction.  $\square$

**269 THEOREM (Local Boundedness)** Let  $X \subseteq \mathbb{R}$ ,  $a \in \mathbb{R}$ , and  $f: X \rightarrow \mathbb{R}$ . If  $\lim_{x \rightarrow a} f(x) = L$  exists and is finite, then  $f$  is bounded in a neighbourhood of  $a$ .

**Proof:** Take  $\varepsilon = 1$  in the definition of limit. Then there is a  $\delta > 0$  such that

$$0 < |x - a| < \delta \implies |f(x) - L| < 1 \implies |f(x)| < 1 + |L|,$$

and so  $f$  is bounded on this neighbourhood.  $\square$

**270 THEOREM (Order Properties of Limits)** Let  $X \subseteq \mathbb{R}$ ,  $a \in \mathbb{R}$ , and  $f: X \rightarrow \mathbb{R}$ . Let  $\lim_{x \rightarrow a} f(x) = L$  exist and be finite. Then

1. If  $s < L$  then there exists a neighbourhood  $\mathcal{N}_a$  of  $a$  contained in  $X$  such that  $\forall x \in \mathcal{N}_a, s < f(x)$ .
2. If  $L < t$  then there exists a neighbourhood  $\mathcal{N}_a$  of  $a$  contained in  $X$  such that  $\forall x \in \mathcal{N}_a, f(x) < t$ .
3. If  $s < L < t$  then there exists a neighbourhood  $\mathcal{N}_a$  of  $a$  contained in  $X$  such that  $\forall x \in \mathcal{N}_a, s < f(x) < t$ .
4. If there exists a neighbourhood  $\mathcal{N}_a \subseteq X$  such that  $\forall x \in \mathcal{N}_a, s \leq f(x)$ , then  $s \leq L$ .

5. If there exists a neighbourhood  $\mathcal{N}_a \subseteq X$  such that  $\forall x \in \mathcal{N}_a, f(x) \leq t$ , then  $L \leq t$ .
6. If there exists a neighbourhood  $\mathcal{N}_a \subseteq X$  such that  $\forall x \in \mathcal{N}_a, s \leq f(x) \leq t$ , then  $s \leq L \leq t$ .

**Proof:** We have

1. Take  $\varepsilon = L - s > 0$  in the definition of limit. There is  $\delta > 0$  such that

$$0 < |x - a| < \delta \implies |f(x) - L| < L - s \implies s - L + L < f(x) < 2L - s \implies s < f(x),$$

as claimed.

2. Take  $\varepsilon = t - L > 0$  in the definition of limit. There is  $\delta > 0$  such that

$$0 < |x - a| < \delta \implies |f(x) - L| < t - L \implies L - t + L < f(x) < t - L + L \implies f(x) < t,$$

as claimed.

3. This follows by (1) and (2).
4. If on the said neighbourhood  $\mathcal{N}_a$  we had, on the contrary,  $L > s$  then (1) asserts that there is a neighbourhood of  $\mathcal{N}'_a \subseteq \mathcal{N}_a$  such that  $f(x) > s$ , a contradiction to the assumption that  $\forall x \in \mathcal{N}_a, s \leq f(x)$ .
5. If on the said neighbourhood  $\mathcal{N}_a$  we had, on the contrary,  $L < t$  then (2) asserts that there is a neighbourhood of  $\mathcal{N}'_a \subseteq \mathcal{N}_a$  such that  $f(x) < t$ , a contradiction to the assumption that  $\forall x \in \mathcal{N}_a, t \geq f(x)$ .
6. This follows by (4) and (5).

□

Analogous to the Sandwich Theorem for sequences we have

**271 THEOREM (Sandwich Theorem)** Assume that  $a, b, c$  are functions defined on a neighbourhood  $\mathcal{N}_{x_0}$  of a point  $x_0$  except possibly at  $x_0$  itself. Assume moreover that in  $\mathcal{N}_{x_0}$  they satisfy the inequalities  $a(x) \leq b(x) \leq c(x)$ . Then

$$\lim_{x \rightarrow x_0} a(x) = L = \lim_{x \rightarrow x_0} c(x) \implies \lim_{x \rightarrow x_0} b(x) = L.$$

**Proof:** For all  $\varepsilon > 0$  there is  $\delta > 0$  such that

$$0 < |x - x_0| < \delta \implies |a(x) - L| < \varepsilon \quad \text{and} \quad |c(x) - L| < \varepsilon \implies L - \varepsilon < a(x) < L + \varepsilon \quad \text{and} \quad L - \varepsilon < c(x) < L + \varepsilon.$$

If we now consider  $x \in \mathcal{N}_{x_0} \cap \{x : 0 < |x - x_0| < \delta\}$  then

$$L - \varepsilon < a(x) \leq b(x) \leq c(x) < L + \varepsilon \implies L - \varepsilon < b(x) < L + \varepsilon \implies |b(x) - L| < \varepsilon,$$

whence  $\lim_{x \rightarrow x_0} b(x) = L$ . □

**272 THEOREM** Let  $X \subseteq \mathbb{R}$ ,  $a \in \mathbb{R}$ , and  $f, g : X \rightarrow \mathbb{R}$ . Let  $(L, L', \lambda) \in \mathbb{R}^3$ . Then

1.  $\lim_{x \rightarrow a} f(x) = L \implies \lim_{x \rightarrow a} |f(x)| = |L|$ .
2.  $\lim_{x \rightarrow a} f(x) = 0 \iff \lim_{x \rightarrow a} |f(x)| = 0$ .
3.  $\lim_{x \rightarrow a} f(x) = L, \lim_{x \rightarrow a} g(x) = L' \implies \lim_{x \rightarrow a} (f(x) + \lambda g(x)) = L + \lambda L'$ .
4.  $\lim_{x \rightarrow a} f(x) = L, \lim_{x \rightarrow a} g(x) = L' \implies \lim_{x \rightarrow a} (f(x)g(x)) = LL'$ .
5. If  $\lim_{x \rightarrow a} f(x) = 0$  and if  $g$  is bounded on a neighbourhood  $\mathcal{N}_a$  of  $a$ , then  $\lim_{x \rightarrow a} f(x)g(x) = 0$ .
6.  $\lim_{x \rightarrow a} f(x) = L, \lim_{x \rightarrow a} g(x) = L' \neq 0 \implies \lim_{x \rightarrow a} \left( \frac{f(x)}{g(x)} \right) = \frac{L}{L'}$ .

**Proof:**

1. This follows from the inequality  $||f(x)| - |L|| \leq |f(x) - L|$ .
2. This follows from the inequalities  $-|f(x)| \leq f(x) \leq |f(x)|$  and  $\min(-f(x), f(x)) \leq |f(x)| \leq \max(-f(x), f(x))$  and the Sandwich Theorem.
3. For all  $\epsilon > 0$  there are  $\delta_1 > 0$  and  $\delta_2 > 0$  such that

$$0 < |x - a| < \delta_1 \implies |f(x) - L| < \epsilon, \quad \text{and} \quad 0 < |x - a| < \delta_2 \implies |g(x) - L'| < \epsilon.$$

Take  $\delta = \min(\delta_1, \delta_2)$ . Then

$$0 < |x - a| < \delta \implies |f(x) + \lambda g(x) - (L + \lambda L')| \leq |f(x) - L| + |\lambda| |g(x) - L'| < (1 + |\lambda|)\epsilon.$$

Since the dextral side can be made arbitrarily small, the assertion follows.

4. For all  $\epsilon > 0$  there are  $\delta_1 > 0$  and  $\delta_2 > 0$  such that

$$0 < |x - a| < \delta_1 \implies |f(x) - L| < \epsilon, \quad \text{and} \quad 0 < |x - a| < \delta_2 \implies |g(x) - L'| < \epsilon.$$

Also, by Theorem 269,  $g$  is locally bounded and so there exists  $B > 0$ , and  $\delta_3 > 0$  such that

$$0 < |x - a| < \delta_3 \implies |g(x)| < B.$$

Take  $\delta = \min(\delta_1, \delta_2, \delta_3)$ . Then

$$|f(x)g(x) - LL'| = |(f(x) - L)g(x) + L(g(x) - L')| \leq |f(x) - L| |g(x)| + |L| |g(x) - L'| < (B + |L|)\epsilon.$$

As the dextral side can be made arbitrarily small, the result follows.

5. For all  $\epsilon > 0$  there are  $\delta_1 > 0$ ,  $B > 0$ , and  $\delta_2 > 0$  such that

$$0 < |x - a| < \delta_1 \implies |f(x)| < \epsilon, \quad \text{and} \quad 0 < |x - a| < \delta_2 \implies |g(x)| < B.$$

Take  $\delta = \min(\delta_1, \delta_2)$ . Then

$$|f(x)g(x)| \leq |B| |f(x)| < B\epsilon.$$

As the dextral side can be made arbitrarily small, the result follows.

6. First  $|g(x)| \rightarrow |L'|$  as  $x \rightarrow a$  by part (1). Hence, for  $\epsilon = \left|\frac{L'}{2}\right| > 0$  there is a sufficiently small  $\delta' > 0$  such that

$$||g(x)| - |L'|| < \frac{|L'|}{2} \implies |L'| - \frac{|L'|}{2} < |g(x)| < |L'| + \frac{|L'|}{2} \implies \frac{|L'|}{2} < |g(x)| < \frac{3|L'|}{2},$$

that is,  $|g(x)|$  is bounded away from 0  $x$  sufficiently close to  $a$ . Now, for all  $\epsilon > 0$  there are  $\delta_1 > 0$  and  $\delta_2 > 0$  such that

$$0 < |x - a| < \delta_1 \implies |f(x) - L| < \epsilon, \quad \text{and} \quad 0 < |x - a| < \delta_2 \implies |g(x) - L'| < \epsilon.$$

For  $\delta = \min(\delta_1, \delta_2, \delta')$ ,

$$0 < |x - a| < \delta \implies L - \epsilon < f(x) < L + \epsilon, \quad \frac{|L'|}{2} < |g(x)| < \frac{3|L'|}{2}, \quad \text{and} \quad L' - \epsilon < g(x) < L' + \epsilon.$$

Hence

$$\left| \frac{f(x)}{g(x)} - \frac{L}{L'} \right| = \left| \frac{L'f(x) - Lg(x)}{g(x)L'} \right| = \left| \frac{L'(f(x) - L) - L(g(x) - L')}{g(x)L'} \right| \leq \frac{|L'| |f(x) - L| + |L| |g(x) - L'|}{|g(x)||L'|} < \frac{2(|L'| + |L|)\epsilon}{|L'||L'|},$$

which gives the result.

□

In the manner of proof of Proposition 265, we may prove the following two propositions.

**273 DEFINITION-PROPOSITION (Cauchy-Heine, Limit at  $+\infty$ )** Let  $f : ]a ; +\infty[ \rightarrow \mathbb{R}$ . The following are equivalent.

1. For each sequence  $\{x_n\}_{n=1}^{+\infty}$  of points in the interval  $]a ; +\infty[$ ,

$$x_n \rightarrow +\infty \implies f(x_n) \rightarrow L.$$

2.  $\forall \varepsilon > 0, \exists M, M > \max(0, a)$ , such that

$$x \geq M \implies |f(x) - L| < \varepsilon.$$

If either condition is fulfilled we say that  $f$  has a limit  $L$  as  $x$  tends towards  $+\infty$  and we write

$$L = \lim_{x \rightarrow +\infty} f(x).$$

**274 DEFINITION-PROPOSITION (Cauchy-Heine, Limit at  $-\infty$ )** Let  $f : ]-\infty ; a[ \rightarrow \mathbb{R}$ . The following are equivalent.

1. For each sequence  $\{x_n\}_{n=1}^{+\infty}$  of points in the interval  $] -\infty ; a[$ ,

$$x_n \rightarrow -\infty \implies f(x_n) \rightarrow L.$$

2.  $\forall \varepsilon > 0, \exists M, M < \min(0, a)$ , such that

$$x \leq M \implies |f(x) - L| < \varepsilon.$$

If either condition is fulfilled we say that  $f$  has a limit  $L$  as  $x$  tends towards  $-\infty$  and we write

$$L = \lim_{x \rightarrow -\infty} f(x).$$

**275 Definition** We write  $\lim_{x \rightarrow a^+} f(x) = +\infty$  or  $\lim_{x \searrow a} f(x) = +\infty$  if  $\forall M > 0, \exists \delta > 0$  such that

$$x \in ]a ; a + \delta[ \implies f(x) > M.$$

Similarly, we write  $\lim_{x \rightarrow a^-} f(x) = +\infty$  or  $\lim_{x \nearrow a} f(x) = +\infty$  if  $\forall M > 0, \exists \delta > 0$  such that

$$x \in ]a - \delta ; a[ \implies f(x) > M.$$

Finally, we write  $\lim_{x \rightarrow a} f(x) = +\infty$  if  $\forall M > 0, \exists \delta > 0$  such that

$$x \in ]a - \delta ; a + \delta[ \implies f(x) > M.$$

**276 Definition** We write  $\lim_{x \rightarrow a^+} f(x) = -\infty$  or  $\lim_{x \searrow a} f(x) = -\infty$  if  $\forall M < 0, \exists \delta > 0$  such that

$$x \in ]a ; a + \delta[ \implies f(x) < M.$$

Similarly, we write  $\lim_{x \rightarrow a^-} f(x) = -\infty$  or  $\lim_{x \nearrow a} f(x) = -\infty$  if  $\forall M < 0, \exists \delta > 0$  such that

$$x \in ]a - \delta ; a[ \implies f(x) < M.$$

Finally, we write  $\lim_{x \rightarrow a} f(x) = -\infty$  if  $\forall M < 0, \exists \delta > 0$  such that

$$x \in ]a - \delta ; a + \delta[ \implies f(x) < M.$$

**277 THEOREM** Let  $X, Y$  be subsets of  $\mathbb{R}$ ,  $a \in X$  and  $b \in Y$ ,  $f : X \rightarrow \mathbb{R}$ ,  $g : Y \rightarrow \mathbb{R}$  such that  $f(X) \subseteq Y$ , and let  $L \in \mathbb{R}$ . Then

$$\lim_{x \rightarrow a} f(x) = a \text{ and } \lim_{x \rightarrow b} g(x) = L \implies \lim_{x \rightarrow a} (g \circ f)(x) = L.$$

**Proof:**

□

**Homework**

**Problem 5.1.1** Prove that  $\lim_{x \rightarrow 0} \sin \frac{1}{x}$  does not exist.

**Problem 5.1.2** Let  $m, n$  be strictly positive integers. Prove that  $\lim_{x \rightarrow 1} \frac{x^n - 1}{x^m - 1} = \frac{n}{m}$ .

**Problem 5.1.3** Let  $X \subseteq \mathbb{R}$ ,  $a \in \mathbb{R}$ , and  $f, g : X \rightarrow \mathbb{R}$ . If  $f(x) \rightarrow +\infty$  and there exists a neighbourhood  $\mathcal{N}_a \subseteq X$  of  $a$  where  $f(x) \leq g(x)$ , prove that  $g(x) \rightarrow +\infty$ .

**Problem 5.1.4** Let  $X \subseteq \mathbb{R}$ ,  $a \in \mathbb{R}$ , and  $f, g : X \rightarrow \mathbb{R}$ . Suppose that

$\lim_{x \rightarrow a} f(x) = +\infty$ . Demonstrate that

1. If  $\lim_{x \rightarrow a} g(x) = +\infty$ , then  $\lim_{x \rightarrow a} (f(x) + g(x)) = +\infty$ .
2. If  $\lim_{x \rightarrow a} g(x) = L \in \mathbb{R}$ , then  $\lim_{x \rightarrow a} (f(x) + g(x)) = +\infty$ .
3. If  $\lim_{x \rightarrow a} g(x) = +\infty$ , then  $\lim_{x \rightarrow a} (f(x)g(x)) = +\infty$ .
4. If  $\lim_{x \rightarrow a} g(x) = L > 0$ , then  $\lim_{x \rightarrow a} (f(x)g(x)) = +\infty$ .

**Problem 5.1.5 (Cauchy Criterion for Functional Limits)** Let  $X \subseteq \mathbb{R}$ ,  $a \in X$ , and  $f : X \rightarrow \mathbb{R}$ . Prove that  $f$  has a limit at  $a$  (finite or infinite) if and only if for all  $\epsilon > 0$  there is a  $\delta > 0$  such that  $|x' - x''| < \delta$  implies  $|f(x') - f(x'')| < \epsilon$ .

## 5.2 Continuity

**278 Definition** A function  $f : ]a ; b[ \rightarrow \mathbb{R}$  is said to be *continuous at the point*  $x_0 \in ]a ; b[$ , if we can exchange limiting operations, as in

$$\lim_{x \rightarrow x_0} f(x) = f\left(\lim_{x \rightarrow x_0} x\right) \quad (= f(x_0)).$$

In other words, a function is continuous at the point  $x_0$  if

$$\forall \epsilon > 0, \exists \delta > 0, \text{ such that } |x - x_0| < \delta \implies |f(x) - f(x_0)| < \epsilon.$$

**279 Definition** A function  $f : ]a ; b[ \rightarrow \mathbb{R}$  is said to be *right continuous at*  $a$ , if

$$f(a) = f(a+).$$

It is said to be *left continuous at*  $b$ , if

$$f(b) = f(b-).$$

In view of the above definitions and Proposition 267, we have the following

**280 THEOREM** The following are equivalent.

1. The function  $f : ]a ; b[ \rightarrow \mathbb{R}$  is continuous at the point  $x_0 \in ]a ; b[$ .
2.  $f(x_0-) = f(x_0) = f(x_0+)$ .
3. If  $\{x_n\}_{n=1}^{+\infty}$ , and for all  $n$ ,  $x_n \in ]a ; b[$ , then  $x_n \rightarrow x_0 \implies f(x_n) \rightarrow f(x_0)$ .

**281 Example** What are the points of discontinuity of the function

$$f : \begin{matrix} ]0 ; +\infty[ & \rightarrow & \mathbb{R} \\ x & \mapsto & \begin{cases} \frac{1}{p+q} & \text{if } x \in \mathbb{Q} \cap ]0 ; +\infty[, x = \frac{p}{q}, \text{ in lowest terms} \quad ? \\ 0 & \text{if } x \in ]0 ; +\infty[ \setminus \mathbb{Q} \end{cases} \end{matrix}$$

**Solution:** Let  $a \in \mathbb{Q}$ . Since  $]0 ; +\infty[ \setminus \mathbb{Q}$  is dense in  $]0 ; +\infty[$ , there exists a sequence  $\{a_n\}_{n=1}^{+\infty}$  of points in  $]0 ; +\infty[ \setminus \mathbb{Q}$  such that  $a_n \rightarrow a$  as  $n \rightarrow +\infty$ . Observe that  $f(a_n) = 0$  but  $f(a) \neq 0$ . Hence  $a_n \rightarrow a$  does not imply  $f(a_n) \rightarrow f(a)$  and  $f$  is not continuous at  $a$ . On the other hand, let  $n \in ]0 ; +\infty[ \setminus \mathbb{Q}$ . Then  $f(b) = 0$ . Let  $\{b_n\}_{n=1}^{+\infty}$  be a sequence in  $]0 ; +\infty[ \cap \mathbb{Q}$  converging to  $b$ ,  $b_n = \frac{p_n}{q_n}$  in lowest terms. By Dirichlet's Approximation Theorem we

must have  $p_n \rightarrow +\infty$  and  $q_n \rightarrow +\infty$ . Hence  $\frac{1}{p_n + q_n} \rightarrow 0$ . So  $f$  is continuous at  $b$ . In conclusion,  $f$  is continuous at every irrational in  $[0; +\infty[$  and discontinuous at every rational in  $[0; +\infty[$ .

**282 DEFINITION-PROPOSITION (Oscillation of a function at a point)** Let  $f$  be bounded. The function  $\omega : \text{Dom}(f) \rightarrow [0; +\infty[$ , called the *oscillation of  $f$  at  $x$*  and given by

$$\omega(f, x) = \lim_{\delta \rightarrow 0^+} \sup\{|f(a) - f(b)| : |a - x| < \delta, |b - x| < \delta\}$$

is well-defined. Moreover,  $f$  is continuous at  $x$  if and only if  $\omega(f, x) = 0$ .

**Proof:** Observe that in fact

$$\omega(f, x) = \lim_{\delta \rightarrow 0^+} \sup\{|f(a) - f(b)| : |a - x| < \delta, |b - x| < \delta\} = \inf_{\delta > 0} \sup\{|f(a) - f(b)| : |a - x| < \delta, |b - x| < \delta\} \leq |f(a) - f(b)| \leq 2|f| < +\infty$$

This says that  $\omega(f, x)$  is well-defined.

□

**283 Definition** We say that a function  $f$  is continuous on the closed interval  $[a; b]$  if it is continuous everywhere on  $]a; b[$ , continuous on the right at  $a$  and continuous on the left at  $b$ . If  $X \subseteq \mathbb{R}$ , then  $f : X \rightarrow \mathbb{R}$  is said to be *continuous on  $X$*  (or *continuous*) if it is continuous at every element of  $X$ .

**284 THEOREM** Let  $X \subseteq \mathbb{R}$ . A function  $f : X \rightarrow \mathbb{R}$  is continuous if and only if the the inverse image of an open set is open in  $X$ .

**Proof:**

$\Rightarrow$  Let  $A \subseteq \mathbb{R}$  be an open set. We must shew that  $f^{-1}(A)$  is open in  $X$ . Let  $a \in f^{-1}(A)$ . Since  $f(a) \in A$  and  $A$  is open in  $\mathbb{R}$ , there exists an  $r > 0$  such that  $]f(a) - r; f(a) + r[ \subseteq A$ . Since  $f$  is continuous at  $a$ , there exists a  $\delta > 0$  such that

$$\begin{aligned} |x - a| < \delta &\implies |f(x) - f(a)| < r, \quad \text{that is, } x \in ]a + \delta; a - \delta[ \implies f(x) \in ]f(a) - r; f(a) + r[, \\ &\text{that is, } x \in ]a + \delta; a - \delta[ \implies x \in f^{-1}\left(]f(a) - r; f(a) + r[\right), \\ &\text{that is, } ]a + \delta; a - \delta[ \subseteq f^{-1}\left(]f(a) - r; f(a) + r[\right) \end{aligned}$$

Since  $f^{-1}\left(]f(a) - r; f(a) + r[\right) \subseteq f^{-1}(A)$ , we have shewn that  $]a + \delta; a - \delta[ \subseteq f^{-1}(A)$ , which means that for any  $a$ , a neighbourhood of  $a$  lies entirely in  $f^{-1}(A)$ , that is,  $f^{-1}(A)$  is open.

$\Leftarrow$  Given  $\varepsilon > 0$ , we must find a  $\delta > 0$  such that for all  $a \in X$ ,

$$|x - a| < \delta \implies |f(x) - f(a)| < \varepsilon.$$

Now

$$|f(x) - f(a)| < \varepsilon \implies f(x) \in ]f(a) - \varepsilon; f(a) + \varepsilon[ \implies x \in f^{-1}\left(]f(a) - \varepsilon; f(a) + \varepsilon[\right).$$

Now,  $]f(a) - \varepsilon; f(a) + \varepsilon[ \subseteq \mathbb{R}$  is open in  $\mathbb{R}$ , and so, by assumption, so is  $f^{-1}\left(]f(a) - \varepsilon; f(a) + \varepsilon[\right)$ . This means that if  $t \in f^{-1}\left(]f(a) - \varepsilon; f(a) + \varepsilon[\right)$  then there is a  $r > 0$  such that

$$]t - r; t + r[ \subseteq f^{-1}\left(]f(a) - \varepsilon; f(a) + \varepsilon[\right).$$

But clearly  $a \in f^{-1}\left(]f(a) - \varepsilon; f(a) + \varepsilon[\right)$ , and hence there is a  $\delta > 0$  such that

$$]a - \delta; a + \delta[ \subseteq f^{-1}\left(]f(a) - \varepsilon; f(a) + \varepsilon[\right).$$

Thus

$$x \in ]a - \delta; a + \delta[ \implies x \in f^{-1}\left(]f(a) - \varepsilon; f(a) + \varepsilon[\right),$$

or equivalently,

$$|x - a| < \delta \implies f(x) \in ]f(a) - \varepsilon; f(a) + \varepsilon[ ,$$

that is,

$$|x - a| < \delta \implies |f(x) - f(a)| < \varepsilon,$$

as we needed to shew.

□

**285 THEOREM** Let  $X \subseteq \mathbb{R}$ . A function  $f: X \rightarrow \mathbb{R}$  is continuous if and only if the the inverse image of a closed set is closed in  $X$ .

**Proof:** Let  $F \subseteq \mathbb{R}$  be a closed set. Then  $\mathbb{R} \setminus F$  is open. By Theorem 284  $f^{-1}(\mathbb{R} \setminus F)$  is open in  $X$ , and so  $X \setminus f^{-1}(\mathbb{R} \setminus F)$  is closed in  $X$ . But  $X \setminus f^{-1}(\mathbb{R} \setminus F) = f^{-1}(F)$ , proving the theorem. □

**286 THEOREM** If two continuous functions agree on a dense set of the reals, then they are identical. That is, if  $X \subseteq \mathbb{R}$  is dense in  $\mathbb{R}$  and if  $f: \mathbb{R} \rightarrow \mathbb{R}$  and  $g: \mathbb{R} \rightarrow \mathbb{R}$  satisfy  $f(x) = g(x)$  for all  $x \in X$ , then  $f(x) = g(x)$  for all  $x \in \mathbb{R}$ .

**Proof:** Let  $a \in \mathbb{R} \setminus X$ . Since  $X$  is dense in  $\mathbb{R}$ , there is a sequence  $\{x_n\}_{n=1}^{+\infty} \subseteq X$  such that  $x_n \rightarrow a$  as  $n \rightarrow +\infty$ . Notice that since  $x_n \in X$ , we have  $f(x_n) = g(x_n)$ . By continuity

$$f(a) = f\left(\lim_{n \rightarrow +\infty} x_n\right) = \lim_{n \rightarrow +\infty} f(x_n) = \lim_{n \rightarrow +\infty} g(x_n) = g\left(\lim_{n \rightarrow +\infty} x_n\right) = g(a),$$

proving the theorem. □

**287 THEOREM (Cauchy's Functional Equation)** Let  $f$  be a continuous function defined over the real numbers that satisfies the Cauchy functional equation:

$$\forall (x, y) \in \mathbb{R}^2, \quad f(x + y) = f(x) + f(y).$$

Then  $f$  is linear, that is, there is a constant  $c$  such that  $f(x) = cx$ .

**Proof:** Our method of proof is as follows. We first prove the assertion for positive integers  $n$  using induction. We then extend our result to negative integers. Thence we extend the result to reciprocals of integers and after that to rational numbers. Finally we extend the result to all real numbers by means of Theorem 286.

We prove by induction that for integer  $n \geq 0$ ,  $f(nx) = nf(x)$ . Using the functional equation,

$$f(0 \cdot x) = f(0 \cdot x + 0 \cdot x) = f(0 \cdot x) + f(0 \cdot x) \implies f(0 \cdot x) = 0f(x),$$

and the assertion follows for  $n = 0$ . Assume  $n \geq 1$  is an integer and that  $f((n-1)x) = (n-1)f(x)$ . Then

$$f(nx) = f((n-1)x + x) = f((n-1)x) + f(x) = (n-1)f(x) + f(x) = nf(x),$$

proving the assertion for all strictly positive integers.

Let  $m < 0$  be an integer. Then  $-m > 0$  is a strictly positive integer, for which the result proved in the above paragraph holds, and thus and by the above paragraph,  $f(-mx) = -mf(x)$ . Now,

$$0 = f(0) \implies 0 = f(mx + (-mx)) = f(mx) + f(-mx) \implies f(mx) = -f(-mx) = -(-mf(x)) = mf(x),$$

and the assertion follows for negative integers. We have thus proved the theorem for all integers.



Assume now that  $x = \frac{a}{b}$ , with  $a \in \mathbb{Z}$  and  $b \in \mathbb{Z} \setminus \{0\}$ . Then  $f(a) = f(a \cdot 1) = af(1)$  and  $f(a) = f\left(b \frac{a}{b}\right) = bf\left(\frac{a}{b}\right)$  by the result we proved for integers and hence

$$af(1) = bf\left(\frac{a}{b}\right) \implies f\left(\frac{a}{b}\right) = f(1)\left(\frac{a}{b}\right).$$

We have established that for all rational numbers  $x \in \mathbb{Q}$ ,  $f(x) = xf(1)$ .

We have not used the fact that the function is continuous so far. Since the rationals are dense in the reals the extension of the result now follows from Theorem 286.  $\square$

## Homework

**Problem 5.2.1** Find all functions  $f: \mathbb{R} \rightarrow \mathbb{R}$ , continuous at  $x = 0$  such that  $\forall x \in \mathbb{R}$ ,  $f(x) = f(3x)$ .

**Problem 5.2.2** Find all functions  $f: \mathbb{R} \rightarrow \mathbb{R}$ , continuous at  $x = 0$  such that  $\forall x \in \mathbb{R}$ ,  $f(x) = f\left(\frac{x}{1+x^2}\right)$ .

**Problem 5.2.3** Determine the set of points of discontinuity of the function  $f: \mathbb{R} \rightarrow \mathbb{R}$ ,  $f: x \mapsto \lfloor x \rfloor + \sqrt{x - \lfloor x \rfloor}$ .

**Problem 5.2.4** What are the points of discontinuity of the function

$$\begin{array}{ccc} \mathbb{R} & \rightarrow & \mathbb{R} \\ f: & & \\ x & \mapsto & \begin{cases} x & \text{if } x \in \mathbb{Q} \\ 0 & \text{if } x \in \mathbb{R} \setminus \mathbb{Q} \end{cases} \end{array} \quad ?$$

**Problem 5.2.5** What are the points of discontinuity of the function

$$\begin{array}{ccc} \mathbb{R} & \rightarrow & \mathbb{R} \\ f: & & \\ x & \mapsto & \begin{cases} 0 & \text{if } x \in \mathbb{Q} \\ x & \text{if } x \in \mathbb{R} \setminus \mathbb{Q} \end{cases} \end{array} \quad ?$$

**Problem 5.2.6** What are the points of discontinuity of the function

$$\begin{array}{ccc} \mathbb{R} & \rightarrow & \mathbb{R} \\ f: & & \\ x & \mapsto & \begin{cases} 0 & \text{if } x \in \mathbb{Q} \\ 1 & \text{if } x \in \mathbb{R} \setminus \mathbb{Q} \end{cases} \end{array} \quad ?$$

**Problem 5.2.7** What are the points of discontinuity of the function

$$\begin{array}{ccc} \mathbb{R} & \rightarrow & \mathbb{R} \\ f: & & \\ x & \mapsto & \begin{cases} \cos x & \text{if } x \in \mathbb{Q} \\ \sin x & \text{if } x \in \mathbb{R} \setminus \mathbb{Q} \end{cases} \end{array} \quad ?$$

**Problem 5.2.8** Find all functions  $f: \mathbb{R} \rightarrow \mathbb{R}$ , continuous at  $x = 1$  such that  $\forall x \in \mathbb{R}$ ,  $f(x) = -f(x^2)$ .

**Problem 5.2.9** Let  $a \in \mathbb{R}$  be fixed. Find all functions  $f: \mathbb{R} \rightarrow \mathbb{R}$ , continuous everywhere such that  $\forall (x, y) \in \mathbb{R}^2$ ,  $f(x - y) = f(x) - f(y) + axy$ .

**Problem 5.2.10** Let  $f: [0; +\infty[ \rightarrow [0; +\infty[$ ,  $x \mapsto \sqrt{x + \sqrt{x + \sqrt{x + \dots}}}$ . Is  $f$  right-continuous at 0?

## 5.3 Algebraic Operations with Continuous Functions

**288 THEOREM (Algebra of Continuous Functions)** Let  $f, g: ]a; b[ \rightarrow \mathbb{R}$  be continuous at the point  $x_0 \in ]a; b[$ . Then

1.  $f + g$  is continuous at  $x_0$ .
2.  $fg$  is continuous at  $x_0$ .
3. if  $g(x_0) \neq 0$ ,  $\frac{f}{g}$  is continuous at  $x_0$ .

**Proof:** This follows directly from Theorem 272.  $\square$

**289 THEOREM** Let  $X, Y$  be subsets of  $\mathbb{R}$ ,  $a \in X$  and  $b \in Y$ ,  $f: X \rightarrow \mathbb{R}$ ,  $g: Y \rightarrow \mathbb{R}$  such that  $f(X) \subseteq Y$ . If  $f$  is continuous at  $a$  and  $g$  is continuous at  $f(a)$ , then  $g \circ f$  is continuous at  $a$ .

**Proof:** This follows at once from Theorem 277.  $\square$

**290 THEOREM** Let  $f : I \rightarrow \mathbb{R}$  be a monotone function, where  $I \subseteq \mathbb{R}$  is a non-empty interval. Then the set of points of discontinuity of  $f$  is either finite or countable.

With Theorems 288 and 289 we can now demonstrate the

## 5.4 Monotonicity and Inverse Image

**291 Definition** Let  $X$  and  $Y$  be subsets of  $\mathbb{R}$ . Let  $f : X \rightarrow Y$ , and assume that  $X$  has at least two elements. Then  $f$  is said to be

- *increasing* if  $\forall (a, b) \in X^2, a < b \implies f(a) \leq f(b)$ . Equivalently, if the ratio  $\frac{f(b) - f(a)}{b - a} \geq 0$ .
- *strictly increasing* if  $\forall (a, b) \in X^2, a < b \implies f(a) < f(b)$ . Equivalently, if the ratio  $\frac{f(b) - f(a)}{b - a} > 0$ .
- *decreasing* if  $\forall (a, b) \in X^2, a < b \implies f(a) \geq f(b)$ . Equivalently, if the ratio  $\frac{f(b) - f(a)}{b - a} \leq 0$ .
- *strictly decreasing* if  $\forall (a, b) \in X^2, a < b \implies f(a) > f(b)$ . Equivalently, if the ratio  $\frac{f(b) - f(a)}{b - a} < 0$ .

$f$  is said to be *monotonic* if it is either increasing or decreasing, and *strictly monotonic* if it is either strictly increasing or strictly decreasing.



Observe that if  $f$  is increasing, then  $-f$  is decreasing, and conversely. Similarly for strictly monotonic functions.

**292 THEOREM** Let  $X \subseteq \mathbb{R}$  and let  $f : X \rightarrow \mathbb{R}$  be strictly monotone. Then  $f$  is injective.

**Proof:** Recall that  $f$  is injective if  $x \neq y \implies f(x) \neq f(y)$ . If  $f$  is strictly increasing then  $x < y \implies f(x) < f(y)$  and if  $f$  is strictly decreasing then  $x < y \implies f(x) > f(y)$ . In either case, the condition for injectivity is fulfilled.

$\square$

**293 THEOREM** Let  $I \subseteq \mathbb{R}$  be an interval and let  $f : I \rightarrow f(I)$  be strictly monotone. Then  $f^{-1}$  is strictly monotone in the same sense as  $f$ .

**Proof:** Assume first that  $f$  is strictly increasing and put  $x = f^{-1}(a)$ ,  $y = f^{-1}(b)$  and that  $a < b$ . If  $x \geq y$ , then, since  $f$  is strictly increasing,  $f(x) \geq f(y)$ . But then,  $f(f^{-1}(a)) \geq f(f^{-1}(b)) \implies a \geq b$ , a contradiction.

A similar argument finishes the theorem for  $f$  strictly decreasing.

$\square$

The following theorem is remarkable, since it does not allude to any possible continuity of the function in question.

**294 THEOREM** Let  $I \subseteq \mathbb{R}$  be an interval and let  $f : I \rightarrow f(I)$  be strictly monotone. Then  $f^{-1}$  is continuous.

**Proof:** Let  $b \in f(I)$ ,  $b = f(a)$ , and  $\varepsilon > 0$ . We must shew that there is  $\delta > 0$  such that

$$|y - b| < \delta \implies |f^{-1}(y) - a| < \varepsilon.$$

If  $a$  is not an endpoint of  $I$ , there is an  $\alpha > 0$  such that  $]a - \alpha; a + \alpha[ \subseteq I$ . Put  $\varepsilon' = \min(\varepsilon, \alpha)$ . Since both  $f$  and  $f^{-1}$  are both strictly monotone

$$|f^{-1}(y) - a| < \varepsilon' \implies b - \varepsilon' < f^{-1}(y) < b + \varepsilon' \implies f(b - \varepsilon') < f(f^{-1}(y)) < f(b + \varepsilon') \implies f(b - \varepsilon') < y < f(b + \varepsilon').$$

Since  $f$  is strictly increasing and  $a - \epsilon' < a$ ,  $f(a - \epsilon') < f(a) = b$ . Thus there must be an  $\eta > 0$  such that  $f(a - \epsilon') = b - \eta < b$ . Similarly, there is an  $\eta'$  such that  $b < b + \eta' = f(a + \epsilon')$ . Putting  $\eta'' = \min(\eta, \eta')$ , we have that for all  $y \in f(I)$ ,

$$\begin{aligned} |y - b| < \eta'' &\implies b - \eta'' < y < b + \eta'' \\ &\implies b - \eta < y < b + \eta' \\ &\implies a - \epsilon' < f^{-1}(y) < a + \epsilon' \\ &\implies |f^{-1}(y) - f^{-1}(b)| < \epsilon', \end{aligned}$$

finishing the proof for when  $a$  is not an endpoint. If  $a$  were an endpoint, the above proof carries by suppressing one of  $\eta$  or  $\eta'$ .  $\square$

**295 THEOREM** A continuous function  $f: [a; b] \rightarrow f([a; b])$  is invertible if and only if it is strictly monotone.

**Proof:**

$\implies$  Assume  $f$  is continuous and invertible. Since  $f$  is injective,  $f(a) \neq f(b)$ . Assume that  $f(a) < f(b)$ , if  $f(a) > f(b)$  the argument is similar. We would like to show that if  $a' < b' \implies f(a') < f(b')$ . Consider the continuous function  $g: [0; 1] \rightarrow \mathbb{R}$ ,

$$g(t) = f((1-t)a + ta') - f((1-t)b + tb').$$

We have

$$g(0) = f(a) - f(b) < 0 \quad \text{and} \quad g(1) = f(a') - f(b').$$

If  $g(1) = 0$ , then we must have  $a' = b'$ , contradicting  $a' < b'$ . If  $g(1) > 0$ , then by the Intermediate Value Theorem there must be an  $s \in ]0; 1[$  such that  $g(s) = 0$ . This entails

$$(1-s)a + sa' = (1-s)b + sb' \implies 0 > (1-s)(a-b) = s(b' - a') > 0,$$

absurd. This entails that  $g(1) < 0 \implies f(a') < f(b')$ , as wanted.

$\Leftarrow$  Trivially,  $f$  is surjective. If  $f$  is strictly monotone, then  $f$  is injective by Theorem 292, and thus  $f$  is invertible, by Theorem 27.

$\square$

## 5.5 Convex Functions

**296 Definition** Let  $A \times B \subseteq \mathbb{R}^2$ . A function  $f: A \rightarrow B$  is *convex* in  $A$  if  $\forall (a, b, \lambda) \in A^2 \times [0; 1]$ ,

$$f(\lambda a + (1-\lambda)b) \leq f(a)\lambda + (1-\lambda)f(b).$$

It is *strictly convex* if the inequality above is strict. Similarly, a function  $g: A \rightarrow B$  is *concave* in  $A$  if  $\forall (a, b, \lambda) \in A^2 \times [0; 1]$ ,

$$g(\lambda a + (1-\lambda)b) \geq g(a)\lambda + (1-\lambda)g(b).$$

It is *strictly concave* if the inequality above is strict.

### 5.5.1 Graphs of Functions

**297 Definition** Given a function  $f$ , its *graph* is the set on the plane

$$\Gamma_f = \{(x, y) \in \mathbb{R}^2 : y = f(x)\}.$$

**298 Example** Figures ?? through ?? show the graphs of a few standard functions, with which we presume the reader to be familiar.

## 5.6 Classical Functions

### 5.6.1 Affine Functions

**299 Definition** An *affine function* is one with assignment rule of the form  $x \mapsto ax + b$ , where  $a, b$  are real constants.

**300 THEOREM** The graph of an affine function is a line on the plane. Conversely, any non-vertical straight line on the plane is the graph on an affine function.

### 5.6.2 Quadratic Functions

### 5.6.3 Polynomial Functions

### 5.6.4 Exponential Functions

**301 DEFINITION-PROPOSITION** Let  $x \in \mathbb{R}$  be fixed. The sequence  $\left\{ \left(1 + \frac{x}{n}\right)^n \right\}_{n > -x}^{+\infty}$  is bounded and strictly increasing. Thus it converges and we define *the natural exponential function* by

$$\exp: \mathbb{R} \rightarrow \mathbb{R}, \quad \exp(x) := \lim_{n \rightarrow +\infty} \left(1 + \frac{x}{n}\right)^n.$$

**Proof:** Observe that  $1 + \frac{x}{n} > 0$  for  $n > -x$ . Using the AM-GM Inequality with  $x_1 = 1, x_2 = \dots = x_{n+1} = 1 + \frac{x}{n}$

$$\left(1 + \frac{x}{n}\right)^{n/(n+1)} < \frac{1 + n\left(1 + \frac{x}{n}\right)}{n+1} = 1 + \frac{x}{n+1} \implies \left(1 + \frac{x}{n}\right)^n < \left(1 + \frac{x}{n+1}\right)^{n+1},$$

whence the sequence is increasing.

For  $0 < x \leq 1$  then  $\left(1 + \frac{x}{n}\right)^n \leq \left(1 + \frac{1}{n}\right)^n < e$ , by Theorem 177.

If  $x > 1$  then by the already proved monotonicity,

$$\left(1 + \frac{x}{n}\right)^n \leq \left(1 + \frac{\lfloor x \rfloor + 1}{n}\right)^n < \left(1 + \frac{\lfloor x \rfloor + 1}{n(\lfloor x \rfloor + 1)}\right)^{n(\lfloor x \rfloor + 1)} < e^{\lfloor x \rfloor + 1}.$$

If  $x \leq 0$  then  $1 + \frac{x}{n} \leq 1$  and so  $\left(1 + \frac{x}{n}\right)^n \leq 1$ .  $\square$

 By Theorem 177,  $\exp(1) = e$ . We will later prove, in ????, that for all  $x \in \mathbb{R}$ ,  $\exp(x) = e^x$ .

### 5.6.5 Logarithmic Functions

### 5.6.6 Trigonometric Functions

**302 THEOREM** Let  $x \in \left]0; \frac{\pi}{2}\right[$ . Then  $\sin x < x < \tan x$ .

**Proof:**

$\square$

## Homework

**Problem 5.6.1** How many solutions does the equation

$$\sin x = \frac{x}{100}$$

have?

**Problem 5.6.2** Prove that

$$\frac{2}{\pi}x \leq \sin(x) \leq x, \forall x \in \left[0; \frac{\pi}{2}\right].$$

**Problem 5.6.3** How many solutions does the equation

$$\sin x = \log x$$

have?

**Problem 5.6.4** How many solutions does the equation

$$\sin(\sin(\sin(\sin(\sin(x)))))) = \frac{x}{3}$$

have?

**Problem 5.6.5 (Chebyshev Polynomials)**

**Problem 5.6.6 (Cardano's Formula)**

### 5.6.7 Inverse Trigonometric Functions

## 5.7 Continuity of Some Standard Functions.

### 5.7.1 Continuity Polynomial Functions

**303 LEMMA** Let  $K \in \mathbb{R}$  be a constant. The constant function  $f: \mathbb{R} \rightarrow \mathbb{R}$ ,  $f(x) = K$  is everywhere continuous.

**Proof:** Given  $a \in \mathbb{R}$  and  $\varepsilon > 0$ , take  $\delta = \varepsilon$ . Then clearly

$$|x - a| < \delta \implies |f(x) - f(a)| < \varepsilon,$$

since  $f(x) = f(a) = K$  and the quantity after the implication is  $0 < \varepsilon$  and we obtain a tautology.  $\square$

**304 LEMMA** The identity function  $f: \mathbb{R} \rightarrow \mathbb{R}$ ,  $f(x) = x$  is everywhere continuous.

**Proof:** Given  $a \in \mathbb{R}$  and  $\varepsilon > 0$ , take  $\delta = \varepsilon$ . Then clearly

$$|x - a| < \delta \implies |f(x) - f(a)| < \varepsilon,$$

since the quantity after the implication is  $|x - a| < \delta$  and we obtain a tautology.  $\square$

**305 LEMMA** Given a strictly positive integer  $n$ , the power function  $f: \mathbb{R} \rightarrow \mathbb{R}$ ,  $f(x) = x^n$  is everywhere continuous.

**Proof:** By Lemma 304, the function  $x \mapsto x$  is continuous. Applying this Lemma and the product rule from Theorem 288  $n$  times, we obtain the result.  $\square$

**306 THEOREM (Continuity of Polynomial Functions)** Let  $n$  be a fixed positive integer. Let  $a_k \in \mathbb{R}$ ,  $0 \leq k \leq n$  be constants. Then the polynomial function  $f: \mathbb{R} \rightarrow \mathbb{R}$ ,  $f(x) = a_0 + a_1x + a_2x^2 + \dots + a_nx^n$  is everywhere continuous.

**Proof:** This follows from Lemma 305 and the sum rule from Theorem 288 applied  $n + 1$  times.  $\square$

### 5.7.2 Continuity of the Exponential and Logarithmic Functions

**307 LEMMA** Let  $a > 1$ . The exponential function  $\mathbb{R} \rightarrow \mathbb{R}$ ,  $x \mapsto a^x$  is continuous at  $x = 0$ .

**Proof:** For integral  $n > 0$  we know that  $\lim_{n \rightarrow +\infty} a^{1/n} = 1$  by virtue of Theorem 174. We wish to shew that  $a^x \rightarrow 1$  as  $x \rightarrow 0$ . Observe first that  $\lim_{n \rightarrow +\infty} a^{-1/n} = \lim_{n \rightarrow +\infty} \frac{1}{a^{1/n}} = 1$  also. Thus given  $\varepsilon > 0$ , and since  $a > 1$ , there is  $N > 0$  such that

$$1 - \varepsilon < a^{-1/N} < a^{1/N} < 1 + \varepsilon.$$

If  $x \in ]-\frac{1}{N}; \frac{1}{N}[$  then,

$$a^{-1/N} < a^x < a^{1/N}.$$

By the above, this implies that

$$1 - \varepsilon < a^x < 1 + \varepsilon \implies |a^x - 1| < \varepsilon \implies |a^x - a^0| < \varepsilon,$$

finishing the proof.  $\square$

**308 THEOREM (Continuity of the Exponential Function)** Let  $a > 0$ ,  $a \neq 1$ . The exponential function  $f: \mathbb{R} \rightarrow ]0; +\infty[$ ,  $x \mapsto a^x$  is everywhere continuous.

**Proof:** Assume first that  $a > 1$ . Let us show that it is continuous at an arbitrary  $u \in \mathbb{R}$ . If  $x \rightarrow u$  then  $x - u \rightarrow 0$ . Thus

$$\lim_{x \rightarrow u} a^x = a^u \lim_{x \rightarrow u} a^{x-u} = a^u \lim_{x-u \rightarrow 0} a^{x-u} = a^u \lim_{t \rightarrow 0} a^t = a^u \cdot 1 = a^u,$$

by Lemma 307, and so the continuity is established for  $a > 1$ .

If  $0 < a < 1$  then  $\frac{1}{a} > 1$  and by what we have proved,  $x \mapsto \frac{1}{a^x}$  is continuous. Then

$$\lim_{x \rightarrow u} a^x = \lim_{x \rightarrow u} \frac{1}{\frac{1}{a^x}} = \frac{1}{\frac{1}{a^u}} = a^u,$$

proving continuity in the case  $0 < a < 1$ .  $\square$

**309 LEMMA** Let  $a > 0$ ,  $a \neq 1$ . Then  $]0; +\infty[ \rightarrow \mathbb{R}$ ,  $x \mapsto \log_a x$  is everywhere continuous.

**Proof:** Its inverse function  $\mathbb{R} \rightarrow ]0; +\infty[$ ,  $x \mapsto a^x$ , is everywhere continuous and strictly monotone. The result then follows from Theorem 294.  $\square$

### 5.7.3 Continuity of the Power Functions

**310 THEOREM** Let  $p \in \mathbb{R}$ . Then  $]0; +\infty[ \rightarrow ]0; +\infty[$ ,  $x \mapsto x^p$  is everywhere continuous.

**Proof:** This follows by the continuity of compositions:  $x^p = e^{p \log x}$ .  $\square$

### Homework

**Problem 5.7.1** Prove the continuity of the function  $\mathbb{R} \rightarrow [-1; 1]$ ,  $x \mapsto \sin x$ .

**Problem 5.7.2** Prove the continuity of the function  $[-1; 1] \rightarrow [-\frac{\pi}{2}; \frac{\pi}{2}]$ ,  $x \mapsto \arcsin x$ .

**Problem 5.7.3** Prove the continuity of the function  $\mathbb{R} \rightarrow [-1; 1]$ ,  $x \mapsto \cos x$ .

**Problem 5.7.4** Prove the continuity of the function  $[-1; 1] \rightarrow ]0; \pi]$ ,  $x \mapsto \arccos x$ .

**Problem 5.7.5** Prove the continuity of the function  $\mathbb{R} \setminus (2\mathbb{Z} + 1)\frac{\pi}{2} \rightarrow \mathbb{R}$ ,  $x \mapsto \tan x$ .

**Problem 5.7.6** Prove the continuity of the function  $\mathbb{R} \rightarrow ]-\frac{\pi}{2}; \frac{\pi}{2}[$ ,  $x \mapsto \arctan x$ .

## 5.8 Inequalities Obtained by Continuity Arguments

The technique used Theorem 287, of proving results in a dense set of the real numbers and extending the result by continuity can be exploited in a variety of situations. We now use it to give a generalisation of Bernoulli's Inequality.

**311 THEOREM (Generalisation of Bernoulli's Inequality)** Let  $(\alpha, x) \in \mathbb{R}^2$  with  $x \geq -1$ . If  $0 < \alpha < 1$  then

$$(1+x)^\alpha \leq 1 + \alpha x.$$

If  $\alpha \in ]-\infty; 0[ \cup ]1; +\infty[$  then

$$(1+x)^\alpha \geq 1 + \alpha x.$$

Equality holds in either case if and only if  $x = 0$ .

**Proof:** Let  $\alpha \in \mathbb{Q}$ ,  $0 < \alpha < 1$ . Then  $\alpha = \frac{m}{n}$  for integers  $m, n$  with  $1 \leq m < n$ . Since  $x+1 \geq 0$ , we may use the AM-GM Inequality to obtain

$$\begin{aligned} (1+x)^\alpha &= (1+x)^{m/n} \\ &= \left( (1+x)^m \cdot 1^{n-m} \right)^{1/n} \\ &\leq \frac{m(1+x) + (n-m) \cdot 1}{n} \\ &= \frac{n + mx}{n} \\ &= 1 + \frac{m}{n}x \\ &= 1 + \alpha x. \end{aligned}$$

Equality holds when the factors are the same, that is, when  $1+x = 1 \implies x = 0$ .

Assume now that  $\alpha \in \mathbb{R} \setminus \mathbb{Q}$  with  $0 < \alpha < 1$ . We can find a sequence of rational numbers  $\{a_n\}_{n=1}^{+\infty} \subseteq \mathbb{Q}$  such that  $a_n \rightarrow \alpha$  as  $n \rightarrow +\infty$ . Then

$$(1+x)^{a_n} \leq 1 + a_n x,$$

whence by the continuity of the power functions (Theorem 310),

$$(1+x)^\alpha = \lim_{n \rightarrow +\infty} (1+x)^{a_n} \leq \lim_{n \rightarrow +\infty} (1 + a_n x) = 1 + \alpha x,$$

giving the result for all real numbers  $\alpha$  with  $0 < \alpha < 1$ , except that we need to prove that equality holds only for  $x = 0$ . Take a rational number  $r$  with  $0 < \alpha < r < 1$ , and recall that we are assuming that  $\alpha$  is irrational. Then

$$(1+x)^\alpha = (1+x)^{\alpha/r} \leq \left(1 + \frac{\alpha}{r}x\right)^r.$$

Since the exponent on the right is rational, by what we have proved above  $\left(1 + \frac{\alpha}{r}x\right)^r \leq 1 + x$  with equality if and only if  $x = 0$ . Hence the full result has been proved for the case  $\alpha \in \mathbb{R}$  with  $0 < \alpha < 1$ .

Let  $\alpha > 1$ . If  $1 + \alpha x < 0$ , then obviously  $(1+x)^\alpha > 0 > 1 + \alpha x$ , and there is nothing to prove. Hence we will assume that  $\alpha x \geq -1$ . By the first part of the theorem, since  $0 < \frac{1}{\alpha} < 1$ ,

$$(1+\alpha x)^{1/\alpha} \leq 1 + \frac{1}{\alpha} \cdot \alpha x = 1 + x \implies 1 + \alpha x \leq (1+x)^\alpha,$$

with equality only if  $x = 0$ . The theorem has been proved for  $\alpha > 1$ .

Finally, let  $\alpha < 0$ . Again, if  $1 + \alpha x < 0$ , then obviously  $(1+x)^\alpha > 0 > 1 + \alpha x$ , and there is nothing to prove. Assume thus  $\alpha x \geq -1$ . Choose a strictly positive integer  $n$  satisfying  $0 < -\alpha < n$ . Now,

$$1 \geq 1 - \frac{\alpha^2}{n^2} x^2 = \left(1 - \frac{\alpha}{n} x\right) \left(1 + \frac{\alpha}{n} x\right) \Rightarrow \frac{1}{1 - \frac{\alpha}{n} x} \geq 1 + \frac{\alpha}{n} x,$$

and so by the first part of the theorem

$$\begin{aligned} (1+x)^{-\alpha/n} \leq 1 - \frac{\alpha}{n} x &\Rightarrow (1+x)^{\alpha/n} \geq \frac{1}{1 - \frac{\alpha}{n} x} \\ &\Rightarrow (1+x)^{\alpha/n} \geq 1 + \frac{\alpha}{n} x \\ &\Rightarrow (1+x)^\alpha \geq \left(1 + \frac{\alpha}{n} x\right)^n, \end{aligned}$$

and since  $n$  is a positive integer,  $\left(1 + \frac{\alpha}{n} x\right)^n \geq 1 + n \cdot \frac{\alpha}{n} x = 1 + \alpha x$  and so  $(1+x)^\alpha \geq 1 + \alpha x$  also when  $\alpha < 0$ . This finishes the proof of the theorem.  $\square$

**312 THEOREM (Monotonicity of Power Means)** Let  $a_1, a_2, \dots, a_n$  be strictly positive real numbers and let  $(\alpha, \beta) \in \mathbb{R}^2$  be such that  $\alpha \cdot \beta \neq 0$  and  $\alpha < \beta$ . Then

$$\left(\frac{a_1^\alpha + a_2^\alpha + \dots + a_n^\alpha}{n}\right)^{1/\alpha} \leq \left(\frac{a_1^\beta + a_2^\beta + \dots + a_n^\beta}{n}\right)^{1/\beta},$$

with equality if and only is  $a_1 = a_2 = \dots = a_n$ .

**Proof:** Assume first that  $0 < \alpha < \beta$ . Put  $c_\alpha = \left(\frac{a_1^\alpha + a_2^\alpha + \dots + a_n^\alpha}{n}\right)^{1/\alpha}$  and  $d_k = \left(\frac{a_k}{c_\alpha}\right)^\alpha$ . Observe that

$$\frac{c_\beta}{c_\alpha} = \left(\frac{\left(\frac{a_1}{c_\alpha}\right)^\beta + \left(\frac{a_2}{c_\alpha}\right)^\beta + \dots + \left(\frac{a_n}{c_\alpha}\right)^\beta}{n}\right)^{1/\beta} = \left(\frac{d_1^{\beta/\alpha} + d_2^{\beta/\alpha} + \dots + d_n^{\beta/\alpha}}{n}\right)^{1/\beta},$$

and that

$$\left(\frac{d_1 + d_2 + \dots + d_n}{n}\right)^{1/\alpha} = \frac{1}{c_\alpha} \left(\frac{a_1^\alpha + a_2^\alpha + \dots + a_n^\alpha}{n}\right)^{1/\alpha} = 1 \Rightarrow d_1 + d_2 + \dots + d_n = n.$$

Put  $d_k = 1 + x_k$ . Then  $x_1 + x_2 + \dots + x_n = 0$ . By Theorem 311,

$$d_k^{\beta/\alpha} = (1 + x_k)^{\beta/\alpha} \geq 1 + \frac{\beta}{\alpha} x_k. \quad (5.1)$$

Letting  $k$  run from 1 through  $n$  and adding,

$$d_1^{\beta/\alpha} + d_2^{\beta/\alpha} + \dots + d_n^{\beta/\alpha} \geq n + \frac{\beta}{\alpha} (x_1 + x_2 + \dots + x_n) = n.$$

Hence

$$\frac{d_1^{\beta/\alpha} + d_2^{\beta/\alpha} + \dots + d_n^{\beta/\alpha}}{n} \geq 1 \Rightarrow \frac{c_\beta}{c_\alpha} \geq 1,$$

proving the theorem when  $0 < \alpha < \beta$ .

If  $\alpha < \beta < 0$ , then  $0 < \frac{\beta}{\alpha} < 1$ . The inequality in (5.1) is reversed, giving  $\frac{d_1^{\beta/\alpha} + d_2^{\beta/\alpha} + \dots + d_n^{\beta/\alpha}}{n} \leq 1$ , and since  $\beta < 0$ ,

$$\frac{c_\beta}{c_\alpha} = \left(\frac{d_1^{\beta/\alpha} + d_2^{\beta/\alpha} + \dots + d_n^{\beta/\alpha}}{n}\right)^{1/\beta} \geq 1^{1/\beta} = 1,$$



proving the theorem when  $\alpha < \beta < 0$ .

Finally, we tackle the case  $\alpha < 0 < \beta$ . By the AM-GM Inequality, putting  $G = (a_1 a_2 \cdots a_n)^{1/n}$

$$G^\alpha = (a_1^\alpha a_2^\alpha \cdots a_n^\alpha)^{1/n} \leq \frac{a_1^\alpha + a_2^\alpha + \cdots + a_n^\alpha}{n}.$$

Raising the quantities at the extreme of the inequalities to the power  $-1/\alpha$  and remembering that  $-1/\alpha > 0$ , we gather that

$$\left( \frac{a_1^\alpha + a_2^\alpha + \cdots + a_n^\alpha}{n} \right)^{1/\alpha} \leq G.$$

In a similar manner,

$$G^\beta = (a_1^\beta a_2^\beta \cdots a_n^\beta)^{1/n} \leq \frac{a_1^\beta + a_2^\beta + \cdots + a_n^\beta}{n},$$

and

$$G \leq \left( \frac{a_1^\beta + a_2^\beta + \cdots + a_n^\beta}{n} \right)^{1/\beta},$$

since  $\beta > 0$ . This finishes the proof.  $\square$

**313 LEMMA** Let  $\alpha, a, x$  be real numbers with  $\alpha > 1$ ,  $a > 0$ , and  $x \geq 0$ . Then

$$x^\alpha - ax \geq (1 - \alpha) \left( \frac{a}{\alpha} \right)^{\alpha/(\alpha-1)}.$$

**Proof:** By Theorem 311, since  $\alpha > 1$ ,

$$(1 + z)^\alpha \geq 1 + \alpha z, \quad z \geq -1,$$

with equality only if  $z = 0$ . Putting  $z = 1 + y$ ,

$$y^\alpha \geq 1 + \alpha(y - 1) \implies y^\alpha - \alpha y \geq 1 - \alpha, \quad y \geq 0,$$

with equality only if  $y = 1$ . Let  $c > 0$  be a constant. Multiplying the above inequality by  $c^\alpha$  we obtain

$$(cy)^\alpha - \alpha c^{\alpha-1}(cy) \geq (1 - \alpha)c^\alpha, \quad \text{for } y \geq 0.$$

Putting  $x = cy$  and  $a = \alpha c^{\alpha-1}$ , we get

$$x^\alpha - ax \geq (1 - \alpha) \left( \frac{a}{\alpha} \right)^{\alpha/(\alpha-1)},$$

with equality if and only if  $x = c = \left( \frac{a}{\alpha} \right)^{\alpha/(\alpha-1)}$ .

$\square$

**314 THEOREM (Young's Inequality)** Let  $p > 1$  and put  $\frac{1}{p} + \frac{1}{q} = 1$ . Then for  $(x, y) \in ([0; +\infty])^2$  we have

$$xy \leq \frac{x^p}{p} + \frac{y^q}{q}.$$

**Proof:** Put  $\alpha = p$ ,  $a = py$  in Lemma 313, obtaining

$$x^p - (py)x \geq (1 - p) \left( \frac{py}{p} \right)^{p/(p-1)} = (1 - p)y^{p/(p-1)}.$$

Now,  $\frac{1}{q} = \frac{p-1}{p} \implies q = \frac{p}{p-1}$  and  $p-1 = \frac{p}{q}$ . Hence

$$x^p - (py)x \geq (1 - p)y^{p/(p-1)} \implies (1 - p)y^{p/(p-1)} \geq -\frac{p}{q}y^q,$$

and rearranging gives the result sought.  $\square$

We now derive a generalisation of the Cauchy-Bunyakovsky-Schwarz Inequality.

**315 THEOREM (Hölder Inequality)** Let  $x_j, y_k, 1 \leq j, k \leq n$ , be real numbers. Let  $p > 1$  and put  $\frac{1}{p} + \frac{1}{q} = 1$ . Then

$$\left| \sum_{k=1}^n x_k y_k \right| \leq \left( \sum_{k=1}^n |x_k|^p \right)^{1/p} \left( \sum_{k=1}^n |y_k|^q \right)^{1/q}.$$

**Proof:** If either  $\sum_{k=1}^n |x_k|^p = 0$  or  $\sum_{k=1}^n |y_k|^q = 0$  there is nothing to prove, so assume otherwise. From Young's Inequality we have

$$\frac{|x_k|}{\left( \sum_{k=1}^n |x_k|^p \right)^{1/p}} \frac{|y_k|}{\left( \sum_{k=1}^n |y_k|^q \right)^{1/q}} \leq \frac{|x_k|^p}{\left( \sum_{k=1}^n |x_k|^p \right)^p} + \frac{|y_k|^q}{\left( \sum_{k=1}^n |y_k|^q \right)^q}.$$

Adding, we deduce

$$\begin{aligned} \sum_{k=1}^n \frac{|x_k|}{\left( \sum_{k=1}^n |x_k|^p \right)^{1/p}} \frac{|y_k|}{\left( \sum_{k=1}^n |y_k|^q \right)^{1/q}} &\leq \frac{1}{\left( \sum_{k=1}^n |x_k|^p \right)^p} \sum_{k=1}^n |x_k|^p + \frac{1}{\left( \sum_{k=1}^n |y_k|^q \right)^q} \sum_{k=1}^n |y_k|^q \\ &= \frac{\sum_{k=1}^n |x_k|^p}{\left( \sum_{k=1}^n |x_k|^p \right)^p} + \frac{\left( \sum_{k=1}^n |y_k|^q \right)^q}{\left( \sum_{k=1}^n |y_k|^q \right)^q} \\ &= \frac{1}{p} + \frac{1}{q} \\ &= 1. \end{aligned}$$

This gives

$$\sum_{k=1}^n |x_k y_k| \leq \left( \sum_{k=1}^n |x_k|^p \right)^{1/p} \left( \sum_{k=1}^n |y_k|^q \right)^{1/q}.$$

The result follows by observing that

$$\left| \sum_{k=1}^n x_k y_k \right| \leq \sum_{k=1}^n |x_k y_k| \leq \left( \sum_{k=1}^n |x_k|^p \right)^{1/p} \left( \sum_{k=1}^n |y_k|^q \right)^{1/q}.$$

□

Finally, we derive a generalisation of Minkowski's Inequality.

**316 THEOREM (Generalised Minkowski Inequality)** Let  $p \in ]1; +\infty[$ . Let  $x_j, y_k, 1 \leq j, k \leq n$ , be real numbers. Then the following inequality holds

$$\left( \sum_{k=1}^n |x_k + y_k|^p \right)^{1/p} \leq \left( \sum_{k=1}^n |x_k|^p \right)^{1/p} + \left( \sum_{k=1}^n |y_k|^p \right)^{1/p}.$$

**Proof:** From the triangle inequality for real numbers

$$|x_k + y_k|^p = |x_k + y_k| |x_k + y_k|^{p-1} \leq (|x_k| + |y_k|) |x_k + y_k|^{p-1}.$$

Adding

$$\sum_{k=1}^n |x_k + y_k|^p \leq \sum_{k=1}^n |x_k| |x_k + y_k|^{p-1} + \sum_{k=1}^n |y_k| |x_k + y_k|^{p-1}. \quad (5.2)$$

By the Hölder Inequality

$$\begin{aligned} \sum_{k=1}^n |x_k| |x_k + y_k|^{p-1} &\leq \left( \sum_{k=1}^n |x_k|^p \right)^{1/p} \left( \sum_{k=1}^n |x_k + y_k|^{(p-1)q} \right)^{1/q} \\ &= \left( \sum_{k=1}^n |x_k|^p \right)^{1/p} \left( \sum_{k=1}^n |x_k + y_k|^p \right)^{1/q} \end{aligned} \quad (5.3)$$

In the same manner we deduce

$$\sum_{k=1}^n |y_k| |x_k + y_k|^{p-1} \leq \left( \sum_{k=1}^n |y_k|^p \right)^{1/p} \left( \sum_{k=1}^n |x_k + y_k|^p \right)^{1/q}. \quad (5.4)$$

Hence (5.2) gives

$$\begin{aligned} \sum_{k=1}^n |x_k + y_k|^p &\leq \left( \sum_{k=1}^n |x_k|^p \right)^{1/p} \left( \sum_{k=1}^n |x_k + y_k|^p \right)^{1/q} + \left( \sum_{k=1}^n |y_k|^p \right)^{1/p} \left( \sum_{k=1}^n |x_k + y_k|^p \right)^{1/q} \\ &= \left( \left( \sum_{k=1}^n |x_k|^p \right)^{1/p} + \left( \sum_{k=1}^n |y_k|^p \right)^{1/p} \right) \left( \sum_{k=1}^n |x_k + y_k|^p \right)^{1/q} \end{aligned}$$

from where we deduce the result.  $\square$

## Homework

**Problem 5.8.1** Prove that if  $\alpha > 0$  and  $n > 0$  an integer then

$$\frac{n^{1+\alpha} - (n-1)^{1+\alpha}}{1+\alpha} < n^\alpha < \frac{(n+1)^{1+\alpha} - n^{1+\alpha}}{1+\alpha}.$$

Deduce that

$$\lim_{n \rightarrow +\infty} \frac{1^\alpha + 2^\alpha + \dots + n^\alpha}{n^{1+\alpha}} = \frac{1}{1+\alpha}.$$

## 5.9 Intermediate Value Property

**317 THEOREM (Intermediate Value Theorem)** Let  $I \subseteq \mathbb{R}$  and let  $(a, b) \in I^2$ . Let  $f : I \rightarrow \mathbb{R}$  be a continuous function such that  $f(a) \leq f(b)$ . Then  $f$  attains every intermediate value between  $f(a)$  and  $f(b)$ , that is,

$$\forall t \in [f(a); f(b)], \exists c \in I, \quad \text{such that} \quad f(c) = t.$$

**Proof:** Suppose on the contrary that there is a  $t \in [f(a); f(b)]$  such that for all  $c \in I$ ,  $f(c) \neq t$ . Hence  $f(a) < t < f(b)$ . Assume, without loss of generality, that  $a < b$ . Consider the sets

$$U = ]-\infty; a[ \cup \{x \in [a; b] : f(x) < t\} = ]-\infty; a[ \cup f^{-1}(] -\infty; t[ \cap ] a; b]),$$

and

$$V = ]b; +\infty[ \cup \{x \in [a; b] : f(x) > t\} = ]b; +\infty[ \cup f^{-1}(] t; +\infty[ \cap ] a; b]).$$

Then  $U, V$  are open sets of  $\mathbb{R}$  by virtue of Theorem 284. But then  $\mathbb{R} = U \cup V$  and  $U \cap V = \emptyset$ ,  $U \neq \emptyset$ ,  $V \neq \emptyset$ , contradicting the fact that  $\mathbb{R}$  is connected. Thus there must exist a  $c$  such that  $f(c) = t$ .  $\square$

**318 COROLLARY** A continuous function defined on an interval maps that interval into an interval.

**Proof:** This follows at once from the Intermediate Value Theorem and the definition of an interval.  $\square$

**319 THEOREM (Bolzano's Theorem)** If  $f : [u; v] \rightarrow \mathbb{R}$  is continuous and  $f(u)f(v) < 0$ , then there is a  $w \in ]u; v[$  such that  $f(w) = 0$ .

**Proof:** This follows at once from the Intermediate Value Theorem by putting  $a = \min(f(u), f(v)) < 0$  and  $b = \max(f(u), f(v)) > 0$ .  $\square$

**320 COROLLARY** Every polynomial  $p(x) \in \mathbb{R}[x]$  with real coefficients and odd degree has at least one real root.

**Proof:** Let  $p(x) = a_0 + a_1x + a_2x^2 + \cdots + a_nx^n$ , with  $a_n \neq 0$  and  $n$  odd. Since  $p$  has odd degree,  $\lim_{x \rightarrow -\infty} p(x) = (-\infty) \text{signum}(a_n)$  and  $\lim_{x \rightarrow +\infty} p(x) = (+\infty) \text{signum}(a_n)$ , which are of opposite sign. The polynomial must then attain positive and negative values and between values of opposite sign, it will have a real root.  $\square$

**321 COROLLARY** If  $f$  is continuous at the point  $a$  and  $f(a) \neq 0$ , then there is a neighbourhood of  $a$  where  $f(x)$  has the same sign as  $f(a)$ .

**Proof:** Take  $\varepsilon = \frac{|f(a)|}{2} > 0$  in the definition of continuity. There is a  $\delta > 0$  such that

$$|x - a| < \delta \implies |f(x) - f(a)| < \frac{|f(a)|}{2} \implies f(a) - \frac{|f(a)|}{2} < f(x) < f(a) + \frac{|f(a)|}{2},$$

from where the result follows.  $\square$

**322 THEOREM** A continuous function defined on a compact set maps that compact set into a compact set.

**Proof:** Let  $f : X \rightarrow \mathbb{R}$  be continuous and  $X \subseteq \mathbb{R}$  compact. Let  $\{y_n\}_{n=1}^{+\infty} \subseteq f(X)$  be an infinite sequence of  $f(X)$ . There are  $x_n \in X$  such that  $x_n = f(y_n)$ . Since  $\{x_n\}_{n=1}^{+\infty} \subseteq X$  is an infinite sequence of  $X$  and  $X$  is compact, it has a convergent subsequence in  $X$ , say,  $\{x_{n_k}\}_{k=1}^{+\infty}$  with  $x_{n_k} \rightarrow x \in X$ , by virtue of Theorem 143. Since  $f$  is continuous

$$x_{n_k} \rightarrow x \implies f(x_{n_k}) \rightarrow f(x).$$

Clearly  $f(x) \in f(X)$ . Thus the arbitrary sequence  $\{y_n\}_{n=1}^{+\infty} \subseteq f(X)$  has the convergent subsequence  $\{y_{n_k}\}_{k=1}^{+\infty}$  in  $f(X)$ , and one more appeal to Theorem 143 proves compactness.  $\square$

**323 THEOREM (Weierstrass Theorem)** A continuous function  $f : [a; b] \rightarrow \mathbb{R}$  attains a maximum and a minimum on  $[a; b]$ .

**Proof:** By Theorem 322,  $f([a; b])$  is compact, and so, by the Heine-Borel Theorem, it is closed and bounded. Thus there exists  $(m, M) \in \mathbb{R}^2$  such that  $m = \inf_{x \in [a; b]} f(x)$  and  $M = \sup_{x \in [a; b]} f(x)$ . We must prove that these are attained in  $[a; b]$ , i.e., that there exist  $\mu \in [a; b]$  and  $\mu' \in [a; b]$  such that  $f(\mu) = m$  and  $f(\mu') = M$ . By the Approximation Property of the Infimum and the Supremum, we may find sequences  $\{m_n\}_{n=1}^{+\infty} \subseteq [a; b]$ , and  $\{M_n\}_{n=1}^{+\infty} \subseteq [a; b]$  such that  $m \leq m_n$  and  $m_n \rightarrow m$ , and also,  $M_n \leq M$ , and  $M_n \rightarrow M$  as  $n \rightarrow +\infty$ . By the Intermediate Value Theorem, there exist  $\mu_n \in [a; b]$  and  $\mu'_n \in [a; b]$  such that  $m_n = f(\mu_n)$  and  $M_n = f(\mu'_n)$ . By the compactness of  $[a; b]$  the sequences  $\{\mu_n\}_{n=1}^{+\infty} \subseteq [a; b]$  and  $\{\mu'_n\}_{n=1}^{+\infty} \subseteq [a; b]$  have convergent subsequences  $\{\mu_{n_k}\}_{k=1}^{+\infty} \subseteq [a; b]$  and  $\{\mu'_{n_k}\}_{k=1}^{+\infty} \subseteq [a; b]$  such that  $\mu_{n_k} \rightarrow \mu \in [a; b]$  and  $\mu'_{n_k} \rightarrow \mu' \in [a; b]$ . By continuity and the uniqueness of limits,

$$\mu_{n_k} \rightarrow \mu \implies m_{n_k} = f(\mu_{n_k}) \rightarrow m = f(\mu), \quad \text{and} \quad \mu'_{n_k} \rightarrow \mu' \implies M_{n_k} = f(\mu'_{n_k}) \rightarrow M = f(\mu'),$$

and so  $f$  attains both extrema in  $[a; b]$ .  $\square$

**324 THEOREM (Fixed Point Theorem)** Let  $f : [a; b] \rightarrow [a; b]$  be continuous. Then  $f$  has a fixed point, that is, there is  $c \in [a; b]$  such that  $f(c) = c$ .

**Proof:** If either  $f(a) = a$  or  $f(b) = b$  we are done. Assume then that  $f(a) > a$  and  $f(b) < b$ . Put  $g(x) = f(x) - x$ . Then  $g$  is continuous,  $g(a) > 0$  and  $g(b) < 0$ . By Bolzano's Theorem, there must be a  $c \in ]a; b[$  such that  $g(c) = 0$ , that is,  $f(c) - c = 0$ , finishing the proof.  $\square$

## Homework

**Problem 5.9.1** Let  $p(x), q(x)$  be polynomials with real coefficients such that

$$p(x^2 + x + 1) = p(x)q(x).$$

Prove that  $p$  must have even degree.

**Problem 5.9.2** A function  $f$  defined over all real numbers is continuous and for all real  $x$  satisfies

$$(f(x)) \cdot ((f \circ f)(x)) = 1.$$

Given that  $f(1000) = 999$ , find  $f(500)$ .

**Problem 5.9.3** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a continuous function such that  $\lim_{x \rightarrow -\infty} f(x) = 0 = \lim_{x \rightarrow +\infty} f(x)$ . If  $f$  is strictly negative somewhere on  $\mathbb{R}$  then  $f$  attains a finite absolute minimum on  $\mathbb{R}$ . If  $f$  is strictly positive somewhere on  $\mathbb{R}$  then  $f$  attains a finite absolute maximum on  $\mathbb{R}$ .

**Problem 5.9.4** Let  $f : [0; 1] \rightarrow [0; 1]$  be continuous. Prove that there is no  $c \in [0; 1]$  such that  $f^{-1}(\{c\})$  has exactly two elements.

**Problem 5.9.5** Let  $f, g$  be continuous functions from  $[0; 1]$  to  $[0; 1]$  such that

$$\forall x \in [0; 1] \quad f(g(x)) = g(f(x)).$$

Prove that  $f$  and  $g$  have a common fixed point in  $[0; 1]$ .

**Problem 5.9.6** A continuous function  $f : \mathbb{R} \rightarrow \mathbb{R}$  satisfies

$$\forall x \in \mathbb{R} \quad f(x + f(x)) = f(x).$$

Prove that  $f$  is constant.

**Problem 5.9.7** Let  $I$  be a closed and bounded interval on the line and let  $f$  be continuous on  $I$ . Suppose that for each  $x \in I$ , there exists a  $y \in I$  such that

$$|f(y)| \leq \frac{1}{2}|f(x)|.$$

Prove the existence of a  $t \in I$  such that  $f(t) = 0$ .

**Problem 5.9.8** Find all continuous functions that satisfy the functional equation

$$f(x) + f(y) = f\left(\frac{x+y}{1-xy}\right),$$

for all  $-1 < x, y < 1$ .

**Problem 5.9.9 (Putnam 1947)** A real valued continuous function satisfies for all real  $x, y$  the functional equation

$$f(\sqrt{x^2 + y^2}) = f(x)f(y).$$

Prove that  $f(x) = (f(x))^{x^2}$ .

**Problem 5.9.10** Suppose that  $f : [0; 1] \rightarrow [0; 1]$  is continuous. Prove that there is a number  $c$  in  $[0; 1]$  such that  $f(c) = 1 - c$ .

**Problem 5.9.11 (Universal Chord Theorem)** Suppose that  $f$  is a continuous function of  $[0; 1]$  and that  $f(0) = f(1)$ . Let  $n$  be a strictly positive integer. Prove that there is some number  $x \in [0; 1]$  such that  $f(x) = f(x + 1/n)$ .

**Problem 5.9.12** Under the same conditions of problems 5.9.11 prove that there are no universal chords of length  $a, 0 < a < 1, a \neq 1/n$ .

## 5.10 Variation of a Function and Uniform Continuity


**325 Definition** A partition  $\mathcal{P}$  of the interval  $[a; b]$  is any finite set of points  $x_0, x_1, \dots, x_n$  such that

$$a = x_0 < x_1 < \dots < x_n = b.$$

A partition  $\mathcal{P}'$  of  $[a; b]$  is said to be *finer* than the partition  $\mathcal{P}$  if  $\mathcal{P} \subseteq \mathcal{P}'$ .

**326 Definition** The *mesh* or *norm* of  $\mathcal{P}$  is

$$\|\mathcal{P}\| = \max_{1 \leq k \leq n} |x_k - x_{k-1}|.$$

 If  $\mathcal{P} \subseteq \mathcal{P}'$  then clearly  $\|\mathcal{P}'\| \leq \|\mathcal{P}\|$ , since the finer partition has probably more points which will make the corresponding subintervals smaller.

**327 Definition** Let  $f$  be a bounded function on an interval  $[a; b]$  and let  $I \subseteq [a; b]$  be a subinterval. The *oscillation of  $f$  on  $I$*  is defined and denoted by

$$\omega(f, I) = \sup_{x \in I} f(x) - \inf_{x \in I} f(x).$$

**328 THEOREM** Let  $f: [a; b] \rightarrow \mathbb{R}$  be a continuous function. Given  $\varepsilon > 0$  there exists a partition of  $[a; b]$  into a finite number of subintervals of equal length such that the oscillation of  $f$  on each of these subintervals is at most  $\varepsilon$ .

**Proof:** Let  $P_\varepsilon$  mean the following: there is an  $\varepsilon > 0$  such that for all partitions of  $[a; b]$  into a finite number of intervals of equal length, the oscillation of  $f$  is  $\geq \varepsilon$ . By bisecting  $[a; b]$ , at least one of the halves must have property  $P_\varepsilon$ , say  $[a_1; b_1]$ . If  $[a; b]$  we to have property  $P_\varepsilon$ , then by bisecting  $[a_1; b_1]$ , at least one of the halves must have property  $P_\varepsilon$ , say  $[a_2; b_2]$ . Continuing in this way we have constructed a sequence of imbricated intervals

$$[a; b] \supseteq [a_1; b_1] \supseteq [a_2; b_2] \supseteq \cdots \supseteq [a_n; b_n] \supseteq \cdots$$

where the length of  $[a_n; b_n]$  is  $b_n - a_n = \frac{b-a}{2^n} \rightarrow 0$  as  $n \rightarrow +\infty$ . By the Cantor Intersection Theorem, there is a point  $c \in \bigcap_{n=1}^{\infty} [a_n; b_n]$ . Moreover, we have  $\omega(f, [a_n; b_n]) \geq \varepsilon$ . Since  $c \in [a; b]$ ,  $f$  is continuous at  $c$ . Hence there is a  $\delta > 0$  such that

$$x \in ]c - \delta; c + \delta[ \implies |f(x) - f(c)| < \frac{\varepsilon}{2}$$

. Taking  $(x', x'') \in ]c - \delta; c + \delta[$  we have

$$|f(x') - f(x'')| \leq |f(x') - f(c)| + |f(c) - f(x'')| < \varepsilon,$$

whence

$$\omega(f, [a; b] \cap ]c - \delta; c + \delta[) < \varepsilon.$$

Now, if there was an  $\varepsilon > 0$  such that for all partitions of  $[a; b]$  into a finite number of intervals of equal length, the oscillation of  $f$  is  $\geq \varepsilon$ , then by taking  $n$  large enough above we could find one of the  $[a_n; b_n]$  completely inside one of the subintervals of the partition. By the above, the oscillation there would be  $< \varepsilon$ , a contradiction.  $\square$

**329 THEOREM** Let  $f: [a; b] \rightarrow \mathbb{R}$  be a continuous function. Given  $\varepsilon > 0$  there exists a  $\delta > 0$  such that on any subinterval  $I \subseteq [a; b]$  having length  $< \delta$  the oscillation of  $f$  on  $I$  is  $< \varepsilon$ .

**Proof:** Let  $\delta = \frac{b-a}{n}$ . By Theorem 328 we may choose  $n$  so large that the oscillation of  $f$  on each of

$$[a; a + \delta], [a + \delta; a + 2\delta], \dots, [a + (n-1)\delta; b], \quad (5.5)$$

is  $< \frac{\varepsilon}{2}$ . Let  $I \subseteq [a; b]$  be any subinterval of length  $< \delta$  and let  $x' \in I$  be the point where  $f$  achieves its largest value and  $x'' \in I$  be the point where  $f$  achieves its smallest value. Then  $x'$  and  $x''$  either belong to the same interval in 5.5—in which case  $|f(x') - f(x'')| < \frac{\varepsilon}{2}$ —or since  $I$  has length smaller than  $\delta$ , to two consecutive subintervals

$$[a + (j-1)\delta; a + j\delta], [a + j\delta; a + (j+1)\delta].$$

In this case

$$f(x') - f(x'') = (f(x') - f(a + j\delta)) + (f(a + j\delta) - f(x'')) < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon.$$

The theorem now follows.  $\square$

**330 Definition** A function  $f$  is said to be *uniformly continuous* on  $[a; b]$  if  $\forall \varepsilon > 0$  there exists  $\delta > 0$  depending only on  $\varepsilon$  such that for any  $(u, v) \in [a; b]^2$ ,

$$|u - v| < \delta \implies |f(u) - f(v)| < \varepsilon.$$

**331 THEOREM** If  $f: [a; b] \rightarrow \mathbb{R}$  is continuous, then  $f$  is uniformly continuous.

**Proof:** This follows from Theorem 329.  $\square$

**332 THEOREM (Heine's Theorem)** If  $f: X \rightarrow \mathbb{R}$  is continuous and  $X$  is compact, then  $f$  is uniformly continuous.

**Proof:** This follows from Theorem 331.  $\square$

**333 THEOREM** Let  $f$  be an increasing function on an open interval  $]a; b[$ . Then, for any  $x$  satisfying  $a < x < b$ ,

$$\sup_{t \in ]a; x[} f(t) = f(x-) \leq f(x) \leq \inf_{t \in ]x; b[} f(t) = f(x+).$$

Moreover, if  $a < x < y < b$ , then  $f(x+) \leq f(y-)$ .

**Proof:** The set  $\{f(t) : a < u < x\}$  is bounded above by  $f(x)$  and hence it has a supremum  $\sup_{t \in ]a; x[} f(t) = A$  and

clearly  $A \leq f(x)$  as  $f$  is increasing. Let us show that  $A = f(x-)$ . By the Approximation Property of the Supremum, there is  $\delta > 0$  such that  $a < x - \delta < x$  and  $A - \varepsilon < f(x - \delta) \leq A$ . But as  $f$  is increasing,

$$x - \delta < t < x \implies f(x - \delta) \leq f(t) < A \implies |f(x) - A|,$$

whence  $f(x-) = A$ .

A similar reasoning gives  $\inf_{t \in ]x; b[} f(t) = f(x+)$ .

Now, if  $a < x < y < b$ , then by what has already been proved we obtain

$$f(x+) = \inf_{x < t < b} f(t) = \inf_{x < t < y} f(t),$$

again, remembering that  $f$  is increasing. Similarly,

$$f(y-) = \sup_{a < t < y} f(t) = \inf_{x < t < y} f(t),$$

from where  $f(x+) \leq f(y-)$ .  $\square$

**334 THEOREM** Let  $f$  be an increasing function defined on the interval  $[a; b]$  and let

$$a = x_0 < x_1 < x_2 < \dots < x_n = b$$

be  $n + 1$  points partitioning the interval. Then

$$\sum_{k=1}^{n-1} (f(x_{k+}) - f(x_{k-})) \leq f(b) - f(a).$$

**Proof:** Let  $y_k \in ]x_k; x_{k+1}[$ . For  $1 \leq k \leq n-1$ , by Theorem ??,

$$f(x_{k+}) \leq f(y_k) \quad \text{and} \quad f(y_{k-1}) \leq f(x_k-) \implies f(x_{k+}) - f(x_k-) \leq f(y_k) - f(y_{k-1}).$$

Adding,

$$\sum_{k=1}^{n-1} (f(x_{k+}) - f(x_k-)) \leq \sum_{k=1}^{n-1} (f(y_k) - f(y_{k-1})) = f(y_{n-1}) - f(y_0).$$

The proof is completed upon noticing that  $f(y_{n-1}) - f(y_0) \leq f(b) - f(a)$ .  $\square$

**335 THEOREM** Let  $f : [a; b] \rightarrow \mathbb{R}$  be a monotone function, Then the set of points of discontinuity of  $f$  is either finite or countable.

**Proof:** Assume  $f$  is increasing, for if  $f$  were decreasing, we may apply the same argument to  $-f$ . Let  $m > 0$  be an integer, and let

$$\mathcal{S}_m = \left\{ x \in [a; b] : f(x_+) - f(x_-) \geq \frac{1}{m} \right\}.$$

If  $x_1 < x_2 < \dots < x_n$  are in  $\mathcal{S}_m$  then by Theorem 334,

$$\frac{n}{m} \leq f(b) - f(a),$$

which implies that  $\mathcal{S}_m$  is a finite set. The set of discontinuities of  $f$  in  $[a; b]$  is  $\bigcup_{m=1}^{\infty} \mathcal{S}_m$ , the countable union of finite sets, and hence it is countable.  $\square$

**336 Definition** Let  $f$  be a function defined on the interval  $[a; b]$  and let

$$a = x_0 < x_1 < x_2 < \dots < x_n = b$$

be  $n+1$  points partitioning the interval. If there exists  $V > 0$  such that

$$\sum_{k=1}^n |f(x_k) - f(x_{k-1})| \leq V$$

for all partitions of  $[a; b]$ , then we say that  $f$  is of bounded variation on  $[a; b]$ .

**337 THEOREM** If  $f$  is monotonic on  $[a; b]$ , then  $f$  is bounded variation on  $[a; b]$ .

**Proof:** Let

$$a = x_0 < x_1 < x_2 < \dots < x_n = b$$

be any partition of  $[a; b]$ . Then

$$\sum_{k=1}^n |f(x_k) - f(x_{k-1})| = \max(f(b) - f(a), f(a) - f(b)),$$

the first choice occurring when  $f$  is increasing and the second when  $f$  is decreasing. Then  $V = |f(b) - f(a)|$  satisfies the definition of bounded variation for an arbitrary partition.  $\square$

**338 THEOREM** If  $f$  is of bounded variation on  $[a; b]$  then  $f$  is bounded on  $[a; b]$ .

**Proof:** Let  $x \in ]a; b[$  and consider the partition  $a < x < b$  of  $[a; b]$ . Since  $f$  is of bounded variation there is a  $V > 0$  such that

$$|f(a) - f(x)| + |f(x) - f(b)| \leq V.$$

But then

$$|f(x)| \leq |f(x) - f(a)| + |f(a)| \leq V + |f(a)|.$$

and so  $f$  is bounded by the constant quantity  $V + |f(a)|$ .  $\square$



## Homework

**Problem 5.10.1** Shew that  $]0; +\infty[ \rightarrow ]0; +\infty[$ ,  $x \mapsto x^2$ , is not uniformly continuous.

## 5.11 Classical Limits

**339 LEMMA** If  $0 < x \leq 1$  then

$$1 \leq \frac{\exp(x) - 1}{x} \leq 1 + x(e - 2).$$

If  $-\frac{1}{2} \leq x < 0$  then

$$1 + x \leq \frac{\exp(x) - 1}{x} \leq 1 + \frac{x}{4}.$$

**Proof:**

Since  $\left(1 + \frac{x}{n}\right)^n \leq \exp(x)$  for  $n > -x$  by Proposition 301, we have  $1 + x \leq \exp(x)$  for all  $x > -1$ . Now, for  $n \geq 2$  and  $0 < x \leq 1$ ,

$$\begin{aligned} \left(1 + \frac{x}{n}\right)^n &= 1 + \binom{n}{1} \frac{x}{n} + \binom{n}{2} \frac{x^2}{n^2} + \binom{n}{3} \frac{x^3}{n^3} + \cdots + \binom{n}{n} \frac{x^n}{n^n} \\ &= 1 + x + x^2 \left( \frac{1}{2!} \left(\frac{1}{n}\right) \left(1 - \frac{1}{n}\right) + \frac{1}{3!} \left(\frac{1}{n}\right) \left(1 - \frac{1}{n}\right) \left(1 - \frac{2}{n}\right) x + \cdots + \frac{1}{n!} \left(\frac{1}{n}\right) \left(1 - \frac{1}{n}\right) \left(1 - \frac{2}{n}\right) \cdots \left(1 - \frac{n-1}{n}\right) x^{n-2} \right) \\ &\leq 1 + x + x^2 \left( \frac{1}{2!} + \frac{1}{3!} + \cdots + \frac{1}{n!} \right) \\ &< 1 + x + x^2(e - 2), \end{aligned}$$

upon using Theorem 180. This proves the first set of inequalities.

For  $x > -2$ ,  $1 + x + \frac{x^2}{4} = \left(1 + \frac{x}{2}\right)^2 \leq \exp(x)$  by Proposition 301. Now we assume that  $-\frac{1}{2} \leq x \leq 0$ . As before,

$$\left(1 + \frac{x}{n}\right)^n = 1 + x + x^2 \left( \frac{1}{2!} \left(\frac{1}{n}\right) \left(1 - \frac{1}{n}\right) + \frac{1}{3!} \left(\frac{1}{n}\right) \left(1 - \frac{1}{n}\right) \left(1 - \frac{2}{n}\right) x + \cdots + \frac{1}{n!} \left(\frac{1}{n}\right) \left(1 - \frac{1}{n}\right) \left(1 - \frac{2}{n}\right) \cdots \left(1 - \frac{n-1}{n}\right) x^{n-2} \right).$$

Since  $x^k \leq 0$  for odd  $k$  and  $x^k \leq \frac{1}{2^k}$  for even  $k$  we may delete the odd terms from the dextral side and so

$$\begin{aligned} \left(1 + \frac{x}{n}\right)^n &\leq 1 + x + x^2 \left( \frac{1}{2!} \left(\frac{1}{n}\right) \left(1 - \frac{1}{n}\right) + 0 + \cdots + \frac{1}{n!} \left(\frac{1}{n}\right) \left(1 - \frac{1}{n}\right) \left(1 - \frac{2}{n}\right) \cdots \left(1 - \frac{2k-1}{n}\right) x^{2k} + \cdots \right) \\ &\leq 1 + x + x^2 \left( \frac{1}{2} + \frac{1}{2^2} + \cdots \right) \\ &\leq 1 + x + x^2. \end{aligned}$$

On taking limits  $\exp(x) \leq 1 + x + x^2$  for  $-\frac{1}{2} \leq x \leq 0$ . Thus we have

$$-\frac{1}{2} \leq x < 0 \implies 1 + x + \frac{x^2}{4} \leq \exp(x) \leq 1 + x + x^2 \implies 1 + x \leq \frac{\exp(x) - 1}{x} \leq 1 + \frac{x}{4},$$

since division by negative  $x$  reverses the sense of the inequalities.  $\square$

**340 THEOREM**  $\lim_{x \rightarrow 0} \frac{\exp(x) - 1}{x} = 1.$

**Proof:** We prove that  $\lim_{x \rightarrow 0^+} \frac{\exp(x) - 1}{x} = 1$  and that  $\lim_{x \rightarrow 0^-} \frac{\exp(x) - 1}{x} = 1$ . Let us start with the first assertion. For  $0 < x \leq 1$  we have, by the Sandwich Theorem, and Lemma 339,

$$1 \leq \frac{\exp(x) - 1}{x} \leq 1 + x(e - 2) \implies \lim_{x \rightarrow 0^+} \frac{\exp(x) - 1}{x} = 1,$$

proving the first assertion.

For  $-\frac{1}{2} \leq x \leq 0$  we have, by the Sandwich Theorem and Lemma 339,

$$1 + x + \frac{x^2}{4} \leq \exp(x) \leq 1 + x + x^2 \implies 1 + \frac{x}{4} \leq \frac{\exp(x) - 1}{x} \leq 1 + x \implies \lim_{x \rightarrow 0^-} \frac{\exp(x) - 1}{x} = 1,$$

proving the second assertion.  $\square$

**341 LEMMA** For  $0 < x \leq 1$ ,

$$1 - \frac{x(e - 2)}{1 + x} \leq \frac{\log(1 + x)}{x} \leq 1$$

and for  $-\frac{1}{2} \leq x \leq 0$ ,

$$1 \leq \frac{\log(1 + x)}{x} \leq 1 - \frac{x}{1 + x}.$$

**Proof:** Since  $x \mapsto \log(1 + x)$  is strictly increasing, we have by Lemma 339 for  $0 < x \leq 1$ ,

$$1 + x \leq \exp(x) \leq 1 + x + x^2(e - 2) \implies \log(1 + x) \leq x \leq \log(1 + x + x^2(e - 2)).$$

Notice that we have established that  $\log(1 + x) \leq x$  for  $0 < x \leq 1$ . Now

$$\log(1 + x + x^2(e - 2)) = \log(1 + x) \left(1 + \frac{x^2(e - 2)}{1 + x}\right) = \log(1 + x) + \left(1 + \frac{x^2(e - 2)}{1 + x}\right).$$

Since for  $x > 0$ ,  $x \mapsto \frac{x^2}{1 + x}$  is strictly increasing,  $\frac{x^2(e - 2)}{1 + x} < \frac{e - 2}{2} < 1$  for  $0 < x < 1$ . Thus we may use  $\log(1 + y) \leq y$ ,  $0 \leq y \leq 1$  with  $y = \frac{x^2(e - 2)}{1 + x}$  obtaining

$$\log\left(1 + \frac{x^2(e - 2)}{1 + x}\right) \leq \frac{x^2(e - 2)}{1 + x}.$$

Hence

$$x \leq \log(1 + x + x^2(e - 2)) \leq \log(1 + x) + \frac{x^2(e - 2)}{1 + x}.$$

In conclusion,

$$0 < x \leq 1 \implies \log(1 + x) \leq x \leq \log(1 + x) + \frac{x^2(e - 2)}{1 + x} \implies 1 - \frac{x(e - 2)}{1 + x} \leq \frac{\log(1 + x)}{x} \leq 1.$$

Similarly, for  $-\frac{1}{2} \leq x < 0$ , by Lemma 339,

$$1 + x + \frac{x^2}{4} \leq \exp(x) \leq 1 + x + x^2 \implies \log\left(1 + x + \frac{x^2}{4}\right) \leq x \leq \log(1 + x + x^2).$$

Since  $x \mapsto \log(1 + x)$  is increasing, plainly

$$\log(1 + x) \leq \log\left(1 + x + \frac{x^2}{4}\right) \leq x.$$

Now observe that  $-\frac{1}{2} \leq x < 0 \implies 0 < \frac{x^2}{1+x} \leq \frac{1}{2} < 1$  and so

$$\log(1+x+x^2) = \log(1+x) + \log\left(1 + \frac{x^2}{1+x}\right) \leq \log(1+x) + \frac{x^2}{1+x} \implies x \leq \log(1+x) + \frac{x^2}{1+x}.$$

In conclusion,

$$-\frac{1}{2} \leq x < 0 \implies \log(1+x) \leq x \leq \log(1+x) + \frac{x^2}{1+x} \implies 1 \leq \frac{\log(1+x)}{x} \leq 1 - \frac{x}{1+x},$$

since division by negative  $x$  reverses the sense of the inequalities.  $\square$

**342 THEOREM**  $\lim_{x \rightarrow 0} \frac{\log(1+x) - x}{x} = 0.$

**Proof:** By Lemma 341, for  $0 < x \leq 1$ ,

$$1 - \frac{x(e-2)}{1+x} \leq \frac{\log(1+x)}{x} \leq 1 \implies \lim_{x \rightarrow 0^+} \frac{\log(1+x)}{x} = 1,$$

by the Sandwich Theorem. Again, by Lemma 341 and the Sandwich Theorem,

$$-\frac{1}{2} \leq x \leq 0 \implies 1 \leq \frac{\log(1+x)}{x} \leq 1 - \frac{x}{1+x} \implies \lim_{x \rightarrow 0^-} \frac{\log(1+x)}{x} = 1.$$

Combining both results, the theorem follows.  $\square$




Figure 5.1: Theorem 344.

**343 THEOREM** If  $a \in \mathbb{R}$ , then  $\lim_{x \rightarrow 0} \frac{(1+x)^a - 1}{x} = a.$

**Proof:** This is evident for  $a = 0$ . Assume now  $a \neq 0$ . Since  $x \mapsto \exp(x)$  is continuous and since  $a \log(1+x) \rightarrow 0$  as  $x \rightarrow 0$ , by Theorems 340 and 342,

$$\lim_{x \rightarrow 0} \frac{(1+x)^a - 1}{x} = a \lim_{x \rightarrow 0} \frac{\exp(a \log(1+x)) - 1}{a \log(1+x)} \cdot \lim_{x \rightarrow 0} \frac{\log(1+x)}{x} = a \cdot 1 \cdot 1 = a.$$

$\square$

**344 THEOREM**  $\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1.$

**Proof:** We first prove that  $\lim_{\theta \rightarrow 0^+} \frac{\sin \theta}{\theta} = 1$ . Since  $\theta \mapsto \frac{\sin \theta}{\theta}$  is an even function it will also follow that  $\lim_{\theta \rightarrow 0^-} \frac{\sin \theta}{\theta} = 1$ .

Assume  $0 < \theta < \frac{\pi}{2}$  and consider  $\triangle OAB$  right-angled at  $A$ , with  $OA = 1$  and  $\angle BOA = \theta$ .  $C$  is the point where line  $OB$  meets the unit circle with centre at  $O$  and  $D$  is its perpendicular projection. The area of  $\triangle OAC$  is smaller than the area of the circular sector  $OAC$ , which is smaller than the area of  $\triangle OAB$ . Hence

$$\frac{1}{2} \sin \theta < \frac{\theta}{2} < \frac{1}{2} \tan \theta \implies \frac{1}{\cos \theta} < \frac{\sin \theta}{\theta} < 1 \implies \lim_{\theta \rightarrow 0^+} \frac{\sin \theta}{\theta} = 1$$

by the Sandwich Theorem, proving the theorem.  $\square$

# Chapter 6

## Differentiable Functions

### 6.1 Derivative at a Point

**345 Definition** Let  $I$  be an interval,  $a \in \overset{\circ}{I}$ , and  $f : I \rightarrow \mathbb{R}$ . We say that  $f$  is differentiable at  $a$  if the limit

$$\lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

exists and is finite. In such a case we denote this limit by  $f'(a)$ ,  $Df(a)$ , or  $\frac{df}{dx}(a)$  and we call this quantity *the derivative of  $f$  at  $a$* .

**346 Definition** Let  $I$  be an interval,  $a \in \overset{\circ}{I}$ , and  $f : I \rightarrow \mathbb{R}$ . If

$$\lim_{x \rightarrow a^+} \frac{f(x) - f(a)}{x - a} = \lim_{h \rightarrow 0^+} \frac{f(a+h) - f(a)}{h}$$

exists and is finite we say that  $f$  is differentiable at  $a$  on the right and write  $f'_+(a)$  for this limit. If

$$\lim_{x \rightarrow a^-} \frac{f(x) - f(a)}{x - a} = \lim_{h \rightarrow 0^-} \frac{f(a+h) - f(a)}{h}$$

exists and is finite we say that  $f$  is differentiable at  $a$  on the left and write  $f'_-(a)$  for this limit.

**347 THEOREM** Let  $I$  be an interval,  $a \in \overset{\circ}{I}$ , and  $f : I \rightarrow \mathbb{R}$ . Then  $f$  is differentiable at  $a$  if and only if both  $f_+(a)$  and  $f_-(a)$  exist and are equal. In this case  $f_+(a) = f'(a) = f_-(a)$ .

**Proof:** *Obvious.*  $\square$

**348 THEOREM** Let  $I$  be an interval,  $a \in \overset{\circ}{I}$ , and  $f : I \rightarrow \mathbb{R}$ . If  $f$  is differentiable at  $a$  then it is continuous at  $a$ .

**Proof:** *We have*

$$\lim_{h \rightarrow 0} f(a+h) - f(a) = \lim_{h \rightarrow 0} \left( \frac{f(a+h) - f(a)}{h} \right) h = \left( \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h} \right) \left( \lim_{h \rightarrow 0} h \right) = f'(a) \cdot 0 = 0.$$

Thus  $\lim_{h \rightarrow 0} f(a+h) - f(a) = 0 \implies \lim_{h \rightarrow 0} f(a+h) = f(a)$  and so  $f$  is continuous.  $\square$

**349 THEOREM** Let  $I \subseteq \mathbb{R}$  be an interval. If  $f : I \rightarrow \mathbb{R}$  is identically constant, then  $f'(I) = 0$ .

**Proof:** Assume that  $f(I) = K$ , a constant. Let  $c \in \overset{\circ}{I}$ . Then  $f'(c) = \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} = \lim_{x \rightarrow c} \frac{K - K}{x - c} = 0$ . If  $c$  is an endpoint of  $I$ , then the argument is modified to be either the left or right derivative.  $\square$

### Homework

**Problem 6.1.1** Let  $f: \mathbb{R} \rightarrow \mathbb{R}$ ,

$$f(x) \begin{cases} x+1 & \text{if } x \in \mathbb{Q} \\ 2-x & \text{if } x \in \mathbb{R} \setminus \mathbb{Q} \end{cases}$$

Prove that  $f$  is nowhere differentiable.

**Problem 6.1.2** Let  $f: \mathbb{R} \rightarrow \mathbb{R}$ ,  $x \mapsto |x|$ . Prove that  $f$  is not differentiable at  $x = 0$  and that for  $x \neq 0$ ,  $f'(x) = \text{signum}(x)$ .

**Problem 6.1.3** Let  $f: \mathbb{R} \rightarrow \mathbb{R}$ ,  $x \mapsto x|x|$ . Determine whether  $f'(0)$  exists.

## 6.2 Differentiation Rules

**350 THEOREM** Let  $I$  be an interval,  $a \in I$ ,  $\lambda \in \mathbb{R}$  a constant, and  $f$ , then,  $g: I \rightarrow \mathbb{R}$ . If  $f$  and  $g$  are differentiable at  $a$  then

1. **(Linearity Rule)**  $f + \lambda g$  is differentiable at  $a$  and  $(f + \lambda g)'(a) = f'(a) + \lambda g'(a)$
2. **(Product Rule)**  $f g$  is differentiable at  $a$  and  $(f g)'(a) = f'(a)g(a) + f(a)g'(a)$
3. if  $g(a) \neq 0$ ,  $\frac{1}{g}$  is differentiable at  $a$  and  $\left(\frac{1}{g}\right)'(a) = -\frac{g'(a)}{(g(a))^2}$
4. **(Quotient Rule)** if  $g(a) \neq 0$ ,  $\frac{f}{g}$  is differentiable at  $a$  and  $\left(\frac{f}{g}\right)'(a) = \frac{f'(a)g(a) - f(a)g'(a)}{(g(a))^2}$

**Proof:**

1. This follows by the linearity of limits.
2. We have

$$\begin{aligned} (fg)'(a) &= \lim_{h \rightarrow 0} \frac{(fg)(a+h) - (fg)(a)}{h} \\ &= \lim_{h \rightarrow 0} \frac{g(a+h)(f(a+h) - f(a)) + f(a)(g(a+h) - g(a))}{h} \\ &= \lim_{h \rightarrow 0} g(a+h) \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h} + \lim_{h \rightarrow 0} f(a) \lim_{h \rightarrow 0} \frac{g(a+h) - g(a)}{h} \\ &= g(a)f'(a) + f(a)g'(a), \end{aligned}$$

as desired.

3. We have

$$\begin{aligned} \left(\frac{1}{g}\right)'(a) &= \lim_{h \rightarrow 0} \frac{\frac{1}{g(a+h)} - \frac{1}{g(a)}}{h} \\ &= \lim_{h \rightarrow 0} \frac{g(a) - g(a+h)}{g(a+h)g(a)h} \\ &= \lim_{h \rightarrow 0} \frac{g(a) - g(a+h)}{h} \lim_{h \rightarrow 0} \frac{1}{g(a+h)g(a)} \\ &= (-g'(a)) \left(\frac{1}{g(a)g(a)}\right) \\ &= -\frac{g'(a)}{g(a)^2}, \end{aligned}$$

as desired.

4. Using (2) and (3),

$$\begin{aligned} \left(\frac{f}{g}\right)'(a) &= f'(a) \left(\frac{1}{g}\right)'(a) + f(a) \left(\frac{1}{g}\right)'(a) \\ &= \frac{f'(a)}{g(a)} - \frac{f(a)g'(a)}{g(a)^2} \\ &= \frac{f'(a)g(a) - f(a)g'(a)}{(g(a))^2}, \end{aligned}$$

as desired.

□

**351 THEOREM (Chain Rule)** Let  $I, J$  be intervals of  $\mathbb{R}$ , with  $a \in I$ . Let  $f: I \rightarrow \mathbb{R}$  and  $g: J \rightarrow \mathbb{R}$  be such that  $f(I) \subseteq J$ . If  $f$  is differentiable at  $a$  and  $g$  is differentiable at  $f(a)$ , then  $g \circ f$  is differentiable at  $a$  and  $(g \circ f)' = g'(f(a))f'(a)$ .

**Proof:** Put  $b = f(a)$ , and

$$\varphi(y) = \begin{cases} \frac{g(y) - g(b)}{y - b} & \text{if } y \neq b \\ g'(b) & \text{if } y = b \end{cases}$$

Since  $g$  is differentiable at  $b$ ,  $\varphi$  is continuous at  $y = b$ . Now, for  $x \neq a$ ,

$$\frac{g(f(x)) - g(f(a))}{x - a} = \varphi(f(x)) \frac{f(x) - f(a)}{x - a}.$$

(If  $f(x) \neq f(a)$  this follows directly from the definition of  $\varphi$ . If  $f(x) = f(a)$ , both sides of the equality are  $0$ .)

By the continuity of  $f$  at  $a$  and of  $\varphi$  at  $b$ ,

$$\lim_{x \rightarrow a} \varphi(f(x)) = \varphi(f(a)) = g'(f(a)),$$

whence

$$\begin{aligned} (g \circ f)'(a) &= \lim_{x \rightarrow a} \frac{g(f(x)) - g(f(a))}{x - a} \\ &= \lim_{x \rightarrow a} \varphi(f(x)) \frac{f(x) - f(a)}{x - a} \\ &= g'(f(a))f'(a), \end{aligned}$$

as desired.

□


**352 THEOREM (Inverse Function Rule)** Let  $I$  be an interval of  $\mathbb{R}$ , with  $a \in I$ . Let  $f: I \rightarrow \mathbb{R}$  be strictly monotonic and continuous over  $I$ . If  $f$  is differentiable at  $a$  and  $f'(a) \neq 0$ , then the inverse  $f^{-1}: f(I) \rightarrow \mathbb{R}$  is differentiable at  $f(a)$  and

$$(f^{-1})'(f(a)) = \frac{1}{f'(a)}.$$

**Proof:** Put  $b = f(a)$ . Observe that  $\lim_{y \rightarrow b} f^{-1}(y) = a$ , and by the composition rule for limits,

$$\lim_{y \rightarrow b} \frac{f^{-1}(y) - f^{-1}(b)}{y - b} = \lim_{y \rightarrow b} \frac{f^{-1}(y) - a}{f(f^{-1}(y)) - a} = \frac{1}{f'(a)},$$

proving the theorem. □

 Once it is known that  $(f^{-1})'$  exists, we may proceed as follows. Since  $f^{-1}(f(x)) = x$ , differentiating on both sides, using the Chain Rule on the sinistral side,

$$(f^{-1})'(f(x))f'(x) = 1,$$

from where the result follows.

**353 Definition** Let  $I$  be an interval of  $\mathbb{R}$ . Let  $f: I \rightarrow \mathbb{R}$  be differentiable at every point of  $I$ . The function  $f': I \rightarrow \mathbb{R}$ ,  $x \mapsto f'(x)$  is called the *derivative function* or *derivative* of the function  $f$ .

**354 THEOREM** Let  $n \geq 0$  be an integer. Let  $f: \mathbb{R} \rightarrow \mathbb{R}$ ,  $x \mapsto x^n$ . Then  $f$  is everywhere differentiable and  $f': \mathbb{R} \rightarrow \mathbb{R}$  is given by  $x \mapsto nx^{n-1}$ .

**Proof:** Assume first  $n$  is strictly positive. By Theorem 55,

$$\begin{aligned}\lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} &= \lim_{x \rightarrow a} \frac{(x - a)(x^{n-1} + ax^{n-2} + a^2x^{n-3} + \cdots + a^{n-2}x + a^{n-1})}{x - a} \\ &= \lim_{x \rightarrow a} (x^{n-1} + ax^{n-2} + a^2x^{n-3} + \cdots + a^{n-2}x + a^{n-1}) \\ &= na^{n-1}.\end{aligned}$$

Observe that this is true for all  $a \in \mathbb{R}$ .

If  $n = 0$  then  $f$  is constant, say  $f(x) = K$  for all  $x$  and so

$$\lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} = \lim_{x \rightarrow a} \frac{K - K}{x - a} = 0.$$

□

**355 THEOREM** Let  $n > 0$  be an integer and  $f: ]0; +\infty[ \rightarrow ]0; +\infty[$ ,  $x \mapsto \frac{1}{x^n}$ . Then  $f'$  exists everywhere in  $]0; +\infty[$  and  $f': ]0; +\infty[ \rightarrow ]0; +\infty[$  is given by  $f'(x) = -\frac{n}{x^{n+1}}$ .

**Proof:** We use the result above, part (3) of Theorem 350, and the Chain Rule, to get

$$\frac{d}{dx} \frac{1}{x^n} = -\frac{nx^{n-1}}{(x^n)^2} = -\frac{n}{x^{n+1}},$$

and the theorem follows. □

**356 LEMMA** Let  $q \in \mathbb{Z}$ ,  $q > 0$  be an integer, and  $f: ]0; +\infty[ \rightarrow ]0; +\infty[$ ,  $x \mapsto x^{1/q}$ . Then  $f'$  exists everywhere in  $]0; +\infty[$  and  $f': ]0; +\infty[ \rightarrow ]0; +\infty[$  is given by  $f'(x) = \frac{x^{1/q-1}}{q}$ .

**Proof:** We have  $(f(x))^q = x$ . Using the Chain Rule  $qf'(x)(f(x))^{q-1} = 1$ . Since  $f(x) \neq 0$ ,

$$f'(x) = \frac{1}{q(f(x))^{q-1}} = \frac{1}{q(x^{1/q})^{q-1}} = \frac{1}{q}x^{1/q-1}.$$

□

**357 THEOREM** Let  $r \in \mathbb{Q}$  and let  $f: ]0; +\infty[ \rightarrow ]0; +\infty[$ ,  $x \mapsto x^r$ . Then  $f'$  exists everywhere in  $]0; +\infty[$  and  $f': ]0; +\infty[ \rightarrow ]0; +\infty[$  is given by  $f'(x) = rx^{r-1}$ .

**Proof:** Let  $r = \frac{a}{b}$ , where  $a, b$  are integers, with  $b > 0$ . We use the Chain Rule, Lemma 356, and Theorem 355. Then

$$\frac{d}{dx} x^{a/b} = \frac{d}{dx} (x^{1/b})^a = a(x^{1/b})^{a-1} \cdot \frac{1}{b} x^{1/b-1} = \frac{a}{b} x^{a/b-1} = rx^{r-1},$$

proving the theorem.

□

**358 THEOREM (Derivative of the Exponential Function)** Let  $\exp: \mathbb{R} \rightarrow \mathbb{R}$ ,  $x \mapsto e^x$ . Then  $\exp$  is everywhere differentiable and  $\exp': \mathbb{R} \rightarrow \mathbb{R}$  is given by  $x \mapsto e^x$ .



**Proof:** Using Theorem 340, we have, with  $h = x - a$ ,

$$\begin{aligned} \lim_{x \rightarrow a} \frac{e^x - e^a}{x - a} &= e^a \lim_{x \rightarrow a} \frac{e^{x-a} - 1}{x - a} \\ &= e^a \lim_{h \rightarrow 0} \frac{e^h - 1}{h} \\ &= e^a \cdot 1 \\ &= e^a. \end{aligned}$$

□

**359 THEOREM (Derivative of the Logarithmic Function)** Let  $f: ]0; +\infty[ \rightarrow ]-\infty; +\infty[$ ,  $x \mapsto \log x$ . Then  $f'$  exists everywhere in  $]0; +\infty[$  and  $f': ]0; +\infty[ \rightarrow \mathbb{R} \setminus \{0\}$  is given by  $f'(x) = \frac{1}{x}$ .

**Proof:** Let  $a > 0$ . Then, with  $h = \frac{x}{a} - 1$ , and using Theorem 342,

$$\begin{aligned} \lim_{x \rightarrow a} \frac{\log x - \log a}{x - a} &= \lim_{x \rightarrow a} \frac{\log \frac{x}{a}}{x - a} \\ &= \frac{1}{a} \cdot \lim_{x \rightarrow a} \frac{\log\left(1 + \frac{x}{a} - 1\right)}{\frac{x}{a} - 1} \\ &= \frac{1}{a} \cdot \lim_{h \rightarrow 0} \frac{\log(1 + h)}{h} \\ &= \frac{1}{a} \cdot 1 \\ &= \frac{1}{a}. \end{aligned}$$

□

**360 THEOREM (Power Rule)** Let  $t \in \mathbb{R}$  and let  $f: ]0; +\infty[ \rightarrow ]0; +\infty[$ ,  $x \mapsto x^t$ . Then  $f'$  exists everywhere in  $]0; +\infty[$  and  $f': ]0; +\infty[ \rightarrow ]0; +\infty[$  is given by  $f'(x) = tx^{t-1}$ .

**Proof:** Using the Chain Rule,

$$\frac{d}{dx} x^t = \frac{d}{dx} (\exp(t \log x)) = \frac{t}{x} \cdot (\exp(t \log x)) = \frac{t}{x} \cdot x^t = tx^{t-1}.$$

□

**361 THEOREM (Derivative of sin)** . Let  $\sin: \mathbb{R} \rightarrow \mathbb{R}$ ,  $x \mapsto \sin x$ . Then  $\sin$  is everywhere differentiable and  $\sin': \mathbb{R} \rightarrow \mathbb{R}$  is given by  $x \mapsto \cos x$ .

**Proof:** We make a change of variables, and use Theorem 344,

$$\begin{aligned}
 \lim_{x \rightarrow a} \frac{\sin x - \sin a}{x - a} &= \lim_{x \rightarrow a} \frac{\sin(x - a + a) - \sin a}{x - a} \\
 &= \lim_{x \rightarrow a} \frac{\sin(x - a) \cos a + \cos(x - a) \sin a - \sin a}{x - a} \\
 &= (\cos a) \lim_{x \rightarrow a} \frac{\sin(x - a)}{x - a} + (\sin a) \lim_{x \rightarrow a} \frac{\cos(x - a) - 1}{x - a} \\
 &= (\cos a) \lim_{h \rightarrow 0} \frac{\sin h}{h} + (\sin a) \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} \\
 &= (\cos a) \cdot 1 + (\sin a) \lim_{h \rightarrow 0} \frac{\cos^2 h - 1}{h(\cos h + 1)} \\
 &= (\cos a) \cdot 1 + (\sin a) \lim_{h \rightarrow 0} \frac{-\sin^2 h}{h(\cos h + 1)} \\
 &= (\cos a) + (\sin a) \lim_{h \rightarrow 0} \frac{\sin h}{h} \cdot \lim_{h \rightarrow 0} \frac{-\sin h}{\cos h + 1} \\
 &= \cos a,
 \end{aligned}$$

and the theorem follows.  $\square$

### 362 THEOREM (Derivatives of the Goniometric Functions)

1.  $\frac{d}{dx} \sin x = \cos x \quad x \in \mathbb{R}$
2.  $\frac{d}{dx} \cos x = -\sin x \quad x \in \mathbb{R}$
3.  $\frac{d}{dx} \tan x = \sec^2 x \quad x \in \mathbb{R} \setminus (2\mathbb{Z} + 1)\frac{\pi}{2}$
4.  $\frac{d}{dx} \sec x = \sec x \tan x \quad x \in \mathbb{R} \setminus (2\mathbb{Z} + 1)\frac{\pi}{2}$
5.  $\frac{d}{dx} \csc x = -\csc x \cot x \quad x \in \mathbb{R} \setminus \mathbb{Z}\pi$
6.  $\frac{d}{dx} \cot x = -\csc^2 x \quad x \in \mathbb{R} \setminus \mathbb{Z}\pi$

**Proof:** (1) is Theorem 361. To prove (2), observe that

$$\frac{d}{dx} \cos x = \frac{d}{dx} \sin\left(\frac{\pi}{2} - x\right) = -\cos\left(\frac{\pi}{2} - x\right) = -\sin x.$$

To prove (3), we use the Quotient Rule,

$$\frac{d}{dx} \tan x = \frac{d}{dx} \frac{\sin x}{\cos x} = \frac{(\cos x)(\cos x) - (-\sin x)(\sin x)}{\cos^2 x} = \frac{1}{\cos^2 x} = \sec^2 x.$$

To prove (4), we use once again the Quotient Rule,

$$\frac{d}{dx} \sec x = \frac{d}{dx} \frac{1}{\cos x} = \frac{(0)(\cos x) - (-\sin x)(1)}{\cos^2 x} = \frac{\sin x}{\cos^2 x} = \sec x \tan x.$$

To prove (5), observe that

$$\frac{d}{dx} \csc x = \frac{d}{dx} \sec\left(\frac{\pi}{2} - x\right) = -\sec\left(\frac{\pi}{2} - x\right) \tan\left(\frac{\pi}{2} - x\right) = -\csc x \cot x.$$

To prove (6), observe that

$$\frac{d}{dx} \cot x = \frac{d}{dx} \tan\left(\frac{\pi}{2} - x\right) = -\sec^2\left(\frac{\pi}{2} - x\right) = -\csc^2 x.$$

$\square$

**363 Definition (Higher Order Derivatives)** Let  $I$  be an interval of  $\mathbb{R}$  and let  $f : I \rightarrow \mathbb{R}$ . For  $a \in I$  we define the successive derivatives of  $f$  at  $a$ , inductively. Put  $f(a) = f^{(0)}(a)$ . If  $n \geq 1$ ,

$$f^{(n)}(a) = f'(f^{(n-1)}(a)),$$

provided  $f$  is differentiable at  $f^{(n-1)}(a)$ .



We usually write  $f''$  instead of  $f^{(2)}$ .

**364 THEOREM (Leibniz's Rule)** Let  $n$  be a positive integer.

$$(fg)^{(n)} = \sum_{k=0}^n \binom{n}{k} f^{(k)} g^{(n-k)}$$

**Proof:** This is a generalisation of the Product Rule. The proof is by induction on  $n$ . For  $n = 0$  and  $n = 1$  the assertion is obvious. Assume that  $(fg)^{(n)} = \sum_{k=0}^n \binom{n}{k} f^{(k)} g^{(n-k)}$ . Observe that

$$\begin{aligned} (fg)^{(n+1)} &= ((fg)^{(n)})' \\ &= \left( \sum_{k=0}^n \binom{n}{k} f^{(k)} g^{(n-k)} \right)' \\ &= \sum_{k=0}^n \binom{n}{k} (f^{(k+1)} g^{(n-k)} + f^{(k)} g^{(n-k+1)}) \\ &= \sum_{k=0}^n \binom{n}{k} f^{(k+1)} g^{(n-k)} + \sum_{k=0}^n \binom{n}{k} f^{(k)} g^{(n-k+1)} \\ &= f^{(0)} g^{(n+1)} + \sum_{k=0}^n \left( \binom{n}{k} + \binom{n}{k+1} \right) f^{(k)} g^{(n+1-k)} + f^{(n+1)} g^{(0)} \\ &= \sum_{k=0}^{n+1} \binom{n+1}{k} f^{(k)} g^{(n+1-k)}, \end{aligned}$$

proving the statement.

□

## Homework

**Problem 6.2.1** Prove that

$$\frac{2}{x^2-1} = \frac{1}{x-1} - \frac{1}{x+1}$$

and use this result to find the 100th derivative of  $f(x) = \frac{2}{x^2-1}$ .

**Problem 6.2.2** Find the 100-th derivative of  $x \mapsto x^2 \sin x$ .

**Problem 6.2.3** Demonstrate that the polynomial  $p(x) \in \mathbb{R}[x]$  has a zero at  $x = a$  of multiplicity  $k$  if and only if

$$p(a) = p'(a) = \dots = p^{(k-1)}(a) = 0.$$

**Problem 6.2.4** Demonstrate that if for all  $x \in \mathbb{R}$  there holds the identity

$$\sum_{k=0}^n a_k (x-a)^k = \sum_{k=0}^n b_k (x-b)^k,$$

then  $a_k = \sum_{j=k}^n \binom{n}{j} b_j (a-b)^{j-k}$ .

**Problem 6.2.5** Let  $p$  be a polynomial of degree  $r$  and consider the polynomial  $F$  with

$$F(x) = p(x) + p'(x) + p''(x) + \dots + p^{(r)}(x).$$

Prove that

$$\frac{d(F(x) \exp(-x))}{dx} = -\exp(-x)p(x).$$

### 6.3 Rolle's Theorem and the Mean Value Theorem

**365 THEOREM (Rolle's Theorem)** Let  $(a, b) \in \mathbb{R}^2$  such that  $a < b$ ,  $f : [a; b] \rightarrow \mathbb{R}$  be such that  $f$  is continuous on  $[a; b]$  and differentiable in  $]a; b[$ , and  $f(a) = f(b)$ . Then there exists  $c \in ]a; b[$  such that  $f'(c) = 0$ .

**Proof:** Since  $f$  is continuous on  $[a; b]$ , by Weierstrass' Theorem 323,

$$m = \inf_{x \in [a; b]} f(x), \quad M = \sup_{x \in [a; b]} f(x),$$

exist. If  $m = M$ , then  $f$  is constant and so by Theorem 349,  $f'$  is identically 0 and there is nothing to prove. Assume that  $m < M$ . Since  $f(a) = f(b)$ , one may not simultaneously have  $M = f(a)$  and  $m = f(a)$ . Assume thus without loss of generality that  $M \neq f(a)$ . Then there exists  $c \in ]a; b[$  such that  $f(c) = M$ . Now

$$\lim_{x \rightarrow c^-} \frac{f(x) - f(c)}{x - c} \geq 0, \quad \lim_{x \rightarrow c^+} \frac{f(x) - f(c)}{x - c} \leq 0,$$

whence it follows that  $f'(c) = 0$ , proving the theorem.  $\square$

**366 THEOREM (Mean Value Theorem)** Let  $(a, b) \in \mathbb{R}^2$  such that  $a < b$ ,  $f : [a; b] \rightarrow \mathbb{R}$  be such that  $f$  is continuous on  $[a; b]$  and differentiable on  $]a; b[$ . Then there exists  $c \in ]a; b[$  such that  $f'(c) = \frac{f(b) - f(a)}{b - a}$ .

**Proof:** Put

$$g : [a; b] \rightarrow \mathbb{R}, \quad g(x) = f(x) - \frac{f(b) - f(a)}{b - a}x.$$

Then  $g$  is continuous on  $[a; b]$  and differentiable on  $]a; b[$ , and  $g(a) = g(b)$ . Since  $g$  satisfies the hypotheses of Rolle's Theorem, there is  $c \in ]a; b[$  such that

$$g'(c) = 0 \implies f'(c) - \frac{f(b) - f(a)}{b - a} = 0 \implies f'(c) = \frac{f(b) - f(a)}{b - a},$$

proving the theorem.  $\square$

**367 THEOREM** If  $f : I \rightarrow \mathbb{R}$  is continuous on the interval  $I$ , differentiable on  $\dot{I}$ , and if  $\forall x \in \dot{I}$ ,  $f'(x) = 0$  then  $f$  is constant on  $I$ .

**Proof:** Let  $(a, b) \in I^2$ ,  $a < b$ . By the Mean Value Theorem, there is  $c \in ]a; b[$  such that

$$f(b) - f(a) = f'(c)(b - a) = 0 \cdot (b - a) \implies f(b) = f(a),$$

thus any two outputs have exactly the same value and  $f$  is constant.  $\square$

**368 THEOREM** If  $f : I \rightarrow \mathbb{R}$  is continuous on the interval  $I$ , and differentiable on  $\dot{I}$ . Then  $f$  is increasing on  $I$  if and only if  $\forall x \in \dot{I}$ ,  $f'(x) \geq 0$  and  $f$  is decreasing on  $I$  if and only if  $\forall x \in \dot{I}$ ,  $f'(x) \leq 0$ .

**Proof:**

$\implies$  Suppose  $f$  is increasing. Let  $x_0 \in \dot{I}$ . If  $h \neq 0$  is so small that  $x_0 + h \in \dot{I}$ , then

$$\frac{f(x_0 + h) - f(x_0)}{h} \geq 0 \implies \lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h} \geq 0 \implies f'(x_0) \geq 0.$$

If  $f$  is decreasing we apply what has just been proved to  $-f$ .

$\Leftarrow$  Suppose that for all  $x \in \overset{\circ}{I}$ ,  $f'(x) \geq 0$ . Let  $(a, b) \in I^2$ ,  $a < b$ . By the Mean Value Theorem, there is  $c \in ]a; b[$  such that

$$f(b) - f(a) = (b - a)f'(c) \geq 0,$$

and so  $f$  is increasing. If for all  $x \in \overset{\circ}{I}$ ,  $f'(x) \leq 0$  we apply what we just proved to  $-f$ .

□

**369 THEOREM** If  $f : I \rightarrow \mathbb{R}$  is continuous on the interval  $I$ , and differentiable on  $\overset{\circ}{I}$ . Then  $f$  is strictly increasing on  $I$  if and only if  $\forall x \in I$ ,  $f'(x) \geq 0$  and the set  $\overline{\{x \in \overset{\circ}{I} : f'(x) = 0\}} = \emptyset$ . Also,  $f$  is strictly decreasing on  $I$  if and only if  $\forall x \in I$ ,  $f'(x) \leq 0$  and  $\overline{\{x \in \overset{\circ}{I} : f'(x) = 0\}} = \emptyset$ .

**Proof:**

$\Rightarrow$  Suppose  $f$  is strictly increasing. From Theorem 368 we know that  $\forall x \in \overset{\circ}{I}$ ,  $f'(x) \geq 0$ . Assume that  $\overline{\{x \in \overset{\circ}{I} : f'(x) = 0\}} \neq \emptyset$ . Then there is  $c \in \overline{\{x \in \overset{\circ}{I} : f'(x) = 0\}}$  and  $\varepsilon > 0$  such that  $]c - \varepsilon; c + \varepsilon[ \subseteq I$  and  $\forall x \in ]c - \varepsilon; c + \varepsilon[$ ,  $f'(x) = 0$ .

By Theorem 367,  $f$  must be constant on  $]c - \varepsilon; c + \varepsilon[$  and so it is not strictly increasing, a contradiction. If  $f$  is strictly decreasing, we apply what has been proved to  $-f$ .

$\Leftarrow$  Conversely, suppose that  $\forall x \in I$ ,  $f'(x) \geq 0$ . and the set  $\overline{\{x \in \overset{\circ}{I} : f'(x) = 0\}} = \emptyset$ . From Theorem 368,  $f$  is increasing on  $I$ . Suppose that there exist  $(a, b) \in I^2$ ,  $a < b$  such that  $f(a) = f(b)$ . Since  $f$  is increasing, we have  $\forall x \in ]a; b[$ ,  $f(x) = f(a)$ . But then  $]a; b[ \subseteq \overline{\{x \in \overset{\circ}{I} : f'(x) = 0\}}$ , a contradiction, since this last set was assumed empty. If  $f'(x) \leq 0$  we apply what has been proved to  $-f$ .

□

## Homework

**Problem 6.3.1** Show, by means of Rolle's Theorem, that  $5x^4 - 4x + 1 = 0$  has a solution in  $[0; 1]$ .

**Problem 6.3.2** Let  $a_0, a_1, \dots, a_n$  be real numbers satisfying

$$a_0 + \frac{a_1}{2} + \frac{a_2}{3} + \dots + \frac{a_n}{n+1} = 0.$$

Show that the polynomial

$$a_0 + a_1x + \dots + a_nx^n$$

has a root in  $]0; 1[$ .

**Problem 6.3.3** Let  $a, b, c$  be three functions such that  $a' = b$ ,  $b' = c$ , and  $c' = a$ . Prove that the function  $a^3 + b^3 + c^3 - 3abc$  is constant.

**Problem 6.3.4** Suppose that  $f : ]0; 1[ \rightarrow \mathbb{R}$  is differentiable,  $f(0) = 0$  and  $f(x) > 0$  for  $x \in ]0; 1[$ . Is there a number  $d \in ]0; 1[$  such that

$$\frac{2f'(c)}{f(c)} = \frac{f'(1-c)}{f(1-c)}?$$

**Problem 6.3.5** Let  $n \geq 1$  be an integer and let  $f : ]0; 1[ \rightarrow \mathbb{R}$  be differentiable and such that  $f(0) = 0$  and  $f(1) = 1$ . Prove that there exist distinct points  $0 < a_0 < a_1 < \dots < a_{n-1} < 1$  such that

$$\sum_{k=0}^{n-1} f'(a_k) = n.$$

**Problem 6.3.6** Let  $n \geq 1$  be an integer and let  $f : ]0; 1[ \rightarrow \mathbb{R}$  be differentiable and such that  $f(0) = 0$  and  $f(1) = 1$ . Prove that there exist distinct points  $0 < a_0 < a_1 < \dots < a_{n-1} < 1$  such that

$$\sum_{k=0}^{n-1} \frac{1}{f'(a_k)} = n.$$

**Problem 6.3.7 (Putnam 1946)** Let  $p(x)$  is a quadratic polynomial with real coefficients satisfying  $\max_{x \in [-1; 1]} |p(x)| \leq 1$ . Prove that

$$\max_{x \in [-1; 1]} |p'(x)| \leq 4.$$

**Problem 6.3.8 (Generalised Mean Value Theorem)** Let  $f, g$  be continuous of  $]a; b[$  and differentiable on  $]a; b[$ . Then there is  $c \in ]a; b[$  such that

$$(f(b) - f(a))g'(c) = (g(b) - g(a))f'(c).$$

**Problem 6.3.9 (First L'Hôpital Rule)** Let  $I$  be an open interval (finite or infinite) having  $c$  has an endpoint (which may be finite or infinite). Assume  $f, g$  are differentiable on  $I$ ,  $g$  and  $g'$  never vanish on  $I$  and that  $\lim_{x \rightarrow c} f(x) = 0 = \lim_{x \rightarrow c} g(x)$ . Prove that if  $\lim_{x \rightarrow c} \frac{f'(x)}{g'(x)} = L$  (where  $L$  is finite or infinite), then  $\lim_{x \rightarrow c} \frac{f(x)}{g(x)} = L$

**Problem 6.3.10 (Second L'Hôpital Rule)** Let  $I$  be an open interval (finite or infinite) having  $c$  as an endpoint (which may be finite or infinite). Assume  $f, g$  are differentiable on  $I$ ,  $g$  and  $g'$  never vanish on  $I$  and that  $\lim_{x \rightarrow c} |f(x)| = \lim_{x \rightarrow c} |g(x)| = +\infty$ . Prove that if

$$\lim_{x \rightarrow c} \frac{f'(x)}{g'(x)} = L \text{ (where } L \text{ is finite or infinite), then } \lim_{x \rightarrow c} \frac{f(x)}{g(x)} = L$$

**Problem 6.3.11** If  $f'$  exists on an interval containing  $c$ , then

$$f'(c) = \lim_{h \rightarrow 0} \frac{f(c+h) - f(c-h)}{2h}.$$

**Problem 6.3.12** If  $f''$  exists on an interval containing  $c$ , then

$$f''(c) = \lim_{h \rightarrow 0} \frac{f(c+h) + f(c-h) - 2f(c)}{h^2}.$$

## 6.4 Extrema

**370 Definition** Let  $X \subseteq \mathbb{R}$ ,  $f: X \rightarrow \mathbb{R}$ .

1. We say that  $f$  has a *local maximum at  $a$*  if there exists a neighbourhood of  $a$ ,  $\mathcal{N}_a$  such that  $\forall x \in \mathcal{N}_a, f(x) \leq f(a)$ .
2. We say that  $f$  has a *local minimum at  $a$*  if there exists a neighbourhood of  $a$ ,  $\mathcal{N}_a$  such that  $\forall x \in \mathcal{N}_a, f(x) \geq f(a)$ .
3. We say that  $f$  has a *strict local maximum at  $a$*  if there exists a neighbourhood of  $a$ ,  $\mathcal{N}_a$  such that  $\forall x \in \mathcal{N}_a, f(x) < f(a)$ .
4. We say that  $f$  has a *strict local minimum at  $a$*  if there exists a neighbourhood of  $a$ ,  $\mathcal{N}_a$  such that  $\forall x \in \mathcal{N}_a, f(x) > f(a)$ .
5. We say that  $f$  has a *local extremum at  $a$*  if  $f$  has either a local maximum or a local minimum at  $a$ .
6. We say that  $f$  has a *strict local extremum at  $a$*  if  $f$  has either a strict local maximum or a strict local minimum at  $a$ . The plural of extremum is *extrema*.

**371 THEOREM** If  $f: I \rightarrow \mathbb{R}$  is continuous on the interval  $I$ , differentiable on  $\overset{\circ}{I}$ , and if  $f$  has a local extremum at  $a \in \overset{\circ}{I}$ , then  $f'(a) = 0$ .

**Proof:** Suppose  $f$  admits a local maximum at  $a$ . Let  $h \neq 0$  be so small that  $a+h \in I$ . Now

$$h > 0 \implies \frac{f(a+h) - f(a)}{h} \leq 0, \quad h < 0 \implies \frac{f(a+h) - f(a)}{h} \geq 0.$$

Upon taking limits as  $h \rightarrow 0$ ,  $f'(a) \leq 0$  and  $f'(a) \geq 0$ , whence  $f'(a) = 0$ .  $\square$

**372 Definition** Let  $f: I \rightarrow \mathbb{R}$ . The points  $x \in I$  where  $f'(x) = 0$  are called *critical points* or *stationary points* of  $f$ .

**373 THEOREM** Let  $f: [a; b] \rightarrow \mathbb{R}$  be a twice differentiable function having a critical point at  $c \in ]a; b[$ . If  $f''(c) < 0$  then  $f$  has a relative maximum at  $x = c$ , and if  $f''(c) > 0$  then  $f$  has a relative minimum at  $x = c$ .

**Proof:** Assume that  $f'(c) = 0$  and  $f''(c) < 0$ . Since

$$\lim_{x \rightarrow c} \frac{f'(x)}{x-c} = \lim_{x \rightarrow c} \frac{f'(x) - f'(c)}{x-c} = f''(c) < 0,$$

there exists  $\delta > 0$  such that  $f'(x) > 0$  when  $c - \delta < x < c$  and  $f'(x) < 0$  when  $c < x < c + \delta$ . Consequently,  $f$  is strictly increasing on  $]c - \delta; c[$  and strictly decreasing on  $]c; c + \delta[$ . Hence

$$|x - c| < \delta \implies f(x) \leq f(c),$$

and so  $x = c$  is a local maximum. If  $f'' > 0$  then we apply what has been proved to  $-f$ .  $\square$

**374 THEOREM (Darboux's Theorem)** Let  $f$  be differentiable on  $[a; b]$  and suppose that  $f'(a) < C < f'(b)$ . Then there exists  $c \in ]a; b[$  such that  $f'(c) = C$ .

**Proof:** Put  $g(x) = f(x) - Cx$ . Then  $g$  is differentiable on  $[a; b]$ . Now  $g'(a) = f'(a) - C < 0$  so  $g$  is strictly increasing at  $x = a$ . Similarly,  $g'(b) = f'(b) - C < 0$  so  $g$  is strictly decreasing at  $x = b$ . Since  $g$  is continuous,  $g$  must have a local maximum at some point  $c \in ]a; b[$ , where  $g'(c) = f'(c) - C = 0$ , proving the theorem.  $\square$

## Homework

**Problem 6.4.1** Let  $f$  be a polynomial with real coefficients of degree  $n$  such that  $\forall x \in \mathbb{R} \quad f(x) \geq 0$ . Prove that

$$\forall x \in \mathbb{R} \quad f(x) + f'(x) + f''(x) + \dots + f^{(n)}(x) \geq 0.$$


**Problem 6.4.2** Put  $f(0) = 1$ ,  $f(x) = x^x$  for  $x > 0$ . Find the minimum value of  $f$ .

## 6.5 Convex Functions

**375 Definition** Let  $I \subseteq \mathbb{R}$  be an interval. A function  $f: I \rightarrow \mathbb{R}$  is said to be *convex* if

$$\forall (a, b) \in I^2, \forall \lambda \in [0; 1], f(\lambda a + (1 - \lambda)b) \leq \lambda f(a) + (1 - \lambda)f(b).$$

We say that  $f$  is *concave* if  $-f$  is convex.

  $f$  is convex if given any two points on its graph, the straight line joining these two points lies above the graph of  $f$ . See figure 6.1.




Figure 6.1: A convex curve




Figure 6.2: A concave curve.

**376 Definition** Let  $(x_1, x_2, \dots, x_n) \in \mathbb{R}^n$  and let  $\lambda_k \in [0; 1]$  be such that  $\sum_{k=1}^n \lambda_k = 1$ . The sum

$$\sum_{k=1}^n \lambda_k x_k$$

is called a *convex combination* of the  $x_k$ .

**377 THEOREM** If  $(x_1, x_2, \dots, x_n) \in [a; b]^n$ , then any convex combination of the  $x_k$  also belongs to  $[a; b]$ .

**Proof:** Assume  $\lambda_k \in [0; 1]$  be such that  $\sum_{k=1}^n \lambda_k = 1$ . Since the  $\lambda_k \geq 0$  we have

$$a \leq x_k \leq b \implies \lambda_k a \leq \lambda_k x_k \leq \lambda_k b.$$

Adding, and bearing in mind that  $\sum_{k=1}^n \lambda_k = 1$ ,

$$\left( \sum_{k=1}^n \lambda_k \right) a \leq \sum_{k=1}^n \lambda_k x_k \leq \left( \sum_{k=1}^n \lambda_k \right) b \implies a \leq \sum_{k=1}^n \lambda_k x_k \leq b,$$

proving the theorem.  $\square$

**378 THEOREM (Jensen's Inequality)** Let  $I \subseteq \mathbb{R}$  be an interval and let  $f : I \rightarrow \mathbb{R}$  be a convex function. Let  $n \geq 1$  be an integer,  $x_k \in I$ , and  $\lambda_k \in [0; 1]$  be such that  $\sum_{k=1}^n \lambda_k = 1$ . Then

$$f\left(\sum_{k=1}^n \lambda_k x_k\right) \leq \sum_{k=1}^n \lambda_k f(x_k).$$

**Proof:** The proof is by induction on  $n$ . For  $n = 2$  we must show that given  $(x_1, x_2) \in [a; b]^2$ ,

$$f(\lambda_1 x_1 + \lambda_2 x_2) \leq \lambda_1 f(x_1) + \lambda_2 f(x_2).$$

As  $\lambda_1 + \lambda_2 = 1$ , we may put  $\lambda = \lambda_2 = 1 - \lambda_1$  and so the above inequality becomes

$$f(\lambda x_1 + (1 - \lambda) x_2) \leq \lambda f(x_1) + (1 - \lambda) f(x_2),$$

retrieving the definition of convexity.

Assume now that  $f\left(\sum_{k=1}^{n-1} \mu_k x_k\right) \leq \sum_{k=1}^{n-1} \mu_k f(x_k)$ , when  $\sum_{k=1}^{n-1} \mu_k = 1$ ,  $\mu_k \in [0; 1]$ . We must prove that  $f\left(\sum_{k=1}^n \lambda_k x_k\right) \leq \sum_{k=1}^n \lambda_k f(x_k)$ , when  $\sum_{k=1}^n \lambda_k = 1$ ,  $\lambda_k \in [0; 1]$ .

If  $\lambda_n = 1$  the assertion is trivial, since then  $\lambda_1 = \dots = \lambda_{n-1} = 0$ . So assume that  $\lambda_n \neq 1$ . Observe that  $\sum_{k=1}^{n-1} \frac{\lambda_k}{1 - \lambda_n} = \frac{(\sum_{k=1}^n \lambda_k) - \lambda_n}{1 - \lambda_n} = \frac{1 - \lambda_n}{1 - \lambda_n} = 1$  so that  $\sum_{k=1}^{n-1} \frac{\lambda_k}{1 - \lambda_n} x_k$  is a convex combination of the  $x_k$  and hence also belongs to  $[a; b]$ , by Theorem 377. Since  $f$  is convex,

$$\begin{aligned} f\left(\sum_{k=1}^n \lambda_k x_k\right) &= f\left(\sum_{k=1}^{n-1} \lambda_k x_k + \lambda_n x_n\right) \\ &= f\left((1 - \lambda_n) \sum_{k=1}^{n-1} \frac{\lambda_k}{1 - \lambda_n} x_k + \lambda_n x_n\right) \\ &\leq (1 - \lambda_n) f\left(\sum_{k=1}^{n-1} \frac{\lambda_k}{1 - \lambda_n} x_k\right) + \lambda_n f(x_n) \end{aligned}$$

By the inductive hypothesis, with  $\mu_k = \frac{\lambda_k}{1 - \lambda_n} = 1$ ,

$$f\left(\sum_{k=1}^{n-1} \frac{\lambda_k}{1 - \lambda_n} x_k\right) \leq \sum_{k=1}^{n-1} \frac{\lambda_k}{1 - \lambda_n} f(x_k).$$

Finally, we gather,

$$\begin{aligned} f\left(\sum_{k=1}^n \lambda_k x_k\right) &\leq (1 - \lambda_n) f\left(\sum_{k=1}^{n-1} \frac{\lambda_k}{1 - \lambda_n} x_k\right) + \lambda_n f(x_n) \\ &\leq (1 - \lambda_n) \sum_{k=1}^{n-1} \frac{\lambda_k}{1 - \lambda_n} f(x_k) + \lambda_n f(x_n) \\ &= \sum_{k=1}^{n-1} \lambda_k f(x_k) + \lambda_n f(x_n) \\ &= \sum_{k=1}^n \lambda_k f(x_k), \end{aligned}$$

proving the theorem.  $\square$



**379 THEOREM** Let  $I \subseteq \mathbb{R}$  be an interval and let  $f: I \rightarrow \mathbb{R}$ . For  $a \in I$  we put

$$T_a: \begin{array}{l} I \setminus \{a\} \rightarrow \mathbb{R} \\ x \mapsto \frac{f(x) - f(a)}{x - a} \end{array}.$$

Then  $f$  is convex if and only if  $\forall a \in I$ ,  $T_a$  is increasing over  $I \setminus \{a\}$ .

**Proof:** Let  $a < b < c$  as in figure 6.3. Consider the points  $A(a, f(a))$ ,  $B(b, f(b))$ , and  $C(c, f(c))$ . The slopes

$$m_{AB} = \frac{f(b) - f(a)}{b - a}, \quad m_{BC} = \frac{f(c) - f(b)}{c - b}, \quad m_{CA} = \frac{f(c) - f(a)}{c - a},$$

satisfy

$$m_{AB} \leq m_{AC}, \quad m_{AC} \leq m_{BC}, \quad m_{AB} \leq m_{BC},$$

and the theorem follows. An analytic proof may be obtained by observing that from Theorem 377, any  $\lambda a + (1 - \lambda)c$  lies in the interval  $[a; c]$  for  $\lambda \in [0; 1]$ . Conversely, given  $b \in [a; c]$ , we may solve for  $\lambda$  the equation

$$b = \lambda a + (1 - \lambda)c \implies \lambda = \frac{c - b}{c - a} \in [0; 1].$$

Hence

$$f(\lambda a + (1 - \lambda)c) \leq \lambda f(a) + (1 - \lambda)f(c) \iff f(b) \leq \frac{c - b}{c - a}f(a) + \frac{b - a}{c - a}f(c) \iff \frac{f(b) - f(a)}{b - a} \leq \frac{f(c) - f(b)}{c - b}. \quad (6.1)$$

This gives

$$\frac{f(b) - f(a)}{b - a} \leq \frac{f(c) - f(a)}{c - a} \leq \frac{f(c) - f(b)}{c - b} \quad (6.2)$$

from where the theorem follows.

□




Figure 6.3: Theorem 379.

**380 THEOREM** Let  $I \subseteq \mathbb{R}$  be an interval and let  $f: I \rightarrow \mathbb{R}$  be a convex function. Then  $f$  is left and right differentiable on every point of  $\overset{\circ}{I}$  and for  $(a, b, c) \in I^3$  with  $a < b < c$ ,

$$\frac{f(b) - f(a)}{b - a} \leq f_-(b) \leq f_+(b) \leq \frac{f(c) - f(b)}{c - b}.$$

**Proof:** Since  $f$  is convex,  $\forall b \in \overset{\circ}{I}$ ,  $T_b: \begin{array}{l} I \setminus \{b\} \rightarrow \mathbb{R} \\ x \mapsto \frac{f(x) - f(b)}{x - b} \end{array}$  is increasing, by virtue of Theorem 379. Thus

$$\forall u \in [a; b], \forall v \in [b; c]$$

$$T_b(a) \leq T_b(u) \leq T_b(v) \leq T_b(c).$$

This means that  $T_b$  is increasing on  $[b; c]$  and bounded below by  $T_b(u)$ . It follows by Theorem 333 that  $T_b(b+)$  exists, and so  $f$  is right-differentiable at  $b$ . Moreover,

$$T_b(a) \leq T_b(u) \leq f'_+(b) \leq T_b(c).$$

Similarly,  $T_b$  is increasing and bounded above by  $f'_+(b)$ . Appealing again to Theorem 333,  $f$  is left-differentiable at  $b$  and

$$T_b(a) \leq f'_-(b) \leq f'_+(b) \leq T_b(c).$$

□

**381 COROLLARY** If  $f$  is convex on an interval  $I$ , then  $f$  is continuous on  $\overset{\circ}{I}$ .

**Proof:** Given  $b \in \overset{\circ}{I}$ , we know that  $f$  is both left and right differentiable at  $b$  (though we may have  $f'_-(b) < f'_+(b)$ ). Regardless, this makes  $f$  left and right continuous at  $b$ : hence both  $f(b-) = f(b)$  and  $f(b+) = f(b)$ . But then  $f(b-) = f(b+) = f(b)$  and so  $f$  is continuous at  $b$ . □

**382 THEOREM** Let  $I \subseteq \mathbb{R}$  be an interval and let  $f: I \rightarrow \mathbb{R}$  be differentiable on  $I$ . Then  $f$  is convex if and only if  $f'$  is increasing on  $I$ .

**Proof:**

⇒ Assume  $f$  is convex. Let  $a < x < c$ . By (6.2),

$$\frac{f(x) - f(a)}{x - a} \leq \frac{f(c) - f(a)}{c - a} \leq \frac{f(c) - f(x)}{c - x}.$$

Taking limits as  $x \rightarrow a+$ ,

$$f'_+(a) \leq \frac{f(c) - f(a)}{c - a}.$$

Taking limits as  $x \rightarrow c-$ ,

$$\frac{f(c) - f(a)}{c - a} \leq f'_-(c).$$

Thus  $f'_+(a) \leq f'_-(c)$ . Since  $f$  is differentiable,  $f'_+(a) = f'(a)$  and  $f'_-(c) = f'(c)$ , and so  $f'(a) \leq f'(c)$  proving that  $f'$  is increasing.

⇐ Assume  $f'$  is increasing and that  $a < x < b$ . By the Mean Value Theorem, there exists  $\alpha \in ]a; x[$  and  $\alpha' \in ]x; b[$  such that

$$\frac{f(x) - f(a)}{x - a} = f(\alpha), \quad \frac{f(b) - f(x)}{b - x} = f(\alpha').$$

Since  $f'(\alpha) \leq f'(\alpha')$  we must have

$$\frac{f(x) - f(a)}{x - a} \leq \frac{f(b) - f(x)}{b - x},$$

and so  $f$  is convex in view of (6.1).

□

**383 COROLLARY** Let  $I \subseteq \mathbb{R}$  be an interval and let  $f: I \rightarrow \mathbb{R}$  be twice differentiable on  $I$ . Then  $f$  is convex if and only if  $f'' \geq 0$ .

**Proof:** This follows from Theorems 368 and 382. □

**384 Definition** An inflexion point is a point on the graph of a function where the graph changes from convex to concave or viceversa.

## Homework

**Problem 6.5.1 (Putnam 1991)** Are there any polynomials  $p(x)$  with real coefficients of degree  $n \geq 2$  all whose  $n$  roots are distinct real numbers and all whose  $n - 1$  zeroes of  $p'(x)$  are the midpoints between consecutive roots of  $p(x)$ ?

**Problem 6.5.2** Prove that the inflexion points of  $x \mapsto \frac{x}{\tan x}$  are aligned.

**Problem 6.5.3** By considering  $f: [0; +\infty[ \rightarrow \mathbb{R}$   
 $x \mapsto x^k - k(x-1)$   
 for  $0 < k < 1$  and using first and second derivative arguments, obtain a new proof of Young's Inequality 314.

## 6.6 Inequalities Obtained Through Differentiation

**385 THEOREM** Let  $x > 0$ . Then  $\frac{x^2}{2} < \exp(x)$ .

**Proof:** Let  $f(x) = \exp(x) - \frac{x^2}{2}$ . Then  $f'(x) = \exp(x) - x$  and  $f''(x) = \exp(x) - 1$ . Since  $x > 0$ ,  $f''(x) > 0$  and so  $f'$  is strictly increasing. Thus  $f'(x) > f'(0) = 1 > 0$  and so  $f$  is increasing. Thus

$$f(x) > f(0) \implies \exp(x) - \frac{x^2}{2} > 0,$$

proving the theorem.  $\square$

**386 THEOREM**  $\lim_{x \rightarrow +\infty} \frac{x}{\exp(x)} = 0$ .

**Proof:** From Theorem 385, for  $x > 0$ ,

$$0 < \frac{x}{\exp(x)} < \frac{2}{x} \implies 0 \leq \lim_{x \rightarrow +\infty} \frac{x}{\exp(x)} \leq \lim_{x \rightarrow +\infty} \frac{2}{x} = 0,$$

and the theorem follows from the Sandwich Theorem.  $\square$

**387 THEOREM** Let  $\alpha \in \mathbb{R}$ . Then  $\lim_{x \rightarrow +\infty} \frac{x^\alpha}{\exp(x)} = 0$ .

**Proof:** If  $\alpha < 1$  then

$$\frac{x^\alpha}{\exp(x)} = \frac{x}{\exp(x)} \cdot x^{\alpha-1} \rightarrow 0 \cdot 0,$$

by Lemma 386. If  $\alpha \geq 1$  then

$$\frac{x^\alpha}{\exp(x)} = \alpha^{-\alpha} \left( \frac{\alpha x}{\exp(\alpha x)} \right)^\alpha \rightarrow \alpha^{-\alpha} \cdot 0^\alpha = 0,$$

by continuity and by Lemma 386.  $\square$

**388 THEOREM** Let  $x > 0$ . Then  $\log x < x$ .

**Proof:** Put  $f(x) = x - \log x$ . Then  $f'(x) = 1 - \frac{1}{x}$ . For  $x < 1$ ,  $f'(x) < 0$ , for  $x = 1$ ,  $f'(x) = 0$ , and for  $x > 1$ ,  $f'(x) > 0$ , which means that  $f$  has a minimum at  $x = 1$ . Thus

$$f(x) > f(1) \implies x - \log x > 1.$$

Since  $x - \log x > 1$  then a fortiori we must have  $x - \log x > 0$  and the theorem follows.  $\square$

**389 LEMMA**  $\lim_{x \rightarrow +\infty} \frac{\log x}{x} = 0$ .

**Proof:** From Theorem 388,  $\log x^2 < x^2$ . For  $x > 1$ ,  $\log x > 0$  and hence,

$$x > 1 \implies 0 < \frac{\log x}{x} < \frac{1}{2x},$$

whence  $\lim_{x \rightarrow +\infty} \frac{\log x}{x} = 0$  by the Sandwich Theorem.  $\square$

**390 THEOREM** Let  $\alpha \in ]0; +\infty[$ . Then  $\lim_{x \rightarrow +\infty} \frac{\log x}{x^\alpha} = 0$ .

**Proof:** If  $\alpha > 1$  then

$$\frac{\log x}{x^\alpha} = \frac{\log x}{x} \cdot x^{1-\alpha} \rightarrow 0 \cdot 0,$$

by Lemma 389. If  $0 < \alpha \leq 1$  then

$$\frac{\log x}{x^\alpha} = \frac{\log x^\alpha}{\alpha x^\alpha} \rightarrow \frac{1}{\alpha} \cdot 0 = 0,$$

by continuity and by Lemma 389.  $\square$

**391 THEOREM** For  $x \in ]0; \frac{\pi}{2}[$ ,  $\sin x < x < \tan x$ .

**Proof:** Observe that we gave a geometrical argument for this inequality in Theorem 344. First, let  $f(x) = \sin x - x$ . Then  $f'(x) = \cos x - 1 < 0$ , since for  $x \in ]0; \frac{\pi}{2}[$ , the cosine is strictly positive. This means that  $f$  is strictly decreasing. Thus for all  $x \in ]0; \frac{\pi}{2}[$ ,

$$f(0) > f(x) \implies 0 > \sin x - x \implies \sin x < x,$$

giving the first half of the inequality.

For the second half, put  $g(x) = \tan x - x$ . Then  $g'(x) = \sec^2 x - 1$ . Now, since  $|\cos x| < 1$  for  $x \in ]0; \frac{\pi}{2}[$ ,  $\sec^2 x > 1$ . Hence  $g'(x) > 0$ , and so  $g$  is strictly increasing. This gives

$$g(0) < g(x) \implies 0 < \tan x - x \implies x < \tan x,$$

obtaining the second inequality.  $\square$




Figure 6.4: Jordan's Inequality

**392 THEOREM (Jordan's Inequality)** For  $x \in ]0; \frac{\pi}{2}[$ ,  $\frac{2}{\pi}x < \sin x < x$ .

**Proof:** This inequality says that the straight line joining  $(0,0)$  to  $(\frac{\pi}{2},1)$  lies below the curve  $y = \sin x$  for  $x \in ]0; \frac{\pi}{2}[$ . See figure 6.4. Put  $f(x) = \frac{\sin x}{x}$  for  $x \neq 0$  and  $f(0) = 1$ . Then  $f'(x) = (\cos x) \left( \frac{x - \tan x}{x^2} \right) < 0$  since  $\cos x > 0$  and  $x - \tan x < 0$  for  $x \in ]0; \frac{\pi}{2}[$ . Thus  $f$  is strictly decreasing for  $x \in ]0; \frac{\pi}{2}[$  and so

$$f(x) > f\left(\frac{\pi}{2}\right) \implies \frac{\sin x}{x} > \frac{2}{\pi},$$

proving the theorem.  $\square$

**393 Definition** If  $w_1, w_2, \dots, w_n$  are positive real numbers such that  $w_1 + w_2 + \dots + w_n = 1$ , we define the  $r$ -th weighted power mean of the  $x_i$  as:

$$M_w^r(x_1, x_2, \dots, x_n) = (w_1 x_1^r + w_2 x_2^r + \dots + w_n x_n^r)^{1/r}.$$

When all the  $w_i = \frac{1}{n}$  we get the standard power mean. The weighted power mean is a continuous function of  $r$ , and taking limit when  $r \rightarrow 0$  gives us

$$M_w^0 = x_1^{w_1} x_2^{w_2} \dots x_n^{w_n}.$$

**394 THEOREM (Generalisation of the AM-GM Inequality)** If  $r < s$  then

$$M_w^r(x_1, x_2, \dots, x_n) \leq M_w^s(x_1, x_2, \dots, x_n).$$

**Proof:** Suppose first that  $0 < r < s$  are real numbers, and let  $w_1, w_2, \dots, w_n$  be positive real numbers such that  $w_1 + w_2 + \dots + w_n = 1$ .

Put  $t = \frac{s}{r} > 1$  and  $y_i = x_i^r$  for  $1 \leq i \leq n$ . This implies that  $y_i^t = x_i^s$ . The function  $f: ]0; +\infty[ \rightarrow ]0; +\infty[$ ,  $f(x) = x^t$  is strictly convex, since its second derivative is  $f''(x) = \frac{1}{t(t-1)} x^{t-2} > 0$  for all  $x \in ]0; +\infty[$ . By Jensen's inequality,

$$\begin{aligned} (w_1 y_1 + w_2 y_2 + \dots + w_n y_n)^t &= f(w_1 y_1 + w_2 y_2 + \dots + w_n y_n) \\ &\leq w_1 f(y_1) + w_2 f(y_2) + \dots + w_n f(y_n) \\ &= w_1 y_1^t + w_2 y_2^t + \dots + w_n y_n^t. \end{aligned}$$

with equality if and only if  $y_1 = y_2 = \dots = y_n$ . By substituting  $t = \frac{s}{r}$  and  $y_i = x_i^r$  back into this inequality, we get

$$(w_1 x_1^r + w_2 x_2^r + \dots + w_n x_n^r)^{s/r} \leq w_1 x_1^s + w_2 x_2^s + \dots + w_n x_n^s$$

with equality if and only if  $x_1 = x_2 = \dots = x_n$ . Since  $s$  is positive, the function  $x \mapsto x^{1/s}$  is strictly increasing, so raising both sides to the power  $1/s$  preserves the inequality:

$$(w_1 x_1^r + w_2 x_2^r + \dots + w_n x_n^r)^{1/r} \leq (w_1 x_1^s + w_2 x_2^s + \dots + w_n x_n^s)^{1/s},$$

which is the inequality we had to prove. Equality holds if and only if all the  $x_i$  are equal.

The cases  $r < 0 < s$  and  $r < s < 0$  can be reduced to the case  $0 < r < s$ .  $\square$

## Homework

**Problem 6.6.1** Complete the following steps (due to George Pólya) in order to prove the AM-GM Inequality (Theorem 86).

2. Put

1. Prove that  $\forall x \in \mathbb{R}, x \leq e^{x-1}$ .

$$A_k = \frac{na_k}{a_1 + a_2 + \dots + a_n},$$

and  $G_n = a_1 a_2 \cdots a_n$ . Prove that

$$A_1 A_2 \cdots A_n = \frac{n^n G_n}{(a_1 + a_2 + \cdots + a_n)^n},$$

and that

$$A_1 + A_2 + \cdots + A_n = n.$$

3. Deduce that

$$G_n \leq \left( \frac{a_1 + a_2 + \cdots + a_n}{n} \right)^n.$$

4. Prove the AM-GM inequality by assembling the results above.

## 6.7 Asymptotic Preponderance

**395 Definition** Let  $I \subseteq \overline{\mathbb{R}}$  be an interval, and let  $a \in I$ . A function  $\alpha : I \rightarrow \mathbb{R}$  is said to be *infinitesimal* as  $x \rightarrow a$  if  $\lim_{x \rightarrow a} \alpha(x) = 0$ . We say that  $\alpha$  is *negligible* in relation to  $\beta$  as  $x \rightarrow a$  or that  $\beta$  is *preponderant* in relation to  $\alpha$  as  $x \rightarrow a$ , if  $\forall \varepsilon > 0, \exists \delta > 0$  such that

$$x \in ]a - \delta; a + \delta[ \implies |\alpha(x)| \leq \varepsilon |\beta(x)|.$$

We express the condition above with the notation  $\alpha(x) = o_{x \rightarrow a}(\beta(x))$  (read “ $\alpha$  of  $x$  is small oh of  $\beta$  of  $x$  as  $x$  tends to  $a$ ”).

Finally, we say that  $\alpha$  is *Big Oh* of  $\beta$  around  $x = a$ —written  $\alpha(x) = O_{x \rightarrow a}(\beta(x))$ , or  $\alpha(x) \ll_{x \rightarrow a}(\beta(x))$ —if  $\exists C > 0$  and  $\exists \delta > 0$  such that  $\forall x \in ]a - \delta; a + \delta[$ ,  $|\alpha(x)| \leq C |\beta(x)|$ .



Notice that  $a$  above may be finite or  $\pm\infty$ . If  $a$  is understood, we prefer to write  $\alpha(x) = o(\beta(x))$  rather  $\alpha(x) = o_{x \rightarrow a}(\beta(x))$ .  
Also

$$\alpha = o_{x \rightarrow a}(\beta) \iff \lim_{x \rightarrow a} \frac{\alpha(x)}{\beta(x)} = 0 \quad \text{and} \quad \beta(a) = 0 \implies \alpha(a) = 0.$$

**396 Example**  $\sin : \mathbb{R} \rightarrow [-1; 1]$  is infinitesimal as  $x \rightarrow 0$ , since  $\lim_{x \rightarrow 0} \sin x = 0$ .

**397 Example**  $f : \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}, x \mapsto \frac{1}{x}$  is infinitesimal as  $x \rightarrow +\infty$ , since  $\lim_{x \rightarrow +\infty} \frac{1}{x} = 0$ .

**398 Example** We have  $x^2 = o(x)$  as  $x \rightarrow 0$  since

$$\lim_{x \rightarrow 0} \frac{x^2}{x} = \lim_{x \rightarrow 0} x = 0.$$

**399 Example** We have  $x = o(x^2)$  as  $x \rightarrow +\infty$  since

$$\lim_{x \rightarrow +\infty} \frac{x}{x^2} = \lim_{x \rightarrow +\infty} \frac{1}{x} = 0.$$

**400 Definition** We write  $\alpha(x) = \gamma(x) + o(\beta(x))$  as  $x \rightarrow a$  if  $\alpha(x) - \gamma(x) = o(\beta(x))$  as  $x \rightarrow a$ . Similarly,  $\alpha(x) = \gamma(x) + O(\beta(x))$  as  $x \rightarrow a$  means that  $\alpha(x) - \gamma(x) = O(\beta(x))$  as  $x \rightarrow a$ .

**401 Example** We have  $\sin x = x + o(x)$  as  $x \rightarrow 0$  since

$$\lim_{x \rightarrow 0} \frac{\sin x - x}{x} = \lim_{x \rightarrow 0} \frac{\sin x}{x} - \lim_{x \rightarrow 0} 1 = 1 - 1 = 0.$$

**402 THEOREM** Let  $f, g, \alpha, \beta, u, v$  be real-valued functions defined on an interval containing  $a \in \overline{\mathbb{R}}$ . Let  $\lambda \in \mathbb{R}$  be a constant. Let  $h$  be a real valued function defined on an interval containing  $b \in \overline{\mathbb{R}}$ . Then

1.  $f = o(g) \implies f = O(g)$ .
2.  $f = o(\alpha) \implies \lambda f = o(\alpha)$ .
3.  $f = o(\alpha), g = o(\alpha) \implies f + g = o(\alpha)$ .
4.  $f = o(\alpha), g = o(\beta) \implies fg = o(\alpha\beta)$ .

5.  $f = O(\alpha) \implies \lambda f = O(\alpha)$ .
6.  $f = O(\alpha), g = O(\alpha) \implies f + g = O(\alpha)$ .
7.  $f = O(\alpha), g = O(\beta) \implies fg = O(\alpha\beta)$ .
8.  $f = O(\alpha), g = o(\beta) \implies fg = o(\alpha\beta)$ .
9.  $f = O(\alpha), \alpha = O(\beta) \implies f = O(\beta)$ .
10.  $f = o(\alpha), \alpha = O(\beta) \implies f = o(\beta)$ .
11.  $f = O(\alpha), \alpha = o(\beta) \implies f = o(\beta)$ .
12.  $f = o(\alpha), \lim_{x \rightarrow b} h(x) = a \implies f \circ h = o_{x \rightarrow b}(\alpha \circ h)$ .
13.  $f = O(\alpha), \lim_{x \rightarrow b} h(x) = a \implies f \circ h = O_{x \rightarrow b}(\alpha \circ h)$ .

**Proof:** These statements follow directly from the definitions.

1. If  $f = o(g)$  then  $\forall \varepsilon > 0$  there exists  $\delta > 0$  such that

$$x \in ]a - \delta; a + \delta[ \implies \left| \frac{f(x)}{g(x)} - 0 \right| < \varepsilon \implies |f(x)| < \varepsilon |g(x)| \implies f = o(g),$$

using  $C = \varepsilon$  in the definition of Big Oh.

2. This follows by Theorem 272.
3. This follows by Theorem 272.

4. Both  $\lim_{x \rightarrow a} \frac{f(x)}{\alpha(x)} = 0$  and  $\lim_{x \rightarrow a} \frac{g(x)}{\beta(x)} = 0$ . Hence  $\lim_{x \rightarrow a} \frac{f(x)g(x)}{\alpha(x)\beta(x)} = \lim_{x \rightarrow a} \frac{f(x)}{\alpha(x)} \cdot \lim_{x \rightarrow a} \frac{g(x)}{\beta(x)} = 0 \implies fg = o(\alpha\beta)$ .

5. If  $f = O(\alpha)$  then there is  $\delta > 0$  and  $C > 0$  such that

$$x \in ]a - \delta; a + \delta[ \implies |f(x)| \leq C |g(x)| \implies |\lambda f(x)| \leq C |\lambda| \cdot |g(x)| \implies \lambda f = O(\alpha)$$

6. There exists  $\delta_1 > 0, \delta_2 > 0$  and  $C_1 > 0, C_2 > 0$  such that

$$x \in ]a - \delta_1; a + \delta_1[ \implies |f(x)| \leq C_1 |\alpha(x)| \quad \text{and} \quad x \in ]a - \delta_2; a + \delta_2[ \implies |g(x)| \leq C_2 |\alpha(x)|.$$

Thus if  $\delta = \min(\delta_1, \delta_2)$ ,

$$x \in ]a - \delta; a + \delta[ \implies |f(x) + g(x)| \leq |f(x)| + |g(x)| \leq C_1 \alpha(x) + C_2 \alpha(x) = (C_1 + C_2) \alpha(x) \implies f + g = O(\alpha).$$

7. There exists  $\delta_1 > 0, \delta_2 > 0$  and  $C_1 > 0, C_2 > 0$  such that

$$x \in ]a - \delta_1; a + \delta_1[ \implies |f(x)| \leq C_1 |\alpha(x)| \quad \text{and} \quad x \in ]a - \delta_2; a + \delta_2[ \implies |g(x)| \leq C_2 |\beta(x)|.$$

Thus if  $\delta = \min(\delta_1, \delta_2)$ ,

$$x \in ]a - \delta; a + \delta[ \implies |f(x)g(x)| = |f(x)| |g(x)| \leq C_1 |\alpha(x)| \cdot C_2 |\beta(x)| = (C_1 C_2) |\alpha(x)\beta(x)| \implies fg = O(\alpha\beta).$$

8. There exists  $\delta_1 > 0, \delta_2 > 0$  and  $C_1 > 0$ , such that  $\forall \varepsilon > 0$

$$x \in ]a - \delta_1; a + \delta_1[ \implies |f(x)| \leq C_1 |\alpha(x)| \quad \text{and} \quad x \in ]a - \delta_2; a + \delta_2[ \implies |g(x)| \leq \varepsilon |\beta(x)|.$$

Thus if  $\delta = \min(\delta_1, \delta_2)$ ,

$$x \in ]a - \delta; a + \delta[ \implies |f(x)g(x)| = |f(x)| |g(x)| \leq C_1 |\alpha(x)| \cdot \varepsilon |\beta(x)| = \varepsilon (C_1) |\alpha(x)\beta(x)| \implies fg = o(\alpha\beta).$$

9. There exists  $\delta_1 > 0, \delta_2 > 0$  and  $C_1 > 0, C_2 > 0$  such that

$$x \in ]a - \delta_1; a + \delta_1[ \implies |f(x)| \leq C_1 |\alpha(x)| \quad \text{and} \quad x \in ]a - \delta_2; a + \delta_2[ \implies |\alpha(x)| \leq C_2 |\beta(x)|.$$

Thus if  $\delta = \min(\delta_1, \delta_2)$ ,

$$x \in ]a - \delta; a + \delta[ \implies |f(x)| \leq C_1 |\alpha(x)| \leq C_1 C_2 |\beta(x)| \implies f = O(\beta).$$

10. There exists  $\delta_1 > 0, \delta_2 > 0$  and  $C > 0$ , such that  $\forall \varepsilon > 0$

$$x \in ]a - \delta_1; a + \delta_1[ \implies |f(x)| \leq \varepsilon |\alpha(x)| \quad \text{and} \quad x \in ]a - \delta_2; a + \delta_2[ \implies |\alpha(x)| \leq C |\beta(x)|.$$

Thus if  $\delta = \min(\delta_1, \delta_2)$ ,

$$x \in ]a - \delta; a + \delta[ \implies |f(x)| \leq \varepsilon |\alpha(x)| \leq C\varepsilon |\beta(x)| \implies f = o(\beta).$$

11. There exists  $\delta_1 > 0, \delta_2 > 0$  and  $C > 0$ , such that  $\forall \varepsilon > 0$

$$x \in ]a - \delta_1; a + \delta_1[ \implies |f(x)| \leq C |\alpha(x)| \quad \text{and} \quad x \in ]a - \delta_2; a + \delta_2[ \implies |\alpha(x)| \leq \varepsilon |\beta(x)|.$$

Thus if  $\delta = \min(\delta_1, \delta_2)$ ,

$$x \in ]a - \delta; a + \delta[ \implies |f(x)| \leq C |\alpha(x)| \leq C\varepsilon |\beta(x)| \implies f = o(\beta).$$

12. There exists  $\delta_1 > 0, \delta_2 > 0$  such that  $\forall \varepsilon > 0$

$$x \in ]a - \delta_1; a + \delta_1[ \implies |f(x)| \leq \varepsilon |\alpha(x)| \quad \text{and} \quad x \in ]b - \delta_2; b + \delta_2[ \implies |h(x) - a| \leq \varepsilon \implies h(x) \in ]a - \varepsilon; a + \varepsilon[.$$

Thus if  $\delta = \min(\delta_1, \delta_2, \varepsilon)$ ,

$$x \in ]b - \delta; b + \delta[ \implies |(f \circ h)(x)| \leq \varepsilon |(\alpha \circ h)(x)| \implies f \circ h = o_{x \rightarrow b}(\alpha \circ h).$$

13. There exists  $\delta_1 > 0, \delta_2 > 0, C > 0$  such that  $\forall \varepsilon > 0$

$$x \in ]a - \delta_1; a + \delta_1[ \implies |f(x)| \leq C |\alpha(x)| \quad \text{and} \quad x \in ]b - \delta_2; b + \delta_2[ \implies |h(x) - a| \leq \varepsilon \implies h(x) \in ]a - \varepsilon; a + \varepsilon[.$$

Thus if  $\delta = \min(\delta_1, \delta_2, \varepsilon)$ ,

$$x \in ]b - \delta; b + \delta[ \implies |(f \circ h)(x)| \leq C |(\alpha \circ h)(x)| \implies f \circ h = O_{x \rightarrow b}(\alpha \circ h).$$

□



In the above theorem, (8), (10), and (11) essentially say that  $O(o) = o(O) = o(o) = o$  and (9) says that  $O(O) = O$ .

The following corollary is immediate.

**403 COROLLARY** Let  $\alpha$  and  $\beta$  be infinitesimal functions as  $x \rightarrow a$ . Then the following hold.

1. The sum of two infinitesimals is an infinitesimal:

$$o(\beta(x)) + o(\beta(x)) = o(\beta(x)).$$

2. The difference of two infinitesimals is an infinitesimal:

$$o(\beta(x)) - o(\beta(x)) = o(\beta(x)).$$

3.  $\forall c \in \mathbb{R} \setminus \{0\}, o(c\beta(x)) = o(\beta(x))$ .

4.  $\forall n \in \mathbb{N}, n \geq 2, 1 \leq k \leq n - 1, o((\beta(x))^n) = o((\beta(x))^k)$ .



5.  $o(o(\beta(x))) = o(\beta(x))$ .
6.  $\forall n \in \mathbb{N}, n \geq 1, (\beta(x))^n o(\beta(x)) = o((\beta(x))^{n+1})$ .
7.  $\forall n \in \mathbb{N}, n \geq 2, \frac{o((\beta(x))^n)}{\beta(x)} = o((\beta(x))^{n-1})$ .
8.  $\frac{o(\beta(x))}{\beta(x)} = o(1)$ .
9. If  $c_k$  are real numbers, then  $o\left(\sum_{k=1}^n c_k (\beta(x))^k\right) = o(\beta(x))$ .
10.  $(\alpha\beta)(x) = o(\alpha(x))$  and  $(\alpha\beta)(x) = o(\beta(x))$ .
11. If  $\alpha \sim \beta$ , then  $(\alpha - \beta)(x) = o(\alpha(x))$  and  $(\alpha - \beta)(x) = o(\beta(x))$ .

**404 THEOREM (Canonical small oh Relations)** The following relationships hold

1.  $\forall (\alpha, \beta) \in \mathbb{R}^2, x^\alpha = o_{x \rightarrow +\infty}(x^\beta) \iff \alpha < \beta$ .
2.  $\forall (\alpha, \beta) \in \mathbb{R}^2, x^\alpha = o_{x \rightarrow 0^+}(x^\beta) \iff \alpha > \beta$ .
3.  $\log x = o_{x \rightarrow +\infty}(x)$ .
4.  $\forall (\alpha, \beta) \in \mathbb{R}^2, \beta > 0, (\log x)^\alpha = o_{x \rightarrow +\infty}(x^\beta)$ .
5.  $\forall (\alpha, \beta) \in \mathbb{R}^2, \beta < 0, |\log x|^\alpha = o_{x \rightarrow 0^+}(x^\beta)$ .
6.  $\forall (\alpha, a) \in \mathbb{R}^2, a > 1, x^\alpha = o_{x \rightarrow +\infty}(a^x)$
7.  $\forall (\alpha, a) \in \mathbb{R}^2, a > 1, a^x = o_{x \rightarrow -\infty}(|x|^\alpha)$

**Proof:**

1. Immediate.
2. Immediate.
3. This follows from Lemma 389.
4. If  $\alpha = 0$  then eventually  $(\log x)^\alpha = 1$  and so the assertion is immediate. If  $\alpha < 0$  the assertion is also immediate, since then  $(\log x)^\alpha \rightarrow 0$  as  $x \rightarrow +\infty$ . If  $\alpha > 0$ , by Theorem 390,

$$\frac{\log x}{x^{\beta/\alpha}} \rightarrow 0,$$

whence

$$\frac{(\log x)^\alpha}{x^\beta} = \left(\frac{\log x}{x^{\beta/\alpha}}\right)^\alpha \rightarrow 0^\alpha = 0.$$

5. If  $x \rightarrow 0^+$  then  $\frac{1}{x} \rightarrow +\infty$ . Hence by the preceding part and by continuity, as  $x \rightarrow 0^+$  and for  $\gamma > 0$ ,

$$\frac{\left(\log \frac{1}{x}\right)^\alpha}{\left(\frac{1}{x}\right)^\gamma} \rightarrow 0.$$

But

$$\frac{\left(\log \frac{1}{x}\right)^\alpha}{\left(\frac{1}{x}\right)^\gamma} = \frac{(|-\log x|)^\alpha}{\left(\frac{1}{x}\right)^\gamma} = x^\gamma |\log x|^\alpha,$$

and so  $|\log x|^\alpha = o_{x \rightarrow 0^+}(x^{-\gamma})$ , and so putting  $\beta = -\gamma < 0$  we have  $|\log x|^\alpha = o_{x \rightarrow 0^+}(x^\beta)$ .

6. For  $\alpha < 1$  we have

$$\frac{x^\alpha}{a^x} = \frac{x \log a}{\exp(x \log a)} \cdot \frac{x^{\alpha-1}}{\log a} \rightarrow 0 \cdot 0,$$

since  $\frac{x \log a}{\exp(x \log a)} \rightarrow 0$  by continuity and Theorem 387, and  $\frac{x^{\alpha-1}}{\log a} \rightarrow 0$  since  $\alpha - 1 < 0$ . If  $\alpha > 1$  then

$$\frac{x^\alpha}{a^x} = \left( \frac{x}{(a^{1/\alpha})^x} \right)^\alpha = \frac{\alpha^\alpha}{(\log a)^\alpha} \cdot \left( \frac{x \frac{\log a}{\alpha}}{\exp\left(x \frac{\log a}{\alpha}\right)} \right)^\alpha \rightarrow \frac{\alpha^\alpha}{(\log a)^\alpha} \cdot 0^\alpha = 0,$$

by continuity and Theorem 387.

7. If  $\alpha > 0$ ,  $a > 1$  then  $|x|^\alpha \rightarrow +\infty$  but  $a^x \rightarrow 0$  as  $x \rightarrow -\infty$ , hence there is nothing to prove. If  $\alpha = 0$ , again the result is obvious. Assume  $\alpha < 0$ . If  $x \rightarrow -\infty$  then  $-x \rightarrow +\infty$  and so by the preceding part

$$\frac{|x|^{-\alpha}}{a^{-x}} \rightarrow 0$$

since the above result is valid regardless of the sign of  $\alpha$ . Now

$$\frac{a^x}{|x|^\alpha} = \frac{|x|^{-\alpha}}{a^{-x}},$$

proving the result.

□

**405 Example** In view of Corollary 403 and Theorem 404, we have

$$o(-2x^3 + 8x^2) = o(x),$$

as  $x \rightarrow 0$ .

**406 Example** In view of Corollary 403 and Theorem 404, we have

$$o(-2x^3 + 8x^2) = o(x^4),$$


as  $x \rightarrow +\infty$ .

## Homework

**Problem 6.7.1** Which one is faster as  $x \rightarrow +\infty$ ,  $(\log \log x)^{\log x}$  or  $(\log x)^{\log \log x}$ ?

## 6.8 Asymptotic Equivalence

**407 Definition** Let  $I \subseteq \overline{\mathbb{R}}$  be an interval, and let  $a \in I$ . We say that  $\alpha$  is asymptotic to a function  $\beta: I \rightarrow \mathbb{R}$  as  $x \rightarrow a$ , and we write  $\alpha \sim \beta$ , if  $\alpha \sim \beta \iff \alpha - \beta = o_a(\beta)$ .

 If in a neighbourhood  $\mathcal{N}_a$  of a  $\beta \neq 0$  then

$$\alpha \sim \beta \iff \begin{cases} \frac{\alpha}{\beta} \sim 1 \\ \beta(a) = 0 \implies \alpha(a) = 0 \end{cases}$$

**408 Example** We have  $\sin x \sim x$  as  $x \rightarrow 0$ , since  $\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$ .

**409 Example** We have  $x^2 + x \sim x$  as  $x \rightarrow 0$ , since  $\lim_{x \rightarrow 0} \frac{x^2 + x}{x} = 1$ .

**410 Example** We have  $x^2 + x \sim x^2$  as  $x \rightarrow +\infty$ , since  $\lim_{x \rightarrow +\infty} \frac{x^2 + x}{x^2} = 1$ .

**411 THEOREM**

$$\alpha \sim \beta \implies \begin{cases} \alpha = O(\beta) \\ \beta = O(\alpha) \end{cases}$$

**Proof:** If  $\alpha - \beta = o(\beta)$  there is a neighbourhood  $\mathcal{N}_a$  of  $a$  such that

$$\forall \varepsilon > 0, x \in \mathcal{N}_a \implies |\alpha(x) - \beta(x)| \leq \varepsilon |\beta(x)|.$$

In particular, for  $\varepsilon = \frac{1}{2}$ , we have

$$x \in \mathcal{N}_a \implies |\alpha(x) - \beta(x)| \leq \frac{1}{2} |\beta(x)|.$$

Hence

$$x \in \mathcal{N}_a \implies |\alpha(x)| = |\alpha(x) - \beta(x) + \beta(x)| \leq |\alpha(x) - \beta(x)| + |\beta(x)| \leq \frac{3}{2} |\beta(x)| \implies \alpha = O(\beta),$$

and

$$x \in \mathcal{N}_a \implies |\beta(x)| = |\beta(x) - \alpha(x) + \alpha(x)| \leq |\beta(x) - \alpha(x)| + |\alpha(x)| \leq \frac{1}{2} |\beta(x)| + |\alpha(x)| \implies |\beta(x)| \leq 2|\alpha(x)| \implies \beta = O(\alpha).$$

□

**412 THEOREM** The relation of asymptotic equivalence  $\sim$  is an equivalence relation on the set of functions defined on a neighbourhood of  $a$ .

**Proof:** We have

**Reflexivity**  $\alpha - \alpha = 0 = o(\alpha)$ .

**Symmetry**  $\alpha - \beta = o(\beta) \implies \beta = O(\alpha)$  by Theorem 411. Now by (10) of Theorem 402,

$$\alpha - \beta = o(\beta) \quad \text{and} \quad \beta = O(\alpha) \implies \alpha - \beta = o(\alpha) \implies \beta - \alpha = o(\alpha),$$

whence  $\beta \sim \alpha$ .

**Transitivity** Assume  $\alpha - \beta = o(\beta)$  and  $\beta - \gamma = o(\gamma)$ . Then by Theorem 411 we also have  $\beta = O(\gamma)$ . Hence  $\alpha - \beta = o(\gamma)$  by (10) of Theorem 402. Finally  $\alpha - \beta = o(\gamma)$  and  $\beta - \gamma = o(\gamma)$  give  $\alpha - \gamma = o(\gamma)$  by (3) of Theorem 402.

□

The relationship between  $o$ ,  $O$ , and  $\sim$  is displayed in figure 6.5.

**413 THEOREM** The relation of asymptotic equivalence  $\sim$  possesses the following properties.

$$1. \begin{cases} \alpha \sim \beta \\ \gamma \sim \delta \end{cases} \implies \alpha\gamma \sim \beta\delta.$$




Figure 6.5: Diagram of Big Oh relations.

$$2. \begin{cases} \alpha \sim \beta \\ n \in \mathbb{N} \setminus \{0\} \end{cases} \implies \alpha^n \sim \beta^n$$

3. if  $\alpha \sim \beta$  and if there is a neighbourhood  $\mathcal{N}_a$  of  $a$  where  $\forall x \in \mathcal{N}_a \setminus \{a\}, \beta(x) \neq 0$ , then  $\frac{1}{\alpha}$  and  $\frac{1}{\beta}$  are defined on  $\mathcal{N}_a \setminus \{a\}$  and  $\frac{1}{\alpha} \sim_a \frac{1}{\beta}$ .

$$4. \begin{cases} \alpha = o(\beta) \\ \beta \sim \gamma \end{cases} \implies \alpha = o(\gamma).$$

$$5. \begin{cases} \alpha \sim \beta \\ \beta = o(\gamma) \end{cases} \implies \alpha = o(\gamma).$$

6. if  $\alpha \sim \beta$  and if there is a neighbourhood  $\mathcal{N}_a$  of  $a$  where  $\forall x \in \mathcal{N}_a \setminus \{a\}, \beta(x) > 0$ , and if  $r \in \mathbb{R}$  then  $\alpha^r \sim_a \beta^r$ .

7. (**Dextral Composition**) If  $\alpha \sim_a \beta$  and if  $\lim_{x \rightarrow b} \gamma(x) = a$ , then  $\alpha \circ \gamma \sim_a \beta \circ \gamma$ .

**Proof:** We prove the assertions in the given order.

1. Since  $\alpha - \beta = o(\beta)$  and  $\gamma - \delta = o(\delta)$  then  $\alpha = O(\beta)$ , and so

$$\alpha\gamma - \beta\delta = \alpha(\gamma - \delta) - \delta(\beta - \alpha) = O(\beta) o(\delta) - \delta o(\beta) = o(\beta\delta).$$

2. This follows upon applying the preceding product rule  $n - 1$  times, using  $\gamma = \alpha$  and  $\delta = \beta$ .

3. Observe that

$$\frac{1}{\alpha} - \frac{1}{\beta} = \frac{\beta - \alpha}{\alpha\beta} = \frac{o(\alpha)}{\alpha\beta} = o\left(\frac{1}{\beta}\right),$$

upon using  $\beta - \alpha = o(\alpha)$  and (8) of Corollary 403.

4. We have  $\alpha = o(\beta)$  and  $\beta - \gamma = o(\gamma)$ . This last implies that  $\beta = O(\gamma)$  by Theorem 411. Hence

$$\alpha = o(\beta) = o(O(\gamma)) = o(\gamma).$$

5. We have  $\alpha - \beta = o(\beta)$  and  $\beta = o(\gamma)$ . This last implies that  $\alpha = O(\beta)$  by Theorem 411. Hence

$$\alpha = O(\beta) = O(o(\gamma)) = o(\gamma).$$

6. Since  $\beta$  is eventually strictly positive, so is  $\alpha$ . Hence  $\alpha \sim \beta \iff \frac{\alpha}{\beta}(x) \rightarrow 1$  as  $x \rightarrow a$ . Since the function

$x \mapsto x^r$  is continuous in  $]0; +\infty[$ ,

$$\frac{\alpha}{\beta}(x) \rightarrow 1 \implies \frac{\alpha^r}{\beta^r}(x) \rightarrow 1 \implies \alpha^r \sim \beta^r.$$

7. We have  $\frac{\alpha(x) - \beta(x)}{\beta(x)} \rightarrow 0$  as  $x \rightarrow a$ . Now if  $\gamma(x) \rightarrow a$  as  $x \rightarrow b$  then as  $x \rightarrow b$ ,

$$\frac{\alpha(\gamma(x)) - \beta(\gamma(x))}{\beta(\gamma(x))} \rightarrow 0.$$


□

**414 THEOREM (Exponential Composition)**  $\exp(\alpha) \sim_a \exp(\beta) \iff \alpha - \beta \sim_a 0$ .

**Proof:** We have

$$\begin{aligned} \exp(\alpha) \sim_a \exp(\beta) &\iff \exp(\alpha) - \exp(\beta) = o(\exp(\beta)) \\ &\iff (\exp(-\beta))(\exp(\alpha) - \exp(\beta)) = (\exp(-\beta))o(\exp(\beta)) \\ &\iff \exp(\alpha - \beta) - 1 = o(1) \\ &\iff \alpha - \beta = o(0). \end{aligned}$$

□

 The above theorem does not say that  $\alpha \sim \beta \implies \exp(\alpha) \sim \exp(\beta)$ . That this last assertion is false can be seen from the following counterexample:  $x + 1 \sim x$  as  $x \rightarrow 0$ , but  $\exp(x + 1) = e \exp(x)$  is not asymptotic to  $\exp(x)$ .

**415 THEOREM (Logarithmic Composition)** Suppose there is a neighbourhood of  $a \in \mathcal{N}_a$  such that

$\forall x \in \mathcal{N}_a \setminus \{a\}, \beta(x) > 0$ . Suppose, moreover, that  $\alpha \sim_a \beta$  and that  $\lim_{x \rightarrow a} \beta(x) = l$  with  $l \in [0; +\infty] \setminus \{1\}$ . Then  $\log \circ \alpha \sim_a \log \circ \beta$ .

**Proof:** Either  $l \in ]0; +\infty[ \setminus \{1\}$  or  $l = +\infty$  or  $l = 0$ .

In the first case,  $\log \alpha(x) \rightarrow \log l$  and  $\log \beta(x) \rightarrow \log l$  as  $x \rightarrow a$  hence

$$\log \alpha \sim \log l \sim \log \beta, \quad \text{as } x \rightarrow a.$$

In the second case  $\beta(x) > 1$  eventually, and thus  $\log \beta(x) \neq 0$ . Hence

$$\frac{\log \alpha(x)}{\log \beta(x)} - 1 = \frac{\log \alpha(x) - \log \beta(x)}{\log \beta(x)} = \frac{\log \frac{\alpha(x)}{\beta(x)}}{\log \beta(x)} \rightarrow \frac{\log 1}{+\infty} = 0,$$

since  $\frac{\alpha(x)}{\beta(x)} \rightarrow 1$  and  $\log \beta(x) \rightarrow +\infty$  as  $x \rightarrow +\infty$ .

The third case becomes the second case upon considering  $\frac{1}{\alpha}$  and  $\frac{1}{\beta}$ . □

**416 THEOREM (Addition of Positive Terms)** If  $\alpha \sim \beta$  and  $\gamma \sim \delta$  and there exists a neighbourhood of  $a \in \mathcal{N}_a$  such that  $\forall x \in \mathcal{N}_a \setminus \{a\}, \beta(x) > 0, \delta(x) > 0$  then

$$\alpha + \gamma \sim \beta + \delta.$$

**Proof:** We have  $\alpha - \beta = o(\beta)$  and  $\gamma - \delta = o(\delta)$ . Hence

$$\begin{aligned}(\alpha + \gamma) - (\beta + \delta) &= (\alpha - \beta) + (\gamma - \delta) \\ &= o(\beta) + o(\delta) \\ &= o(\beta + \delta),\end{aligned}$$

which means  $\alpha + \gamma \sim \beta + \delta$ .  $\square$

**417 THEOREM** The following asymptotic expansions hold as  $x \rightarrow 0$ :

1.  $\exp(x) - 1 \sim x$  and thus  $\exp(x) = 1 + x + o(x)$
2.  $\log(1 + x) \sim x$  and thus  $\log(1 + x) = x + o(x)$
3.  $\sin x \sim x$  and thus  $\sin(x) = x + o(x)$
4.  $\tan x \sim x$  and thus  $\tan(x) = x + o(x)$
5.  $\arcsin x \sim x$  and thus  $\arcsin(x) = x + o(x)$
6.  $\arctan x \sim x$  and thus  $\tan(x) = x + o(x)$
7. for  $\alpha \in \mathbb{R}$  constant,  $(1 + x)^\alpha - 1 \sim \alpha x$  and thus  $(1 + x)^\alpha = 1 + \alpha x + o(x)$
8.  $1 - \cos x \sim \frac{x^2}{2}$  and thus  $\cos(x) = 1 - \frac{x^2}{2} + o(x^2)$

**Proof:** Results 1—7 follow from the fact that

$$f'(a) \neq 0, \quad \frac{f(x) - f(a)}{x - a} \rightarrow f'(a) \implies f(x) - f(a) \sim f'(a)(x - a).$$

Property 8 follows from the identity  $1 - \cos x = 2 \sin^2 \frac{x}{2}$ .  $\square$

**418 Example** Since  $\tan x = x + o(x)$ , we have

$$\tan \frac{x^2}{2} = \frac{x^2}{2} + o\left(\frac{x^2}{2}\right) = \frac{x^2}{2} + o(x^2),$$

as  $x \rightarrow 0$ . Also,

$$(\tan x)^3 = (x + o(x))^3 = x^3 + 3x^2 o(x) + 3x o(x^2) + (o(x))^3 = x^3 + o(x^3).$$

**419 Example** Since  $\cos x = 1 - \frac{x^2}{2} + o(x^2)$ , we have

$$\cos 3x^2 = 1 - \frac{9x^4}{2} + o(x^4).$$

**420 Example** Find an asymptotic expansion of  $\cot^2 x$  of type  $o(x^{-2})$  as  $x \rightarrow 0$ .

**Solution:** Since  $\tan x \sim x$  we have

$$\cot^2 x \sim \frac{1}{x^2}.$$

We can write this as  $\cot^2 x = \frac{1}{x^2} + o\left(\frac{1}{x^2}\right)$ .

421 Example Calculate

$$\lim_{x \rightarrow 0} \frac{\sin \sin \tan \frac{x^2}{2}}{\log \cos 3x}.$$

**Solution:** We use theorems 417 and 403.

$$\begin{aligned} \sin \sin \tan \frac{x^2}{2} &= \sin \sin \left( \frac{x^2}{2} + o(x^2) \right) \\ &= \sin \left( \frac{x^2}{2} + o(x^2) + o \left( \frac{x^2}{2} + o(x^2) \right) \right) \\ &= \sin \left( \frac{x^2}{2} + o(x^2) \right) \\ &= \frac{x^2}{2} + o(x^2), \end{aligned}$$

and

$$\begin{aligned} \log \cos 3x &= \log \left( 1 - \frac{9x^2}{2} + o(x^2) \right) \\ &= -\frac{9x^2}{2} + o(x^2) + o \left( -\frac{9x^2}{2} + o(x^2) \right) \\ &= -\frac{9x^2}{2} + o(x^2) \end{aligned}$$

The limit is thus equal to

$$\lim_{x \rightarrow 0} \frac{\frac{x^2}{2} + o(x^2)}{-\frac{9x^2}{2} + o(x^2)} = \lim_{x \rightarrow 0} \frac{\frac{1}{2} + o(1)}{-\frac{9}{2} + o(1)} = -\frac{1}{9}.$$

422 Example Find  $\lim_{x \rightarrow 0} (\cos x)^{(\cot^2 x)}$ .

**Solution:** By example 420, we have  $\cot^2 x = \frac{1}{x^2} + o\left(\frac{1}{x^2}\right)$ . Also,

$$\log \cos x = \log \left( 1 - \frac{x^2}{2} + o(x^2) \right) = -\frac{x^2}{2} + o(x^2).$$

Hence

$$\begin{aligned} (\cos x)^{\cot^2 x} &= \exp \left( (\cot^2 x) \log \cos x \right) \\ &= \exp \left( \left( \frac{1}{x^2} + o\left(\frac{1}{x^2}\right) \right) \left( -\frac{x^2}{2} + o(x^2) \right) \right) \\ &= \exp \left( -\frac{1}{2} + o(1) \right) \\ &\rightarrow e^{-1/2}, \end{aligned}$$

as  $x \rightarrow 0$ .

## Homework

**Problem 6.8.1** Prove that  $\frac{\log(1 + 2 \tan x)}{\sin x} \rightarrow 2$  as  $x \rightarrow 0$ .

**Problem 6.8.2** Prove that  $\left(1 + \frac{1}{x}\right)^x \rightarrow e$  as  $x \rightarrow +\infty$ .

**Problem 6.8.3** Prove that  $(\tan x)^{\cot 4x} \rightarrow e^{1/2}$  as  $x \rightarrow \frac{\pi}{4}$ .

## 6.9 Asymptotic Expansions

**423 Definition** Let  $n \in \mathbb{N}$  and let  $f: \mathcal{N}_0 \rightarrow \mathbb{R}$  where  $\mathcal{N}_0$  is a neighbourhood of  $\mathbf{0}$ . We say that  $f$  admits an *asymptotic expansion* of order  $n$  about  $x = \mathbf{0}$  if there exists a polynomial  $p$  of degree  $n$  such that

$$\forall x \in \mathcal{N}_0, \quad f(x) = p(x) + o_0(x^n).$$

The polynomial  $p$  is called the *regular part of the asymptotic expansion about  $x = \mathbf{0}$  of  $f$* .

**424 THEOREM** If  $f$  admits an asymptotic expansion about  $\mathbf{0}$ , its regular part is unique.

**Proof:** Assume  $f(x) = p(x) + o_0(x^n)$  and  $f(x) = q(x) + o_0(x^n)$ , where  $p(x) = p_n x^n + \dots + p_1 x + p_0$  and  $q(x) = q_n x^n + \dots + q_1 x + q_0$  are polynomials of degree  $n$ . If  $p \neq q$  let  $k$  be the largest  $k$  for which  $p_k \neq q_k$ . Then subtracting both equivalencies, as  $x \rightarrow \mathbf{0}$ ,

$$p(x) - q(x) = o(x^n) \implies (p_n - q_n)x^n + (p_{n-1} - q_{n-1})x^{n-1} + \dots + (p_1 - q_1)x = o(x^n) \implies (p_k - q_k)x^k + \dots = o(x^n).$$

But  $(p_k - q_k)x^k + \dots = O(x^k)$  as  $x \rightarrow \mathbf{0}$ , a contradiction, since  $k \leq n$ .  $\square$

**425 Definition** Let  $n \in \mathbb{N}$ ,  $a \in \mathbb{R}$ , and let  $f: \mathcal{N}_a \rightarrow \mathbb{R}$  where  $\mathcal{N}_a$  is a neighbourhood of  $a$ . We say that  $f$  admits an *asymptotic expansion* of order  $n$  about  $x = a$  if there exists a polynomial  $p$  of degree  $n$  such that

$$\forall x \in \mathcal{N}_a, \quad f(x) = p(x - a) + o_a((x - a)^n).$$

The polynomial  $p$  is called the *regular part of the asymptotic expansion about  $x = a$  of  $f$* .

**426 Definition** Let  $n \in \mathbb{N}$ , and let  $f: \mathcal{N}_{+\infty} \rightarrow \mathbb{R}$  where  $\mathcal{N}_{+\infty}$  is a neighbourhood of  $+\infty$ . We say that  $f$  admits an *asymptotic expansion* of order  $n$  about  $+\infty$  if there exists a polynomial  $p$  of degree  $n$  such that

$$\forall x \in \mathcal{N}_a \cap ]\mathbf{0}; +\infty[, \quad f(x) = p\left(\frac{1}{x}\right) + o_{+\infty}\left(\frac{1}{x^n}\right).$$

The polynomial  $p$  is called the *regular part of the asymptotic expansion about  $+\infty$  of  $f$* .

**427 THEOREM** Let  $f: \mathcal{N}_0 \rightarrow \mathbb{R}$  be a function with an asymptotic expansion  $f(x) = p(x) + o_0(x^n)$ , where  $p$  is a polynomial. Then, if  $f$  is even, then  $p$  is even and if  $f$  is odd, then  $p$  is odd.

**Proof:** Let  $f(x) = p(x) + o(x^n)$  as  $x \rightarrow \mathbf{0}$ , where  $p$  is a polynomial of degree  $n$ . Then  $f(-x) = p(-x) + o(x^n)$ . If  $f$  is even then

$$p(x) + o(x^n) = f(x) = f(-x) = p(-x) + o(x^n),$$

and so by uniqueness of the regular part of an asymptotic expansion we must have  $p(x) = p(-x)$ , so  $p$  is even. Similarly if  $f$  is odd then

$$-p(x) + o(x^n) = -f(x) = f(-x) = p(-x) + o(x^n),$$

and so by uniqueness of the regular part of an asymptotic expansion we must have  $-p(x) = p(-x)$ , so  $p$  is odd.

$\square$

We want to expand the function  $f$  in powers of  $x - a$ :

$$f(x) = a_0 + a_1(x - a) + a_2(x - a)^2 + \dots + a_n(x - a)^n + \dots,$$

and that we will truncate at the  $n$ -th term, obtaining thereby a polynomial of degree  $n$  in powers of  $x - a$ . We must determine what the coefficients  $a_k$  are, and what the remainder

$$f(x) - a_0 - a_1(x - a) - a_2(x - a)^2 - \dots - a_n(x - a)^n = R(x)$$



is. We hope that this remainder is  $o_a((x-a)^n)$ . The coefficients  $a_k$  are easily found. For  $0 \leq k \leq n$  since  $f$  is  $n+1$  times differentiable, differentiating  $k$  times,

$$f^{(k)}(x) = k!a_k + ((k+1)(k) \cdots 2)a_{k+1}(x-a) + ((k+2)(k+1) \cdots 3)a_{k+2}(x-a)^2 + \cdots + R^{(k)}(x), \implies \frac{f^{(k)}(a)}{k!} = a_k,$$

as long as  $R(a) = R'(a) = R''(a) = \cdots = R^{(n)}(a) = 0$ . We write our ideas formally in the following theorems.

**428 THEOREM (Taylor-Lagrange Theorem)** Let  $I \subseteq \mathbb{R}$ ,  $I \neq \emptyset$  be an interval of  $\mathbb{R}$  and let  $f: I \rightarrow \mathbb{R}$  be  $n+1$  times differentiable in  $I$ . Then if  $(x, a) \in I^2$ , there exist  $c$  with  $\inf(x, a) < c < \sup(x, a)$  such that

$$f(x) = f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \frac{f^{(3)}(a)}{3!}(x-a)^3 + \cdots + \frac{f^{(n)}(a)}{n!}(x-a)^n + \frac{f^{(n+1)}(c)}{(n+1)!}(x-a)^{n+1}.$$

**Proof:** If  $x = a$  then there is nothing to prove. If  $x < a$  then replace  $x \rightarrow f(x)$  with  $x \rightarrow f(-x)$ , which then verifies the same hypotheses given in the theorem. Thus it remains to prove the theorem for  $x > a$ . Consider the function  $\phi: [a; x] \rightarrow \mathbb{R}$  with

$$\phi(t) = f(x) - \sum_{k=0}^n f^{(k)}(t) \frac{(x-t)^k}{k!} - R \frac{(x-t)^{n+1}}{(n+1)!},$$

where  $R$  is a constant. Observe that  $\phi(x) = 0$ . We now choose the constant  $R$  so that  $\phi(a) = 0$ . Observe that  $\phi$  is differentiable and that it satisfies the hypotheses of Rolle's Theorem on  $[a; x]$ . Therefore, there exists  $c \in ]a; x[$  such that  $\phi'(c) = 0$ . Now

$$\phi'(t) = - \sum_{k=1}^n \left( f^{(k+1)}(t) \frac{(x-t)^k}{k!} - f^{(k)}(t) \frac{(x-t)^{k-1}}{(k-1)!} \right) + R \frac{(x-t)^n}{n!} = - \frac{(x-t)^n}{n!} f^{(n+1)}(t) + R \frac{(x-t)^n}{n!},$$

from where we gather, that  $R = f^{(n+1)}(c)$  and the theorem follows.  $\square$

**429 COROLLARY (Taylor-Young Theorem)** Let  $f: \mathcal{N}_a \rightarrow \mathbb{R}$  be  $n+1$  times differentiable in  $\mathcal{N}_a$ . Then  $f$  admits the asymptotic expansion of order  $n$  about  $a$ :

$$f(x) = f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \frac{f^{(3)}(a)}{3!}(x-a)^3 + \cdots + \frac{f^{(n)}(a)}{n!}(x-a)^n + o_a((x-a)^n).$$

**Proof:** Follows at once from Theorem 428.  $\square$

The following theorem follows at once from Corollary 429.

**430 THEOREM** Let  $x \rightarrow 0$ . Then

$$1. \sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \cdots + (-1)^n \frac{x^{2n+1}}{(2n+1)!} + o(x^{2n+2}).$$

$$2. \cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \cdots + (-1)^n \frac{x^{2n}}{(2n)!} + o(x^{2n+1}).$$

$$3. \tan x = x + \frac{x^3}{3} + \frac{2x^5}{15} + o(x^5).$$

$$4. e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots + \frac{x^n}{n!} + o(x^n)$$

$$5. \log(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \cdots + (-1)^{n+1} \frac{x^n}{n} + o(x^n).$$

$$6. (1+x)^\tau = 1 + \tau x + \frac{\tau(\tau-1)}{2}x^2 + \dots + \frac{\tau(\tau-1)(\tau-2)(\tau-3)\dots(\tau-n+1)}{n!}x^n + o(x^n).$$

**431 Example** Find an asymptotic development of  $\log(2\cos x + \sin x)$  around  $x = 0$  of order  $o(x^4)$ .

**Solution:** By theorem 430,

$$\begin{aligned} 2\cos x + \sin x &= 2\left(1 - \frac{x^2}{2} + \frac{x^4}{24} + o(x^5)\right) + \left(x - \frac{x^3}{6} + o(x^4)\right) \\ &= 2 + x - x^2 - \frac{x^3}{6} + \frac{x^4}{12} + o(x^4) \\ &= 2\left(1 + \frac{x}{2} - \frac{x^2}{2} - \frac{x^3}{12} + \frac{x^4}{24} + o(x^4)\right), \end{aligned}$$

and so,

$$\begin{aligned} \log(2\cos x + \sin x) &= \log 2 \left(1 + \frac{x}{2} - \frac{x^2}{2} - \frac{x^3}{12} + \frac{x^4}{24} + o(x^4)\right) \\ &= \log 2 + \log \left(1 + \frac{x}{2} - \frac{x^2}{2} - \frac{x^3}{12} + \frac{x^4}{24} + o(x^4)\right) \\ &= \log 2 + \left(\frac{x}{2} - \frac{x^2}{2} - \frac{x^3}{12} + \frac{x^4}{24} + o(x^4)\right) \\ &\quad - \frac{1}{2} \left(\frac{x}{2} - \frac{x^2}{2} - \frac{x^3}{12} + \frac{x^4}{24} + o(x^4)\right)^2 \\ &\quad + \frac{1}{3} \left(\frac{x}{2} - \frac{x^2}{2} - \frac{x^3}{12} + \frac{x^4}{24} + o(x^4)\right)^3 \\ &\quad - \frac{1}{4} \left(\frac{x}{2} - \frac{x^2}{2} - \frac{x^3}{12} + \frac{x^4}{24} + o(x^4)\right)^4 + o(x^4) \\ &= \log 2 + \left(\frac{x}{2} - \frac{x^2}{2} - \frac{x^3}{12} + \frac{x^4}{24}\right) - \frac{1}{2} \left(\frac{x^2}{4} - \frac{x^3}{2} + \frac{x^4}{6}\right) \\ &\quad + \frac{1}{3} \left(\frac{x^3}{8} - \frac{3x^4}{8}\right) - \frac{1}{4} \cdot \frac{x^4}{16} + o(x^4) \\ &= \log 2 + \frac{x}{2} - \frac{5x^2}{8} + \frac{5x^3}{24} - \frac{35x^4}{192} + o(x^4) \end{aligned}$$

as  $x \rightarrow 0$ .

## Homework

**Problem 6.9.1** Prove that the limit

$$\lim_{n \rightarrow +\infty} \left(1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n}\right) - \log n,$$

exists. The constant

$$\gamma = \lim_{n \rightarrow +\infty} \left(1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n}\right) - \log n$$

is called the Euler-Mascheroni constant. It is not known whether  $\gamma$  is irrational.

# Chapter 7

## Integrable Functions

### 7.1 The Area Problem

**432 Definition** Let  $f : [a; b] \rightarrow \mathbb{R}$  be bounded, say with  $m \leq f(x) \leq M$  for all  $x \in [a; b]$ . Corresponding to each partition  $\mathcal{P} = \{x_0, x_1, \dots, x_n\}$  of  $[a; b]$ , we define the *upper Darboux sum*

$$U(f, \mathcal{P}) = \sum_{k=1}^n \left( \sup_{x_{k-1} \leq x \leq x_k} f(x) \right) (x_k - x_{k-1}),$$

and the *lower Darboux sum*

$$L(f, \mathcal{P}) = \sum_{k=1}^n \left( \inf_{x_{k-1} \leq x \leq x_k} f(x) \right) (x_k - x_{k-1}).$$

Clearly

$$L(f, \mathcal{P}) \leq U(f, \mathcal{P}).$$

Finally, we put

$$\int_a^b f(x) dx = \inf_{\mathcal{P} \text{ is a partition of } [a; b]} U(f, \mathcal{P}),$$

which we call the *upper Riemann integral of f* and

$$\int_a^b f(x) dx = \sup_{\mathcal{P} \text{ is a partition of } [a; b]} L(f, \mathcal{P}).$$

which we call the *lower Riemann integral of f*.

**433 Definition** Let  $f : [a; b] \rightarrow \mathbb{R}$  be bounded. We say that  $f$  is *Riemann integrable* if  $\int_a^b f(x) dx = \int_a^b f(x) dx$ . In this case, we denote their common value by  $\int_a^b f(x) dx$  and call it the *Riemann integral of f over [a; b]*.

**434 THEOREM** Let  $f$  be a bounded function on  $[a; b]$  and let  $\mathcal{P} \subseteq \mathcal{P}'$  be two partitions of  $[a; b]$ . Then

$$L(f, \mathcal{P}) \leq L(f, \mathcal{P}') \leq U(f, \mathcal{P}') \leq U(f, \mathcal{P}).$$

**Proof:** Clearly is enough to prove this when  $\mathcal{P}'$  has exactly one more point than  $\mathcal{P}$ . Let

$$\mathcal{P} = \{x_0, x_1, \dots, x_n\}$$

with  $a = x_0 < x_1 < \dots < x_{n-1} < x_n = b$ . Let  $\mathcal{P}'$  have the extra point  $x_*$  with  $x_i < x_* < x_{i+1}$ . Observe that we have both  $\inf_{x_i \leq x \leq x_{i+1}} f(x) \leq \inf_{x_i \leq x \leq x_*} f(x)$  and  $\inf_{x_i \leq x \leq x_{i+1}} f(x) \leq \inf_{x_* \leq x \leq x_{i+1}} f(x)$  since the larger interval may contain smaller values of  $f$ . Then

$$\begin{aligned} \inf_{x_i \leq x \leq x_{i+1}} f(x)(x_{i+1} - x_i) &= \inf_{x_i \leq x \leq x_{i+1}} f(x)(x_{i+1} - x_* + x_* - x_i) \\ &= \inf_{x_i \leq x \leq x_{i+1}} f(x)(x_* - x_i) + \inf_{x_i \leq x \leq x_{i+1}} f(x)(x_{i+1} - x_*) \\ &\leq \inf_{x_i \leq x \leq x_*} f(x)(x_* - x_i) + \inf_{x_* \leq x \leq x_{i+1}} f(x)(x_{i+1} - x_*). \end{aligned}$$

Thus

$$\begin{aligned} L(f, \mathcal{P}) &= (\inf_{x_0 \leq x \leq x_1} f(x))(x_1 - x_0) + \dots + (\inf_{x_i \leq x \leq x_{i+1}} f(x))(x_{i+1} - x_i) + \dots + (\inf_{x_{n-1} \leq x \leq x_n} f(x))(x_n - x_{n-1}) \\ &\leq (\inf_{x_0 \leq x \leq x_1} f(x))(x_1 - x_0) + \dots + (\inf_{x_i \leq x \leq x_*} f(x))(x_* - x_i) + (\inf_{x_* \leq x \leq x_{i+1}} f(x))(x_{i+1} - x_*) + \dots + (\inf_{x_{n-1} \leq x \leq x_n} f(x))(x_n - x_{n-1}) \\ &= L(f, \mathcal{P}'). \end{aligned}$$

A similar argument shows that  $U(f, \mathcal{P}') \leq U(f, \mathcal{P})$ . Then we have

$$L(f, \mathcal{P}) \leq L(f, \mathcal{P}') \leq U(f, \mathcal{P}') \leq U(f, \mathcal{P})$$

proving the theorem.  $\square$

**435 THEOREM** Let  $f$  be a bounded function on  $[a; b]$  and let  $\mathcal{P}_1$  and  $\mathcal{P}_2$  be any two partitions of  $[a; b]$ . Then

$$L(f, \mathcal{P}_1) \leq U(f, \mathcal{P}_2)$$

**Proof:** Let  $\mathcal{P}' = \mathcal{P}_1 \cup \mathcal{P}_2$  be a common refinement for  $\mathcal{P}_1$  and  $\mathcal{P}_2$ . By Theorem 434,

$$L(f, \mathcal{P}_1) \leq L(f, \mathcal{P}_1 \cup \mathcal{P}_2) \leq U(f, \mathcal{P}_1 \cup \mathcal{P}_2) \leq U(f, \mathcal{P}_2),$$

and

$$L(f, \mathcal{P}_2) \leq L(f, \mathcal{P}_1 \cup \mathcal{P}_2) \leq U(f, \mathcal{P}_1 \cup \mathcal{P}_2) \leq U(f, \mathcal{P}_2),$$

whence the theorem follows.  $\square$

**436 THEOREM** Let  $f$  be a bounded function on  $[a; b]$ . Then  $\int_{-a}^b f(x) dx \leq \int_a^{-b} f(x) dx$ .

**Proof:** By Theorem 435,

$$L(f, \mathcal{P}_1) \leq U(f, \mathcal{P}_2) \implies \int_{-a}^b f(x) dx = \sup_{\mathcal{P}_1 \text{ is a partition of } [a; b]} L(f, \mathcal{P}_1) \leq U(f, \mathcal{P}_2),$$

and so

$$\int_{-a}^b f(x) dx \leq U(f, \mathcal{P}_2).$$

Taking now the infimum,

$$\int_{-a}^b f(x) dx \leq \inf_{\mathcal{P}_2 \text{ is a partition of } [a; b]} U(f, \mathcal{P}_2) = \int_a^{-b} f(x) dx,$$

and the result is established.  $\square$

**437 THEOREM** Let  $f$  be a bounded function on  $[a; b]$ . Then  $f$  is Riemann integrable if and only if  $\forall \varepsilon > 0, \exists \mathcal{P}$  a partition of  $[a; b]$  such that

$$U(f, \mathcal{P}) - L(f, \mathcal{P}) < \varepsilon.$$

**Proof:**

$\Leftarrow$  If for all  $\varepsilon > 0, U(f, \mathcal{P}) - L(f, \mathcal{P}) < \varepsilon$  then by Theorem 436,

$$L(f, \mathcal{P}) \leq \int_a^b f(x) dx \leq \int_a^b f(x) dx \leq U(f, \mathcal{P}) \Rightarrow 0 \leq \int_a^b f(x) dx - \int_a^b f(x) dx < \varepsilon,$$

and so  $\int_a^b f(x) dx = \int_a^b f(x) dx$ , which means that  $f$  is Riemann-integrable.

$\Rightarrow$  Suppose  $f$  is Riemann integrable. By the Approximation property of the supremum and infimum, for all  $\varepsilon > 0$  there exist partitions  $\mathcal{P}_1$  and  $\mathcal{P}_2$  such that

$$U(f, \mathcal{P}_2) - \int_a^b f(x) dx < \frac{\varepsilon}{2}, \quad \int_a^b f(x) dx - L(f, \mathcal{P}_1) < \frac{\varepsilon}{2}.$$

Hence by taking  $\mathcal{P} = \mathcal{P}_1 \cup \mathcal{P}_2$  then

$$U(f, \mathcal{P}) \leq U(f, \mathcal{P}_2) < \int_a^b f(x) dx + \frac{\varepsilon}{2} < L(f, \mathcal{P}_1) + \varepsilon < L(f, \mathcal{P}) + \varepsilon,$$

from where  $U(f, \mathcal{P}) - L(f, \mathcal{P}) < \varepsilon$ .

□

**438 Example** •  $f(x) = \begin{cases} 0 & x \text{ irrational,} \\ 1 & x \text{ rational.} \end{cases} \quad x \in [0; 1]$

Then  $U(f, \mathcal{P}) = 1, L(f, \mathcal{P}) = 0$ , for any partition  $\mathcal{P}$ , and so  $f$  is not Riemann integrable.

•  $f(x) = \begin{cases} 0 & x \text{ irrational,} \\ \frac{1}{q} & x \text{ rational} = \frac{p}{q} \text{ in lowest terms.} \end{cases} \quad x \in [0; 1]$

is Riemann integrable with

$$\int_0^1 f(x) dx = 0$$

**439 Definition** Let  $f$  be a bounded function on  $[a; b]$  and let  $\mathcal{P} = \{x_0, x_1, \dots, x_n\}$  be a partition of  $[a; b]$ . If  $t_k$  are selected so that  $x_{k-1} \leq t_k \leq x_k$ , put

$$S(f, \mathcal{P}) = \sum_{k=1}^n f(t_k)(x_k - x_{k-1}),$$

is the Riemann sum of  $f$  associated with  $\mathcal{P}$ .

**440 THEOREM** Let  $f_1, f_2, \dots, f_m$  be Riemann integrable over  $[a; b]$ , and let  $f: [a; b] \rightarrow \mathbb{R}$ . If for any subinterval  $I \subseteq [a; b]$  there exists strictly positive numbers  $a_1, a_2, \dots, a_m$  such that

$$\omega(f, I) \leq a_1 \omega(f_1, I) + a_2 \omega(f_2, I) + \dots + a_m \omega(f_m, I),$$

then  $f$  is also Riemann integrable.

**Proof:** Let  $\mathcal{P} = \{a = x_0 < x_1 < \dots < x_n = b\}$  be a partition of  $[a; b]$  selected so that for all  $j$ ,

$$U(f_j, \mathcal{P}) - L(f_j, \mathcal{P}) < \frac{\varepsilon}{a_1 + a_2 + \dots + a_m}.$$

Using the notation of the preceding theorem,

$$\begin{aligned} U(f, \mathcal{P}) - L(f, \mathcal{P}) &= Z(f, \mathcal{P}) \\ &= \sum_{k=1}^n \omega(f, [x_{k-1}; x_k])(x_k - x_{k-1}) \\ &\leq \sum_{k=1}^n \sum_{j=1}^m a_j \omega(f_j, [x_{k-1}; x_k])(x_k - x_{k-1}) \\ &= \sum_{j=1}^m a_j \sum_{k=1}^n \omega(f_j, [x_{k-1}; x_k])(x_k - x_{k-1}) \\ &= \sum_{j=1}^m a_j (U(f_j, \mathcal{P}) - L(f_j, \mathcal{P})) \\ &< \varepsilon, \end{aligned}$$

and the theorem follows from Theorem 437.  $\square$

**441 THEOREM (Algebra of Riemann Integrable Functions)** Let  $f$  and  $g$  be Riemann integrable functions on  $[a; b]$  and let  $\lambda \in \mathbb{R}$  be a constant. Then the following are also Riemann integrable

1.  $f + \lambda g$
2.  $|f|$
3.  $f g$
4. provided  $\inf_{x \in [a; b]} |g(x)| > 0$ , also  $\frac{1}{g}$
5. provided  $\inf_{x \in [a; b]} |g(x)| > 0$ , also  $\frac{f}{g}$

**Proof:** Since

$$|f(x) + \lambda g(x) - f(t) - \lambda g(t)| \leq |f(x) - f(t)| + |\lambda| |g(x) - g(t)|, \quad \text{and} \quad ||f(x) - f(t)|| \leq |f(x) - f(t)|,$$

we have

$$\omega(f + \lambda g, I) \leq \omega(f, I) + |\lambda| \omega(g, I) \quad \text{and} \quad \omega(|f|, I) \leq \omega(f, I),$$

from where the first two assertions follow, upon appealing to Theorem 440.

To prove the third assertion, put  $a_1 = \sup_{u \in [a; b]} |f(u)|$  and  $a_2 = \sup_{u \in [a; b]} |g(u)|$

$$\begin{aligned} |f(x)g(x) - f(t)g(t)| &= |f(x)(g(x) - g(t)) + g(t)(f(x) - f(t))| \\ &\leq |f(x)| |g(x) - g(t)| + |g(t)| |f(x) - f(t)| \\ &\leq \left( \sup_{u \in [a; b]} |f(u)| \right) |g(x) - g(t)| + \left( \sup_{u \in [a; b]} |g(u)| \right) |f(x) - f(t)| \\ &= a_1 |g(x) - g(t)| + a_2 |f(x) - f(t)|, \end{aligned}$$

which gives

$$\omega(fg, I) \leq a_1 \omega(f, I) + a_2 \omega(g, I),$$

and so the third assertion follows from Theorem 440.

To prove the fourth assertion, with  $a = \inf_{x \in [a; b]} |g(x)| > 0$ , observe that we have

$$\begin{aligned} \left| \frac{1}{g(x)} - \frac{1}{g(t)} \right| &= \frac{1}{|g(x)g(t)|} |g(x) - g(t)| \\ &\leq \frac{1}{a^2} |g(x) - g(t)|, \end{aligned}$$

and this gives  $\omega\left(\frac{1}{g}, I\right) \leq \frac{1}{a^2} \omega(g, I)$ . The fourth assertion now follows by again appealing to Theorem 440.

The fifth assertion follows from the third and the fourth.  $\square$

**442 THEOREM** Let  $f$  and  $g$  be Riemann integrable functions on  $[a; b]$  and let  $\lambda \in \mathbb{R}$  be a constant. Then

$$\int_a^b (f(x) + \lambda g(x)) dx = \int_a^b f(x) dx + \lambda \int_a^b g(x) dx.$$

**Proof:** Let  $\mathcal{P} = \{a = x_0 < x_1 < \dots < x_n = b\}$  be a partition of  $[a; b]$  and choose  $t_k$  such that  $t_k \in [x_{k-1}; x_k]$ . Then for any  $\varepsilon > 0$  there exist  $\delta > 0$  and  $\delta' > 0$  such that

$$\begin{aligned} \left| \sum_{k=1}^n f(t_k)(x_k - x_{k-1}) - \int_a^b f(x) dx \right| &< \frac{\varepsilon}{2} \quad \text{if } \|\mathcal{P}\| < \delta, \\ \left| \lambda \sum_{k=1}^n g(t_k)(x_k - x_{k-1}) - \lambda \int_a^b g(x) dx \right| &< \frac{\varepsilon}{2} \quad \text{if } \|\mathcal{P}\| < \delta'. \end{aligned}$$

Hence, if  $\|\mathcal{P}\| < \min(\delta, \delta')$ ,

$$\begin{aligned} &\left| \sum_{k=1}^n (f(t_k) + \lambda g(t_k))(x_k - x_{k-1}) - \int_a^b f(x) dx - \lambda \int_a^b g(x) dx \right| \\ &\leq \left| \sum_{k=1}^n f(t_k)(x_k - x_{k-1}) - \int_a^b f(x) dx \right| + \left| \lambda \sum_{k=1}^n g(t_k)(x_k - x_{k-1}) - \lambda \int_a^b g(x) dx \right| \\ &< \varepsilon \end{aligned}$$

proving the theorem.  $\square$

**443 THEOREM** Let  $f$  and  $g$  be Riemann integrable functions on  $[a; b]$  with  $f(x) \leq g(x)$  for all  $x \in [a; b]$ . Then

$$\int_a^b f(x) dx \leq \int_a^b g(x) dx.$$

**Proof:** The function  $h = g - f$  is positive for all  $x \in [a; b]$  and hence  $L(h, \mathcal{P}) \geq 0$  for all partitions  $\mathcal{P}$ . It is also integrable by Theorem 442. Thus

$$\int_a^b h(x) dx = \int_a^b h(x) dx \geq 0.$$

But

$$\int_a^b h(x) dx \geq 0 \implies 0 \leq \int_a^b (g(x) - f(x)) dx = \int_a^b g(x) dx - \int_a^b f(x) dx,$$

and so  $\int_a^b f(x) dx \leq \int_a^b g(x) dx$ , as claimed.  $\square$

**444 THEOREM (Triangle Inequality for Integrals)** Let  $f$  be a Riemann integrable function on  $[a; b]$ . Then

$$\left| \int_a^b f(x) dx \right| \leq \int_a^b |f(x)| dx.$$

**Proof:** By Theorem 441,  $|f|$  is integrable. Now, since  $-|f| \leq f \leq |f|$  we just need to apply Theorem 443 twice.  $\square$

**445 THEOREM (Chasles' Rule)** Let  $f$  be a Riemann integrable function on  $[a; b]$  and let  $c \in ]a; b[$ . Then  $f$  is Riemann integrable function on  $[a; c]$  and  $[c; b]$ . Moreover,

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx.$$

Conversely, if  $c \in ]a; b[$  and  $f$  is Riemann integrable on  $[a; c]$  and  $[c; b]$  then  $f$  is Riemann integrable on  $[a; b]$  and

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx.$$

**Proof:** Consider the partitions

$$\mathcal{P} = \{a = x_0 < x_1 < \cdots < x_m = c < x_{m+1} < \cdots < x_n = b\}, \quad \mathcal{P}' = \{a = x_0 < x_1 < \cdots < x_m = c\}, \quad \mathcal{P}'' = \{c = x_m < x_{m+1} < \cdots < x_n = b\}.$$

where by virtue of Theorem 437, given  $\varepsilon > 0$ , we choose  $\mathcal{P}$  so that

$$U(f, \mathcal{P}) - L(f, \mathcal{P}) < \varepsilon.$$

It follows that

$$(U(f, \mathcal{P}') - L(f, \mathcal{P}')) + (U(f, \mathcal{P}'') - L(f, \mathcal{P}'')) = U(f, \mathcal{P}) - L(f, \mathcal{P}) < \varepsilon.$$

Hence  $f$  is Riemann-integrable over both  $[a; c]$  and  $[c; b]$ . Observe that

$$0 \leq U(f, \mathcal{P}') - \int_a^c f(x) dx < \varepsilon, \quad 0 \leq U(f, \mathcal{P}'') - \int_c^b f(x) dx < \varepsilon,$$

$$0 \leq \int_a^c f(x) dx - L(f, \mathcal{P}') < \varepsilon, \quad 0 \leq \int_c^b f(x) dx - L(f, \mathcal{P}'') < \varepsilon,$$

and upon addition,

$$0 \leq U(f, \mathcal{P}) - \left( \int_a^c f(x) dx + \int_c^b f(x) dx \right) < 2\varepsilon,$$

$$0 \leq \left( \int_a^c f(x) dx + \int_c^b f(x) dx \right) - L(f, \mathcal{P}) < 2\varepsilon,$$

whence

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx,$$

as required.  $\square$

**446 THEOREM (Converse of Chasles' Rule)** Let  $f$  be a function defined on the interval  $[a; b]$  and let  $c \in ]a; b[$ . If  $f$  is Riemann-integrable on  $[a; c]$  and  $[c; b]$  then it is also Riemann integrable in  $[a; b]$  and

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx.$$



**Proof:** Since  $f$  is Riemann-integrable on both subintervals, it is bounded there, and so it is bounded on the larger subinterval. By Theorem 437, given  $\varepsilon > 0$  there exist partitions  $\mathcal{P}'$  and  $\mathcal{P}''$  such that

$$U_{[a;c]}(f, \mathcal{P}') - L_{[a;c]}(f, \mathcal{P}') < \varepsilon, \quad U_{[c;b]}(f, \mathcal{P}'') - L_{[c;b]}(f, \mathcal{P}'') < \varepsilon.$$

The above inequalities also hold in the refinement  $\mathcal{P} = \mathcal{P}' \cup \mathcal{P}''$ , and

$$U(f, \mathcal{P}) = U_{[a;c]}(f, \mathcal{P}) + U_{[c;b]}(f, \mathcal{P}), \quad L(f, \mathcal{P}) = L_{[a;c]}(f, \mathcal{P}) + L_{[c;b]}(f, \mathcal{P}).$$

We then deduce that

$$\begin{aligned} U(f, \mathcal{P}) - L(f, \mathcal{P}) &= (U_{[a;c]}(f, \mathcal{P}) + U_{[c;b]}(f, \mathcal{P})) - (L_{[a;c]}(f, \mathcal{P}) + L_{[c;b]}(f, \mathcal{P})) \\ &= (U_{[a;c]}(f, \mathcal{P}) - L_{[a;c]}(f, \mathcal{P})) - (U_{[a;c]}(f, \mathcal{P}) - L_{[c;b]}(f, \mathcal{P})) \\ &< 2\varepsilon, \end{aligned}$$

and so  $f$  is Riemann integrable in  $[a; b]$  by virtue of Theorem 437. Now

$$\begin{aligned} \int_a^b f(x) dx &\leq U(f, \mathcal{P}) \\ &< L(f, \mathcal{P}) + \varepsilon \\ &= L_{[a;c]}(f, \mathcal{P}) + L_{[c;b]}(f, \mathcal{P}) + \varepsilon \\ &\leq \int_a^c f(x) dx + \int_c^b f(x) dx + \varepsilon, \end{aligned}$$

and similarly

$$\begin{aligned} \int_a^b f(x) dx &\geq L(f, \mathcal{P}) \\ &> U(f, \mathcal{P}) - \varepsilon \\ &= U_{[a;c]}(f, \mathcal{P}) + U_{[c;b]}(f, \mathcal{P}) - \varepsilon \\ &\geq \int_a^c f(x) dx + \int_c^b f(x) dx - \varepsilon, \end{aligned}$$

hence

$$\int_a^c f(x) dx + \int_c^b f(x) dx - \varepsilon \leq \int_a^b f(x) dx \leq \int_a^c f(x) dx + \int_c^b f(x) dx + \varepsilon$$

giving the desired equality between integrals.  $\square$

**447 THEOREM** Let  $f$  be Riemann integrable over  $[a; b]$  and let  $g: \left[ \inf_{u \in [a; b]} f(u); \sup_{u \in [a; b]} f(u) \right] \rightarrow \mathbb{R}$  be continuous. Then  $g \circ f$  is Riemann integrable on  $[a; b]$ .

**Proof:** Since  $g$  is uniformly continuous on the compact interval  $\left[ \inf_{u \in [a; b]} f(u); \sup_{u \in [a; b]} f(u) \right]$ , for given  $\varepsilon > 0$  we may find  $\delta'$  such that

$$(s, t) \in \left[ \inf_{t \in [a; b]} f(t); \sup_{u \in [a; b]} f(u) \right]^2; \quad |s - t| < \delta' \implies |f(s) - f(t)| < \varepsilon.$$

Let  $\delta = \min(\delta', \varepsilon)$ . Since  $f$  is Riemann-integrable, we may choose a partition  $\mathcal{P} = \{a = x_0 < x_1 < \dots < x_n = b\}$  such that

$$U(f, \mathcal{P}) - L(f, \mathcal{P}) < \delta^2, \tag{7.1}$$

by virtue of Theorem 437. Let

$$m_k = \inf_{x_{k-1} \leq x \leq x_k} f(x); \quad M_k = \sup_{x_{k-1} \leq x \leq x_k} f(x);$$

$$m_k^* = \inf_{x_{k-1} \leq x \leq x_k} (g \circ f)(x); \quad M_k^* = \sup_{x_{k-1} \leq x \leq x_k} (g \circ f)(x).$$

We split the set of indices  $\{1, 2, \dots, n\}$  into two classes:

$$A = \{k : 1 \leq k \leq n, M_k - m_k < \delta\}; \quad B = \{k : 1 \leq k \leq n, M_k - m_k \geq \delta\}.$$

If  $k \in A$  and  $x_{k-1} \leq x \leq y \leq x_k$ , then

$$|f(x) - f(y)| \leq M_k - m_k < \delta \leq \delta' \implies |(g \circ f)(x) - (f \circ g)(y)| < \varepsilon,$$

whence  $M_k^* - m_k^* \leq \varepsilon$ . Therefore

$$\sum_{k \in A} (M_k^* - m_k^*) (x_k - x_{k-1}) \leq \varepsilon \sum_{k=1}^n (x_k - x_{k-1}) = \varepsilon(b - a).$$

If  $k \in B$  then  $M_k - m_k \geq \delta$  and by virtue of (7.1),

$$\delta \sum_{k \in B} (x_k - x_{k-1}) \leq \sum_{k \in B} (M_k - m_k)(x_k - x_{k-1}) \leq \sum_{1 \leq k \leq n} (M_k - m_k)(x_k - x_{k-1}) = U(f, \mathcal{P}) - L(f, \mathcal{P}) < \delta^2,$$

whence

$$\sum_{k \in B} (x_k - x_{k-1}) < \delta \leq \varepsilon.$$

Upon assembling all these inequalities, and letting  $M = \sup_{t \in \left[ \inf_{u \in [a; b]} f(u); \sup_{u \in [a; b]} f(u) \right]} |g(t)|$ , we obtain

$$\begin{aligned} U(g \circ f, \mathcal{P}) - L(g \circ f, \mathcal{P}) &= \sum_{k \in A} (M_k^* - m_k^*) (x_k - x_{k-1}) + \sum_{k \in B} (M_k^* - m_k^*) (x_k - x_{k-1}) \\ &\leq \varepsilon(b - a) + 2M \sum_{k \in B} (x_k - x_{k-1}) \\ &\leq \varepsilon(b - a) + 2M\varepsilon \\ &= \varepsilon(b - a + 2M), \end{aligned}$$

whence the result follows from Theorem 437.  $\square$

**448 Definition** If  $b < a$  we define  $\int_a^b f(x) dx = -\int_b^a f(x) dx$ . Also,  $\int_a^a f(x) dx = 0$ .

**449 THEOREM** A function  $f$  on  $[a; b]$  is Riemann integrable on  $[a; b]$  if and only if its set of discontinuities forms a set of Lebesgue measure 0.

**Proof:**

$\implies$  Given  $\gamma > 0$  and  $\delta > 0$ , put  $\varepsilon = \gamma\delta$ . Let  $f$  be Riemann integrable. There is a partition  $\mathcal{P} = \{a = x_0 < x_1 < \dots < x_n = b\}$  such that

$$U(f, \mathcal{P}) - L(f, \mathcal{P}) < \varepsilon.$$

Let  $x \in ]x_i; x_{i+1}[$  be such that  $\omega(f, x) \geq \gamma$ . Then

$$\sup_{x_i; x_{i+1}[} f(x) - \inf_{x_i; x_{i+1}[} f(x) \geq \gamma.$$

Now observe that

$$\{x \in [a; b] : \omega(f, x) \geq \delta\} = \left( \bigcup_{\sup f - \inf f \geq \gamma} x_i ; x_{i+1} \right] \cup \{x_0, x_1, \dots, x_n\}.$$

Hence

$$\begin{aligned} \mu(\{x \in [a; b] : \omega(f, x) \geq \gamma\}) &\leq \sum_{\sup_{x_i; x_{i+1}} [f(x) - \inf_{x_i; x_{i+1}} f(x)] \geq \gamma} |x_{i+1} - x_i| \\ &\leq \frac{1}{\gamma} \sum_i |x_{i+1} - x_i| \left( \sup_{x_i; x_{i+1}} f(x) - \inf_{x_i; x_{i+1}} f(x) \right) \\ &\leq \frac{1}{\gamma} (U(f, \mathcal{P}) - L(f, \mathcal{P})) \\ &< \frac{\varepsilon}{\gamma} \\ &= \delta. \end{aligned}$$

Letting  $\delta \rightarrow 0+$  and  $\gamma \rightarrow 0+$  we get  $\mu(\{x \in [a; b] : \omega(f, x) \geq 0\}) = 0$ , and in particular,  $\mu(\{x \in [a; b] : \omega(f, x) > 0\}) = 0$  which means that the set of discontinuities is a set of measure 0.

$\Leftarrow$  Conversely, assume  $\mu(\{x \in [a; b] : \omega(f, x) > 0\}) = 0$ . We can write

$$\{x \in [a; b] : \omega(f, x) > 0\} = \bigcup_{K \geq 1} \{x \in [a; b] : \omega(f, x) > \frac{1}{K}\}.$$

Fix  $K$  large enough so that  $\frac{1}{K} < \varepsilon$ . Since  $\mu(\{x \in [a; b] : \omega(f, x) \geq \frac{1}{K}\}) = 0$ , we can find open intervals  $I_j(K)$  such that

$$\{x \in [a; b] : \omega(f, x) \geq \frac{1}{K}\} \subseteq \bigcup_{j \geq 1} I_j(K), \quad \sum_{j \geq 1} \mu(I_j(K)) < \varepsilon.$$

It is easy to show that  $\{x \in [a; b] : \omega(f, x) > \frac{1}{K}\}$  is closed and bounded and hence compact, so we may find a finite subcover with

$$\{x \in [a; b] : \omega(f, x) > \frac{1}{K}\} \subseteq I_{j_1} \cup I_{j_2} \cup \dots \cup I_{j_N}.$$

Now

$$[a; b] \setminus (I_{j_1} \cup I_{j_2} \cup \dots \cup I_{j_N})$$

is a finite disjoint union of closed intervals, say  $J_1 \cup J_2 \cup \dots \cup J_M$ . If  $x \in J_i$  then  $\omega(f, x) < \frac{1}{K}$ . Thus on each of the  $J_i$  we may find so fine a partition that  $\omega(f, L) < \frac{1}{K}$  for every interval such partition. All these partitions and the endpoints of the  $J_{j_k}$  form a partition, say  $\mathcal{P}$ . Write  $\mathcal{S} = S_1 \cup S_2 \cup \dots \cup S_M$  for the intervals of the partition  $\mathcal{P}$  that are not the  $J_{j_k}$ . Observe that  $\omega(f, S_k) < \frac{1}{K}$ . Then

$$\begin{aligned} U(f, \mathcal{P}) - L(f, \mathcal{P}) &= \sum_{I_{j_k}} \left( \sup_{I_{j_k}} f - \inf_{I_{j_k}} f \right) \mu(I_{j_k}) + \sum_{S_k} \left( \sup_{S_k} f - \inf_{S_k} f \right) \mu(S_k) \\ &\leq 2 \sup_{[a; b]} |f| \sum_{k=1}^N \mu(I_{j_k}) + \frac{1}{K} \sum_{S_k} \mu(S_k) \\ &\leq 2 \sup_{[a; b]} |f| \varepsilon + (b-a) \varepsilon \\ &= \left( 2 \sup_{[a; b]} |f| + (b-a) \right) \varepsilon. \end{aligned}$$

This proves the theorem.


□

**450 COROLLARY** Every continuous function  $f$  on  $[a; b]$  is Riemann integrable on  $[a; b]$ .

**Proof:** This is immediate from Theorem 449. □

**451 COROLLARY** Every monotonic function  $f$  on  $[a; b]$  is Riemann integrable on  $[a; b]$ .

**Proof:** Since a countable set has measure 0, and since the set of discontinuities of a monotonic function is countable (Theorem 335), the result is immediate. □



### Homework

**Problem 7.1.1** Let  $f$  be a bounded function on  $[a; b]$ . Then  $f$  is Riemann integrable if and only if  $\forall \epsilon > 0, \exists \delta > 0$  such that for all partitions  $\mathcal{P}$  of  $[a; b]$ ,

$$\|\mathcal{P}\| < \delta \implies U(f, \mathcal{P}) - L(f, \mathcal{P}) < \epsilon.$$

**Problem 7.1.2** Let  $f$  be a bounded function on  $[a; b]$ . Then  $f$  is Riemann integrable on  $[a; b]$  if and only if

$$\lim_{\|\mathcal{P}\| \rightarrow 0} S(f, \mathcal{P})$$

exists and is finite. In this case we write  $\lim_{\|\mathcal{P}\| \rightarrow 0} S(f, \mathcal{P}) =$

$$\int_a^b f(x) dx.$$

**Problem 7.1.3** Let  $f$  be bounded on  $[a; b]$ . Then  $f$  is Riemann integrable on  $[a; b]$  if and only if for every  $\epsilon > 0, \epsilon' > 0$  there is a partition  $\mathcal{P}$  of  $[a; b]$  such that

$$\sum_{k=1}^n (x_k - x_{k-1}) \chi_{\{x \in [a; b]; \omega(f, [x_{k-1}; x_k]) \geq \epsilon'\}} < \epsilon.$$

Here  $\chi(\cdot)$  is the indicator function defined on a set  $E$  as

$$\chi_E(x) = \begin{cases} 1 & \text{if } x \in E \\ 0 & \text{if } x \notin E \end{cases}.$$

## 7.2 Integration

**452 THEOREM (First Fundamental Theorem of Calculus)** Let  $f: [a; b] \rightarrow \mathbb{R}$  be Riemann integrable on  $[a; b]$ . If there exists a differentiable function  $F: [a; b] \rightarrow \mathbb{R}$  such that  $F' = f$  then

$$\int_a^b f(x) dx = F(b) - F(a).$$

**Proof:** Given  $\varepsilon > 0$ , in view of Theorem 437, there is a partition  $\mathcal{P} = \{a = x_0 < x_1 < \cdots < x_n = b\}$  such that

$$U(f, \mathcal{P}) - L(f, \mathcal{P}) < \varepsilon.$$

Since  $F$  is differentiable on  $[a; b]$ , it is continuous on  $[a; b]$ . Applying the Mean Value Theorem to each partition subinterval  $[x_{k-1}; x_k]$ , we obtain  $c_k \in ]x_{k-1}; x_k[$  such that

$$F(x_k) - F(x_{k-1}) = f(c_k)(x_k - x_{k-1}).$$

This gives

$$F(b) - F(a) = \sum_{1 \leq k \leq n} (F(x_k) - F(x_{k-1})) = \sum_{1 \leq k \leq n} f(c_k)(x_k - x_{k-1}),$$

and since  $\inf_{u \in [x_{k-1}; x_k]} f(u) \leq f(c_k) \leq \sup_{u \in [x_{k-1}; x_k]} f(u)$ , we deduce that

$$L(f, \mathcal{P}) \leq F(b) - F(a) \leq U(f, \mathcal{P}).$$

Furthermore, we know that  $L(f, \mathcal{P}) \leq \int_a^b f(x) dx \leq U(f, \mathcal{P})$ . Hence, combining these two last inequalities,

$$\left| F(b) - F(a) - \int_a^b f(x) dx \right| < \varepsilon,$$

and the theorem follows.  $\square$

**453 THEOREM (Second Fundamental Theorem of Calculus)** Let  $f: [a; b] \rightarrow \mathbb{R}$  be Riemann integrable on  $[a; b]$ , and let

$$F(x) = \int_a^x f(t) dt, \quad x \in [a; b].$$

Then  $F$  is continuous on  $[a; b]$ . Moreover, if  $f$  is continuous at  $c \in ]a; b[$ , then  $F$  is differentiable at  $c$  and  $F'(c) = f(c)$ .

**Proof:** There is  $M > 0$  such that  $\forall x \in [a; b]$ ,  $|f(x)| \leq M$ . Now, if  $a \leq x < y \leq b$  with  $|x - y| < \frac{\varepsilon}{M}$ , then

$$|F(y) - F(x)| = \left| \int_x^y f(t) dt \right| \leq \int_x^y |f(t)| dt \leq \int_x^y M dt = M(y - x) < \varepsilon$$

Thus  $F$  is continuous on  $[a; b]$  and by Heine's Theorem, uniformly continuous on  $[a; b]$ . Now, take  $u \in ]a; b[$ , and observe that

$$x \neq u \implies \frac{F(x) - F(u)}{x - u} = \frac{1}{x - u} \int_u^x f(t) dt.$$

Moreover,

$$f(u) = \frac{1}{x - u} \int_u^x f(u) dt,$$

and therefore,

$$\frac{F(x) - F(u)}{x - u} - f(u) = \frac{1}{x - u} \int_u^x (f(t) - f(u)) dt.$$

Since  $f$  is continuous at  $u$ , there is  $\delta > 0$  such that

$$t \in [a; b], |t - u| < \delta \implies |f(t) - f(u)| < \varepsilon.$$

This gives

$$\left| \frac{F(x) - F(u)}{x - u} - f(u) \right| < \varepsilon$$

for  $x \in ]a; b[$  with  $|x - u| < \delta$ . From this it follows that  $F'(u) = f(u)$ .  $\square$

**454 THEOREM (Young's Inequality for Integrals)** Let  $f$  be a strictly increasing continuous function on  $[0; +\infty[$  and let  $f(0) = 0$ . If  $A > 0$  and  $B > 0$  then

$$AB \leq \int_0^A f(x) dx + \int_0^B f^{-1}(x) dx.$$

**Proof:** The inequality is evident from Figure 7.1. The rectangle of area  $AB$  fits nicely in the areas under the curves  $y = f(x), x \in [0; A]$  and  $x = f^{-1}(y), y \in [0; B]$ .  $\square$

**455 THEOREM (Hölder's Inequality for Integrals)** Let  $p > 1$  and put  $\frac{1}{q} = 1 - \frac{1}{p}$ . If  $f$  and  $g$  are Riemann integrable on  $[a; b]$  then

$$\left| \int_a^b f(x)g(x) dx \right| \leq \left( \int_a^b |f(x)|^p dx \right)^{1/p} \left( \int_a^b |g(x)|^q dx \right)^{1/q}.$$

**Proof:** First observe that all of  $|fg|, |f|^p$  and  $|g|^q$  are Riemann-integrable, in view of Theorem 441. Now, with  $f(x) = x^{p-1}$  in Young's Inequality (Theorem 454), we obtain,

$$AB \leq \frac{A^p}{p} + \frac{B^{1/(p-1)+1}}{1/(p-1)+1} = \frac{A^p}{p} + \frac{B^q}{q}. \tag{7.2}$$

If any of the integrals in the statement of the theorem is zero, the result is obvious. Otherwise put  $A^p = \int_a^b |f(x)|^p dx$ ,  $B^q = \int_a^b |g(x)|^q dx$ . Then by (7.2),

$$\frac{|f(x)g(x)|}{AB} \leq \frac{A^{-p} |f(x)|^p}{p} + \frac{B^{-q} |g(x)|^q}{q}.$$

Integrating throughout the above inequality,

$$\frac{1}{AB} \int_a^b |f(x)g(x)| dx \leq \frac{1}{pA^p} \int_a^b |f(x)|^p dx + \frac{1}{qB^q} \int_a^b |g(x)|^q dx = \frac{1}{p} + \frac{1}{q} = 1,$$

whence the theorem follows.  $\square$




Figure 7.1: Young's Inequality (Theorem 454).

**456 THEOREM** Let  $f : [a; b] \rightarrow \mathbb{R}$ . Then

1. If  $f$  is continuous on  $[a; b], \forall x \in [a; b], f(x) \geq 0, \exists c \in [a; b]$  with  $f(c) > 0$  then  $\int_a^b f(x) dx > 0$ .
2. If  $f, g$  are continuous on  $[a; b], \forall x \in [a; b], f(x) \leq g(x)$ , and  $\exists c \in [a; b]$  with  $f(c) < g(c)$  then  $\int_a^b f(x) dx < \int_a^b g(x) dx$ .

**Proof:** The second part follows from the first by considering  $f - g$ . Let us prove the first part.

Assume first that  $c \in ]a; b[$ . Then there is a neighbourhood  $]c - \delta; c + \delta[ \subseteq ]a; b[$  of  $c$ , with  $\delta > 0$ , such that  $\forall x \in ]c - \delta; c + \delta[, f(x) \geq \frac{f(c)}{2}$ . Therefore

$$\int_a^b f(x) dx \geq \int_{c-\delta}^{c+\delta} f(x) dx > \int_{c-\delta}^{c+\delta} \frac{f(c)}{2} dx = \delta f(c) > 0.$$

If  $c = a$  then we consider a neighbourhood of the form  $]a; a + \delta[$ , and similarly if  $c = b$ , we consider a neighbourhood of the form  $]b - \delta; b[$ .  $\square$

**457 THEOREM (First Mean Value Theorem for Integrals)** Let  $f, g$  be continuous on  $[a; b]$ , with  $g$  of constant sign on  $[a; b]$ .

Then there exists  $c \in ]a; b[$  such that

$$\int_a^b f(x)g(x)dx = f(c) \int_a^b g(x)dx.$$

**Proof:** If  $g$  is identically 0, there is nothing to prove. Similarly, if  $f$  is constant in  $[a; b]$  there is nothing to prove. Otherwise,  $g$  is always strictly positive or strictly negative in the interval  $[a; b]$ . Let

$$m = \inf_{x \in [a; b]} f(x); \quad M = \sup_{x \in [a; b]} f(x).$$

Then

$$m < \frac{\int_a^b f(x)g(x)dx}{\int_a^b g(x)dx} < M.$$

By the Intermediate Value Theorem, there is  $c \in ]a; b[$  such that

$$f(c) = \frac{\int_a^b f(x)g(x)dx}{\int_a^b g(x)dx},$$

proving the theorem.  $\square$

**458 THEOREM (Integration by Parts)** Let  $f, g$  be differentiable functions on  $[a; b]$  with  $f'$  and  $g'$  integrable on  $[a; b]$ . Then

$$\int_a^b f'(x)g(x)dx + \int_a^b f(x)g'(x)dx = f(x)g(x) \Big|_a^b = f(b)g(b) - f(a)g(a).$$

**Proof:** This follows at once from the Product Rule for Derivatives and the Second Fundamental Theorem of Calculus, since

$$(fg)' = f'g + fg' \implies f(b)g(b) - f(a)g(a) = \int_a^b (f(x)g(x))' dx = \int_a^b f'(x)g(x)dx + \int_a^b f(x)g'(x)dx.$$

$\square$

**459 COROLLARY (Repeated Integration by Parts)** Let  $n \in \mathbb{N}$ . If the  $n + 1$ -th derivatives  $f^{(n+1)}$  and  $g^{(n+1)}$  are continuous on  $[a; b]$  then

$$\int_a^b f(x)g^{(n+1)}(x)dx = \left( f(x)g^{(n)}(x) - f'(x)g^{(n-1)}(x) + f''(x)g^{(n-2)}(x) - \dots + (-1)^n f^{(n)}(x)g(x) \right) \Big|_a^b + (-1)^{n+1} \int_a^b f^{(n+1)}(x)g(x)dx.$$

**Proof:** Follows by inducting on  $n$  and applying Theorem 458.  $\square$

**460 THEOREM (Integration by Substitution)** Let  $g$  be a differentiable function on an open interval  $I$  such that  $g'$  is continuous on  $I$ . If  $f$  is continuous on  $g(I)$  then  $f \circ g$  is continuous on  $I$  and  $\forall (a, b) \in I^2$ ,

$$\int_a^b (f \circ g)(x) g'(x) dx = \int_{g(a)}^{g(b)} f(u) du.$$

**Proof:** Fix  $c \in I$  and put  $F(x) = \int_c^x f(u) du$ . By The Second Fundamental Theorem of Calculus,  $\forall x \in I, F'(x) = f(x)$ . Furthermore, let  $t(x) = F(g(x))$ . By The Chain Rule,  $t'(x) = (F' \circ g)g' = (f \circ g)g'$ . Therefore

$$\begin{aligned} \int_a^b (f \circ g)(x) g'(x) dx &= \int_a^b t'(x) dx \\ &= t(b) - t(a) \\ &= F(g(b)) - F(g(a)) \\ &= \int_c^{g(b)} f(u) du - \int_c^{g(a)} f(u) du \\ &= \int_{g(a)}^{g(b)} f(u) du, \end{aligned}$$

as was to be shown.  $\square$

**461 THEOREM (Second Mean Value Theorem for Integrals)** Let  $f, g$  be continuous on  $[a; b]$ , with  $g$  monotonic on  $[a; b]$ . Then there exists  $c \in ]a; b[$  such that

$$\int_a^b f(x) g(x) dx = g(a) \int_a^c f(x) dx + g(b) \int_c^b f(x) dx.$$

**Proof:** Put  $F(x) = \int_a^x f(t) dt$ . Then  $F'(x) = f(x)$ . Hence

$$\int_a^b f(x) g(x) dx = \int_a^b F'(x) g(x) dx = F(x) g(x) \Big|_a^b - \int_a^b F(x) g'(x) dx$$

and therefore

$$\int_a^b f(x) g(x) dx = F(b) g(b) - F(a) g(a) - \int_a^b F(x) g'(x) dx.$$

By the First Mean Value Theorem for Integrals and by the First Fundamental Theorem of Calculus, there is a  $c \in ]a; b[$  such that

$$\int_a^b F(x) g'(x) dx = F(c) \int_a^b g'(x) dx = F(c)(g(b) - g(a)).$$

Assembling all the above,

$$\begin{aligned} \int_a^b f(x) g(x) dx &= F(b) g(b) - F(a) g(a) - F(c)(g(b) - g(a)) \\ &= g(b)(F(b) - F(c)) + g(a)(F(c) - F(a)) \\ &= g(b) \int_c^b f(x) dx + g(a) \int_a^c f(x) dx, \end{aligned}$$

as desired.  $\square$

**462 THEOREM (Generalisation of the AM-GM Inequality)** Let  $a_i \geq 0, p_i \geq 0$  with  $p_1 + p_2 + \dots + p_n = 1$ . Then

$$G = a_1^{p_1} a_2^{p_2} \dots a_n^{p_n} \leq p_1 a_1 + p_2 a_2 + \dots + p_n a_n = A.$$

(Here we interpret  $0^0 = 1$ .)



**Proof:** There is a subindex  $k$  such that  $a_k \leq G \leq a_{k+1}$ . Hence

$$\sum_{i=1}^k p_i \int_{a_i}^G \left( \frac{1}{x} - \frac{1}{G} \right) dx + \sum_{i=k+1}^n p_i \int_G^{a_i} \left( \frac{1}{G} - \frac{1}{x} \right) dx \geq 0,$$

as all the integrands are  $\geq 0$ . Upon rearranging

$$\sum_{i=1}^n p_i \int_{a_i}^G \frac{1}{x} dx \leq \sum_{i=1}^n p_i \int_G^{a_i} \frac{1}{G} dx \implies \sum_{i=1}^n p_i (\log a_i - \log G) \leq \sum_{i=1}^n p_i \cdot \frac{a_i - G}{G} \implies 0 \leq \frac{A}{G} - 1,$$

obtaining the inequality  $\square$

## Homework

**Problem 7.2.1** Let  $p$  be a polynomial of degree at most 4 such that  $p(-1) = p(1) = 0$  and  $p(0) = 1$ . If  $p(x) \leq 1$  for  $x \in [-1; 1]$ , find the largest value of  $\int_{-1}^1 p(x) dx$ .

**Problem 7.2.2** Compute  $\int_0^3 x \|x\| dx$ .

**Problem 7.2.3** Let  $f$  be a differentiable function such that

$$f(x+h) - f(x) = e^{x+h} - h - e^x$$

and  $f(0) = 3$ . Find  $f(x)$ .

**Problem 7.2.4** Let  $f$  be a continuous function such that  $f(x)f(a-x) = 1$  and let  $a > 0$ . Find  $\int_0^a \frac{1}{f(x)+1} dx$ .

**Problem 7.2.5** Let  $f$  be a Riemann integrable function over every bounded interval and such that  $f(a+b) = f(a) + f(b)$  for all  $(a, b) \in \mathbb{R}^2$ . Demonstrate that  $f(x) = xf(1)$ .

**Problem 7.2.6** Compute  $\int_0^2 x \|x^2\| dx$ .

**Problem 7.2.7** Find  $\int_{-1}^2 |x^2 - 1| dx$ .

**Problem 7.2.8** Let  $n$  be a fixed integer. Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be given by

$$f(x) = \begin{cases} x & \text{if } x \leq 0 \\ 2^n & \text{if } 2^n - 2^{n-2} < x \leq 2^{n+1} - 2^{n-1} \end{cases}$$

Prove that  $\int_0^{2^n} f(x) dx = \int_0^{2^n} x dx = 2^{2n-1}$ .

**Problem 7.2.9 (Putnam 1938)** Evaluate the limit

$$\lim_{t \rightarrow 0} \frac{\int_0^t (1 + \sin 2x)^{1/x} dx}{t}.$$

**Problem 7.2.10** Find the value of  $\int_0^1 \max(x^2, 1-x) dx$ .

**Problem 7.2.11** Let  $a > 0$ . Let  $f$  be a continuous function on  $[0; a]$  such that  $f(x) + f(a-x)$  does not vanish on  $[0; a]$ . Evaluate  $\int_0^a \frac{f(x) dx}{f(x) + f(a-x)}$ .

**Problem 7.2.12** Let  $a > 0$ . Let  $F$  be a differentiable function such that  $\forall x \in [0; a]$   $F'(a-x) = F'(x)$ . Evaluate  $\int_0^a F(x) dx$ .

**Problem 7.2.13** Let  $n \geq 0$  be an integer. Let  $a$  be the unique differentiable function such that  $\forall x \in \mathbb{R}$

$$(a(x))^{2n+1} + a(x) = x.$$

Evaluate  $\int_0^x a(t) dt$ .

**Problem 7.2.14** Find  $\int_0^{\pi/2} \frac{\sin x dx}{\sin x + \cos x}$ .

**Problem 7.2.15** Find  $\int_0^{\pi/2} \frac{1 dx}{1 + (\tan x)\sqrt{2}}$ .

**Problem 7.2.16** Find  $\int \frac{1}{x\sqrt{x^2-1}} dx$ .

**Problem 7.2.17** Find  $\int \frac{1}{1 + \sqrt{x+1}} dx$ .

**Problem 7.2.18** Find  $\int \frac{x^{1/2}}{x^{1/2} - x^{1/3}} dx$ .

**Problem 7.2.19** Find  $\int \frac{a^{2x}}{\sqrt{a^x+1}} dx$ ,  $a > 0$ .

**Problem 7.2.20** Find  $\int \frac{1}{(e^x - e^{-x})^2} dx$ .

**Problem 7.2.21** Prove that  $\int_1^5 \frac{\lfloor x \rfloor}{x} dx = 4 \log(5) - 3 \log(2) - \log(3)$ .

**Problem 7.2.22** Find  $\int e^{e^x+x} dx$ .

**Problem 7.2.23** Find  $\int \tan x \log(\cos x) dx$ .

**Problem 7.2.24** Find  $\int \frac{\log \log x}{x \log x} dx$ .

**Problem 7.2.25** Find  $\int \frac{x^{18}-1}{x^3-1} dx$ .

**Problem 7.2.26** Find  $\int \frac{1}{x^8+x} dx$ .

**Problem 7.2.27** Find  $\int \frac{4^x}{2^x+1} dx$ .

**Problem 7.2.28** Find  $\int \frac{x^2}{(x+1)^{10}} dx$ .

**Problem 7.2.29** Find  $\int \frac{1}{1+e^x} dx$ .

**Problem 7.2.30** Find  $\int \frac{1}{1-\sin x} dx$ .

**Problem 7.2.31** Find  $\int \sqrt{1+\sin 2x} dx$ .

**Problem 7.2.32** Find  $\int \frac{x}{\sqrt{1-x^4}} dx$ .

**Problem 7.2.33** Find  $\int \sec^4 x dx$ .

**Problem 7.2.34** Find  $\int \sec^5 x dx$ .

**Problem 7.2.35** Find  $\int e^{x^{1/3}} dx$ .

**Problem 7.2.36** Find  $\int \log(x^2+1) dx$ .

**Problem 7.2.37** Find  $\int xe^x \cos x dx$ .

**Problem 7.2.38** Find  $\int x^{2/3} \log x dx$ .

**Problem 7.2.39** Find  $\int \sin(\log x) dx$ .

**Problem 7.2.40** Find  $\int \frac{\log \log x}{x} dx$ .

**Problem 7.2.41** ( $\int \sec x dx$  in three ways) A traditional indefinite integral is

$$\int \sec x dx = \log(\tan x + \sec x) + C.$$

Justify this formula.

Now, prove that  $\frac{1}{\cos x} = \frac{\cos x}{2(1+\sin x)} + \frac{\cos x}{2(1-\sin x)}$ . Use this to find a second formula for  $\int \sec x dx$ .

A third way is as follows. Using  $\sin 2\theta = 2 \sin \theta \cos \theta$  show that  $\int \csc x dx = \log \left| \tan \frac{x}{2} \right| + C$ . Now use  $\csc\left(\frac{\pi}{2} + x\right) = \sec x$  to find yet another formula for  $\int \sec x dx$ .

**Problem 7.2.42** Find  $\int (\arcsin x)^2 dx$ .

**Problem 7.2.43** Find  $\int \frac{dx}{\sqrt{x+1} + \sqrt{x-1}}$ .

**Problem 7.2.44**  $\int x \arctan x dx$ .

**Problem 7.2.45** Find  $\int \sqrt{\tan x} dx$ .

**Problem 7.2.46** Find  $\int \frac{2x+1}{x^2(x-1)} dx$ .

**Problem 7.2.47** Find  $\int \log(x + \sqrt{x}) dx$ .

**Problem 7.2.48** Find  $\int \frac{1}{x^4+1} dx$ .

**Problem 7.2.49** Find  $\int \frac{1}{x^3+1} dx$ .

**Problem 7.2.50** Demonstrate that for all strictly positive integers  $n$ ,

$$\left(1 + \frac{1}{n}\right)^n \left(1 + \frac{1}{4n}\right) < e < \left(1 + \frac{1}{n}\right)^n \left(1 + \frac{1}{2n}\right),$$

that is,  $e$  is contained in the second quarter of the interval  $\left[\left(1 + \frac{1}{n}\right)^n; \left(1 + \frac{1}{n}\right)^{n+1}\right]$ .

**7.3 Riemann-Stieltjes Integration**

**7.4 Euler's Summation Formula**

# Chapter 8

## Sequences and Series of Functions

### 8.1 Pointwise Convergence

**463 Definition** We say that a sequence of functions  $\{f_n\}_{n=1}^{+\infty} : I \rightarrow \mathbb{R}$  defined on an interval  $I \subseteq \mathbb{R}$  converges pointwise to the function  $f$  if  $\forall x \in I, \forall \varepsilon > 0 \exists N > 0$  (depending on  $\varepsilon$  and on  $x$ ) such that

$$n \geq N \implies |f_n(x) - f(x)| < \varepsilon.$$

**464 Example** The sequence of functions  $x \mapsto x^n, n = 1, 2, \dots$  converges pointwise on the interval  $[0; 1]$  to the function  $f : [0; 1] \rightarrow \{0, 1\}$  with

$$f(x) = \begin{cases} 0 & \text{if } x \in [0; 1[ \\ 1 & \text{if } x = 1 \end{cases}$$

### 8.2 Uniform Convergence

**465 Definition** We say that a sequence of functions  $\{f_n\}_{n=1}^{+\infty} : I \rightarrow \mathbb{R}$  defined on an interval  $I \subseteq \mathbb{R}$  converges uniformly to the function  $f$  if  $\forall x \in I, \forall \varepsilon > 0 \exists N > 0$  (depending only on  $\varepsilon$ ) such that

$$n \geq N \implies |f_n(x) - f(x)| < \varepsilon.$$

In this case we write  $f_n \xrightarrow{\text{unif}} f$ .

**466 THEOREM** Let  $\{f_n\}_{n=1}^{+\infty}$  be a sequence of functions defined over a common domain  $I$ . If there exists a numerical sequence  $\{a_n\}_{n=1}^{+\infty}$  with  $a_n \rightarrow 0$  as  $n \rightarrow +\infty$ , and a function  $f$  defined over  $I$  such that eventually

$$|f_n(x) - f(x)| \leq a_n,$$

then  $f_n \xrightarrow{\text{unif}} f$ .

**467 THEOREM** If the sequence of continuous functions  $\{f_n\}_{n=1}^{+\infty} : I \rightarrow \mathbb{R}$  defined on an open interval  $I \subseteq \mathbb{R}$  converges uniformly to  $f$  on  $I$ , then  $f$  is continuous on  $I$ . Moreover, if  $x_0 \in I$  then we may exchange the limits, as in

$$\lim_{n \rightarrow +\infty} \left( \lim_{x \rightarrow x_0} f_n(x) \right) = \lim_{x \rightarrow x_0} \left( \lim_{n \rightarrow +\infty} f_n(x) \right) = \lim_{x \rightarrow x_0} f(x).$$

**468 THEOREM** If the sequence of integrable functions  $\{f_n\}_{n=1}^{+\infty} : I \rightarrow \mathbb{R}$  defined on an open interval  $I \subseteq \mathbb{R}$  converges uniformly to  $f$  on  $I$ , then  $f$  is integrable on  $I$ . Moreover, if  $(a, b) \in I^2$  then we may exchange the limit with the integral, as in

$$\lim_{n \rightarrow +\infty} \left( \int_a^b f_n(x) dx \right) = \int_a^b \left( \lim_{n \rightarrow +\infty} f_n(x) \right) dx = \int_a^b f(x) dx.$$

## 8.3 Integrals and Derivatives of Sequences of Functions

### 8.4 Power Series

A *power series* about  $x = a$  is a series of the form

$$f(x) = \sum_{n=0}^{+\infty} a_n(x-a)^n.$$

This is a function of  $x$ , and truncating it gives polynomial approximations to  $f$ . The goal is to approximate “decent” functions about a given point  $x = a$ .

These expansions don’t necessarily make sense for all  $x$ . The region where the power series converges is called the *interval of convergence*.

**469 Example** Find the interval of convergence of the series  $\sum_{n=1}^{\infty} \frac{2^n(x-3)^n}{\sqrt{n}}$ .

**Solution:** *By the ratio test, the series will converge if*

$$\left| \frac{2^{n+1}(x-3)^{n+1}}{\sqrt{n+1}} \cdot \frac{\sqrt{n}}{2^n(x-3)^n} \right| = 2\sqrt{\frac{n}{n+1}}|x-3| \rightarrow r < 1,$$

that is when

$$2|x-3| < 1 \implies \frac{5}{2} < x < \frac{7}{2}.$$

The series converges absolutely when  $\frac{5}{2} < x < \frac{7}{2}$ . We must also test the endpoints. At  $x = \frac{5}{2}$  the series is  $\sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n}}$ ,

which converges conditionally by Leibniz’s Test. At  $x = \frac{7}{2}$  the series is  $\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}}$ , which diverges.

## 8.5 Maclaurin Expansions to know by inspection

•

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$$

• The sine is an odd function:

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots$$

• The cosine is an even function:

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots$$

• If  $a$  is a real constant,

$$(1+x)^a = 1 + ax + \frac{a(a-1)}{2!}x^2 + \frac{a(a-1)(a-2)}{3!}x^3 + \frac{a(a-1)(a-2)(a-3)}{4!}x^4 + \dots$$

**470 Example** Expand  $f(x) = \cos x$  around  $x = 1$ .

**Solution:** *We have*

$$\begin{aligned} \cos x &= \cos(x-1+1) \\ &= \cos(x-1)\cos 1 - \sin(x-1)\sin 1 \\ &= (\cos 1) \left( 1 - \frac{(x-1)^2}{2!} + \frac{(x-1)^4}{4!} - \dots \right) - (\sin 1) \left( (x-1) - \frac{(x-1)^3}{3!} + \frac{(x-1)^5}{5!} - \dots \right) \end{aligned}$$

## Homework

**Problem 8.5.1** Given a finite collection of closed squares of total area 3, prove that they can be arranged to cover the unit square.

**Problem 8.5.2** Given a finite collection of closed squares of total area  $\frac{1}{2}$ , prove that they can be arranged to cover the unit square, with no overlaps

## 8.6 Comparison Tests

### Homework

**Problem 8.6.1** Let  $\{a_n\}_{n=1}^{\infty}$  be a sequence of real numbers satisfying  $0 < a_n < 1$  for all  $n$ . Assume that  $\sum_{n=1}^{\infty} a_n$  diverges but  $\sum_{n=1}^{\infty} a_n^2$  converges. Let  $f$  be a function defined on  $[0; 1]$  whose second derivative

exists and is bounded on  $[0; 1]$ . Prove that if  $\sum_{n=1}^{\infty} f(a_n)$  converges, so does  $\sum_{n=1}^{\infty} |f(a_n)|$ .

## 8.7 Taylor Polynomials

### Homework

**Problem 8.7.1** Evaluate  $\int_0^1 (\log x)(\log(1-x))dx$ .

**Problem 8.7.3** Find the sum of the infinite series

**Problem 8.7.2** Evaluate the infinite series  $\sum_{n=1}^{\infty} \arctan \frac{2}{n^2}$ .

$$1 - \frac{1}{4} + \frac{1}{6} - \frac{1}{9} + \frac{1}{11} - \frac{1}{14} + \dots$$

## 8.8 Abel's Theorem

### Homework

**Problem 8.8.1** Put

$$a_n = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots + \frac{(-1)^{n+1}}{n} - \log 2.$$

Prove that  $\sum_{n=1}^{\infty} a_n$  converges and find its sum.

**Problem 8.8.3** Evaluate the sum

$$\sum_{n=0}^{\infty} \left( \frac{1}{4n+1} + \frac{1}{4n+3} - \frac{1}{2n+2} \right).$$

**Problem 8.8.2** Evaluate the sum

$$\sum_{n=1}^{\infty} \frac{1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n}}{n(n+1)}.$$

**Problem 8.8.4** Evaluate the limit

$$\lim_{\alpha \rightarrow 0} \frac{1}{\alpha} \cdot \int_0^{\pi} \tan(\alpha \sin x) dx.$$

# Appendix A

## Answers and Hints

1.1.1 Observe that  $A_n = \{0, n, 2n, 3n, \dots\}$ .

1.  $A_6$ .
2.  $\mathbb{N}$ .
3.  $\{0\}$ .

1.1.4 We have,

$$\begin{aligned}
 x \in (A \cup B) \cap C &\iff x \in (A \cup B) \text{ and } x \in C \\
 &\iff (x \in A \text{ or } x \in B) \text{ and } x \in C \\
 &\iff (x \in A \text{ and } x \in C) \text{ or } (x \in B \text{ and } x \in C) \\
 &\iff (x \in A \cap C) \text{ or } (x \in B \cap C) \\
 &\iff x \in (A \cap C) \cup (B \cap C),
 \end{aligned}$$

which establishes the equality.

1.1.13 We check the two statements

$$x \in A \times (B \setminus C) \iff x \in (A \times B) \setminus (A \times C).$$

Let us prove first  $\implies$ . By definition of  $\times$ ,  $x = (a, b)$ , where  $a \in A, b \in B, b \notin C$ . Thus  $x \in A \times B$  but  $x \notin A \times C$ . By definition of  $\setminus$  we are done. Now we prove the assertion  $\impliedby$ . By definition of  $\times$  and  $\setminus$ ,  $x = (a, b)$  where  $a \in A, b \in B$ . Since  $x \notin A \times C$ , we observe that  $b \notin C$ . Thus  $a \in A, b \in B \setminus C$ , and we gather that  $x \in A \times (B \setminus C)$ .

1.1.14 Attach a binary code to each element of the subset, **1** if the element is in the subset and **0** if the element is not in the subset. The total number of subsets is the total number of such binary codes, and there are  $2^N$  in number.

1.2.1 There are  $2^2 = 4$  such functions, namely:

- $f_1$  given by  $f_1(a) = f_1(b) = c$ . Observe that  $\text{Im}(f_1) = \{c\}$ .
- $f_2$  given by  $f_2(a) = f_2(b) = d$ . Observe that  $\text{Im}(f_2) = \{d\}$ .
- $f_3$  given by  $f_3(a) = c, f_3(b) = d$ . Observe that  $\text{Im}(f_3) = \{c, d\}$ .
- $f_4$  given by  $f_4(a) = d, f_4(b) = c$ . Observe that  $\text{Im}(f_4) = \{c, d\}$ .

1.2.2 Each of the  $n$  elements of  $A$  must be assigned an element of  $B$ , and hence there are  $\underbrace{m \cdot m \cdots m}_n = m^n$  possibilities, and thus  $m^n$  functions. If a function from  $A$  to  $B$  is injective then we must have  $n \leq m$  in view of Theorem 16. If

to different inputs we must assign different outputs then to the first element of  $A$  we may assign any of the  $m$  elements of  $B$ , to the second any of the  $m - 1$  remaining ones, to the third any of the  $m - 2$  remaining ones, etc., and so we have  $m(m - 1) \cdots (m - n + 1)$  injective functions.

1.2.4 Rename the independent variable, say  $h(1 - s) = 2s$ . Now, if  $1 - s = 3x$  then  $s = 1 - 3x$ . Hence

$$h(3x) = h(1 - s) = 2s = 2(1 - 3x) = 2 - 6x.$$

1.2.5 Put

$$p(x) = (1 - x^2 + x^4)^{2003} = a_0 + a_1x + a_2x^2 + \cdots + a_{8012}x^{8012}.$$

Then

$$\bullet a_0 = p(0) = (1 - 0^2 + 0^4)^{2003} = 1.$$

$$\bullet a_0 + a_1 + a_2 + \cdots + a_{8012} = p(1) = (1 - 1^2 + 1^4)^{2003} = 1.$$

•

$$\begin{aligned}
 a_0 - a_1 + a_2 - a_3 + \cdots - a_{8011} + a_{8012} &= p(-1) \\
 &= (1 - (-1)^2 + (-1)^4)^{2003} \\
 &= 1.
 \end{aligned}$$

$$\bullet \text{ The required sum is } \frac{p(1) + p(-1)}{2} = 1.$$

$$\bullet \text{ The required sum is } \frac{p(1) - p(-1)}{2} = 0.$$

1.2.6 We have

$$\begin{aligned} \left[ f\left(\frac{27+y^3}{y^3}\right) \right] \sqrt{\frac{27}{y^3}} &= \left[ f\left(\left(\frac{3}{y}\right)^3 + 1\right) \right]^3 \sqrt{\frac{3}{y}} \\ &= \left( \left[ f\left(\left(\frac{3}{y}\right)^3 + 1\right) \right] \sqrt{\frac{3}{y}} \right)^3 \\ &= 5^3 \\ &= 125. \end{aligned}$$

1.2.7 We have

$$\begin{aligned} f(2) &= (-1)^2 1 - 2f(1) &&= 1 - 2f(1) \\ f(3) &= (-1)^3 2 - 2f(2) &&= -2 - 2f(2) \\ f(4) &= (-1)^4 3 - 2f(3) &&= 3 - 2f(3) \\ f(5) &= (-1)^5 4 - 2f(4) &&= -4 - 2f(4) \\ \vdots & && \vdots \\ \vdots & && \vdots \\ f(999) &= (-1)^{999} 998 - 2f(998) &&= -998 - 2f(998) \\ f(1000) &= (-1)^{1000} 999 - 2f(999) &&= 999 - 2f(999) \\ f(1001) &= (-1)^{1001} 1000 - 2f(1000) &&= -1000 - 2f(1000) \end{aligned}$$

Adding columnwise,

$$f(2) + f(3) + \dots + f(1001) = 1 - 2 + 3 - \dots + 999 - 1000 - 2(f(1) + f(2) + \dots + f(1000)).$$

This gives

$$2f(1) + 3f(2) + f(3) + \dots + f(1000) + f(1001) = -500.$$

Since  $f(1) = f(1001)$  we have  $2f(1) + f(1001) = 3f(1)$ . Therefore

$$f(1) + f(2) + \dots + f(1000) = -\frac{500}{3}.$$

1.2.8 Set  $a = b = 0$ . Then  $(f(0))^2 = f(0)f(0) = f(0+0) = f(0)$ . This gives  $f(0)^2 = f(0)$ . Since  $f(0) > 0$  we can divide both sides of this equality to get  $f(0) = 1$ .

Further, set  $b = -a$ . Then  $f(a)f(-a) = f(a-a) = f(0) = 1$ . Since  $f(a) \neq 0$ , may divide by  $f(a)$  to obtain  $f(-a) = \frac{1}{f(a)}$ .

Finally taking  $a = b$  we obtain  $(f(a))^2 = f(a)f(a) = f(a+a) = f(2a)$ . Hence  $f(2a) = (f(a))^2$

1.2.9 To prove that  $f$  is injective, we prove that  $f(a) = f(b) \implies a = b$ . We have

$$\begin{aligned} f(a) = f(b) &\implies \frac{a-1}{a+1} = \frac{b-1}{b+1} \\ &\implies (a-1)(b+1) = (a+1)(b-1) \\ &\implies ab + a - b - 1 = ab - a + b - 1 \\ &\implies 2a = 2b \\ &\implies a = b, \end{aligned}$$

where  $f$  is injective. To prove that  $f$  is surjective we must prove that any  $y \in \mathbb{R} \setminus \{1\}$  has a pre-image  $a \in \mathbb{R} \setminus \{-1\}$  such that  $f(a) = y$ . That is,

$$\frac{a-1}{a+1} = y \implies a-1 = ya+y \implies a-ya = 1+y \implies a(1-y) = 1+y \implies a = \frac{1+y}{1-y}.$$

Thus  $f\left(\frac{1+y}{1-y}\right) = y$ , and  $f$  is surjective. This also serves to prove that  $f^{-1}(x) = \frac{1+x}{1-x}$ .

1.2.10 We have  $f^{[2]}(x) = f(x+1) = (x+1)+1 = x+2$ ,  $f^{[3]}(x) = f(x+2) = (x+2)+1 = x+3$  and so, recursively,  $f^{[n]}(x) = x+n$ .

1.2.14 We have  $f^{[2]}(x) = f(2x) = 2^2 x$ ,  $f^{[3]}(x) = f(2^2 x) = 2^3 x$  and so, recursively,  $f^{[n]}(x) = 2^n x$ .

1.2.15 Let  $y = 0$ . Then  $2g(x) = 2x^2$ , that is,  $g(x) = x^2$ . Let us check that  $g(x) = x^2$  works. We have

$$g(x+y) + g(x-y) = (x+y)^2 + (x-y)^2 = x^2 + 2xy + y^2 + x^2 - 2xy + y^2 = 2x^2 + 2y^2,$$

which is the functional equation given. Our choice of  $g$  works.

1.2.16 Let  $x = 1$ . Then  $f(y) = yf(1)$ . Since  $f(1)$  is a constant, we may let  $k = f(1)$ . So all the functions satisfying the above equation satisfy  $f(y) = ky$ .

1.2.17 From  $f(x) + 2f\left(\frac{1}{x}\right) = x$  we obtain  $f\left(\frac{1}{x}\right) = \frac{x}{2} - \frac{1}{2}f(x)$ . Also, substituting  $1/x$  for  $x$  on the original equation we get

$$f(1/x) + 2f(x) = 1/x.$$

Hence

$$f(x) = \frac{1}{2x} - \frac{1}{2}f(1/x) = \frac{1}{2x} - \frac{1}{2}\left(\frac{x}{2} - \frac{1}{2}f(x)\right),$$

which yields  $f(x) = \frac{2}{3x} - \frac{x}{3}$ .



1.2.18 We have

$$(f(x))^2 \cdot f\left(\frac{1-x}{1+x}\right) = 64x,$$

whence

$$(f(x))^4 \cdot \left(f\left(\frac{1-x}{1+x}\right)\right)^2 = 64^2 x^2 \quad (I)$$

Substitute  $x$  by  $\frac{1-x}{1+x}$ . Then

$$f\left(\frac{1-x}{1+x}\right)^2 f(x) = 64 \left(\frac{1-x}{1+x}\right). \quad (II)$$

Divide (I) by (II),

$$f(x)^3 = 64x^2 \left(\frac{1+x}{1-x}\right),$$

from where the result follows.

1.2.19 We have (i)  $f^{[2]}(x) = (f \circ f)(x) = f(f(x)) = \frac{1}{1 - \frac{1}{1-x}} = \frac{x-1}{x}$ .

(ii)  $f^{[3]}(x) = (f \circ f \circ f)(x) = f(f^{[2]}(x)) = f\left(\frac{x-1}{x}\right) = \frac{1}{1 - \frac{x-1}{x}} = x$ .

(iii) Notice that  $f^{[4]}(x) = (f \circ f^{[3]})(x) = f(f^{[3]}(x)) = f(x) = f^{[1]}(x)$ . We see that  $f$  is cyclic of period 3, that is,  $f^{[1]} = f^{[4]} = f^{[7]} = \dots, f^{[2]} = f^{[5]} = f^{[8]} = \dots, f^{[3]} = f^{[6]} = f^{[9]} = \dots$ . Hence  $f^{[69]}(x) = f^{[3]}(x) = x$ .

1.2.20 To see (i) observe that

$$f(a) = f(b) \implies g(f(a)) = g(f(b)) \implies a = b,$$

whence  $f$  is injective. (The first implication is clear, the second implication follows because  $g \circ f$  is injective.)

To see (ii), given  $y \in C$ ,  $\exists x \in A$  such that  $g(f(x)) = y$ , since  $g \circ f$  is surjective. But then, letting  $a = f(x) \in B$  we have  $g(a) = y$  and  $g$  is surjective.

1.3.1 The map  $f: [0; 1] \rightarrow [a; b]$   $f(x) = \frac{x-a}{b-a}$  is a bijection.

1.3.2 The map  $f: ]-\infty; +\infty[ \rightarrow ]0; +\infty[$   $f(x) = e^x$  is a bijection.

1.4.1 Both answers are "no." If  $a = -b = \sqrt{2}$ , which we will prove later on to be irrational, we have  $a + b = 0$ , rational, and  $ab = -2$ , also rational.

1.4.2 Let  $\omega = -\frac{1}{2} + i\frac{\sqrt{3}}{2}$ . Then  $\omega^2 + \omega + 1 = 0$  and  $\omega^3 = 1$ . Then

$$\begin{aligned} x &= a^3 + b^3 + c^3 - 3abc = (a+b+c)(a+\omega b+\omega^2 c)(a+\omega^2 b+c\omega), \\ y &= u^3 + v^3 + w^3 - 3uvw = (u+v+w)(u+\omega v+\omega^2 w)(u+\omega^2 v+\omega w). \end{aligned}$$

Then

$$\begin{aligned} (a+b+c)(u+v+w) &= au+av+aw+bu+bv+bw+cu+cv+cw, \\ (a+\omega b+\omega^2 c)(u+\omega v+\omega^2 w) &= au+bu+cv \\ &\quad +\omega(av+bu+cw) \\ &\quad +\omega^2(av+bu+cw), \end{aligned}$$

and

$$\begin{aligned} (a+\omega^2 b+\omega c)(u+\omega^2 v+\omega w) &= au+bu+cv \\ &\quad +\omega(av+bu+cw) \\ &\quad +\omega^2(av+bu+cw). \end{aligned}$$

This proves that

$$\begin{aligned} xy &= (au+bu+cv)^3 + (av+bu+cw)^3 + (av+bu+cw)^3 \\ &\quad - 3(au+bu+cv)(av+bu+cw)(av+bu+cw), \end{aligned}$$

which proves that  $S$  is closed under multiplication.

1.4.3 We have

$$\begin{aligned} x \circ y &= (x \circ y) \circ (x \circ y) \\ &= [y \circ (x \circ y)] \circ x \\ &= [(x \circ y) \circ x] \circ y \\ &= [(y \circ x) \circ x] \circ y \\ &= [(x \circ x) \circ y] \circ y \\ &= (y \circ y) \circ (x \circ x) \\ &= y \circ x, \end{aligned}$$

proving commutativity.

1.4.4 By (1.4)

$$x * y = ((x * y) * x) * x.$$

By (1.4) again

$$((x * y) * x) * x = ((x * y) * ((x * y) * y)) * x.$$

By (1.3)

$$((x * y) * ((x * y) * y)) * x = (y) * x = y * x,$$

which is what we wanted to prove.

To show that the operation is not necessarily associative, specialise  $\mathcal{S} = \mathbb{Z}$  and  $x * y = -x - y$  (the opposite of  $x$  minus  $y$ ). Then clearly in this case  $*$  is commutative, and satisfies (1.3) and (1.4) but

$$0 * (0 * 1) = 0 * (-0 - 1) = 0 * (-1) = -0 - (-1) = 1,$$

and

$$(0 * 0) * 1 = (-0 - 0) * 1 = (0) * 1 = -0 - 1 = -1,$$

evinced that the operation is not associative.

1.4.5 1. Clearly, if  $a, b$  are rational numbers,

$$|a| < 1, |b| < 1 \implies |ab| < 1 \implies -1 < ab < 1 \implies 1 + ab > 0,$$

whence the denominator never vanishes and since sums, multiplications and divisions of rational numbers are rational,  $\frac{a+b}{1+ab}$  is also rational. We must prove now that  $-1 < \frac{a+b}{1+ab} < 1$  for  $(a, b) \in ]-1; 1[$ . We have

$$\begin{aligned} -1 < \frac{a+b}{1+ab} < 1 &\iff -1 - ab < a + b < 1 + ab \\ &\iff -1 - ab - a - b < 0 < 1 + ab - a - b \\ &\iff -(a+1)(b+1) < 0 < (a-1)(b-1). \end{aligned}$$

Since  $(a, b) \in ]-1; 1[$ ,  $(a+1)(b+1) > 0$  and so  $-(a+1)(b+1) < 0$  giving the sinistral inequality. Similarly  $a-1 < 0$  and  $b-1 < 0$  give  $(a-1)(b-1) > 0$ , the dextral inequality. Since the steps are reversible, we have established that indeed  $-1 < \frac{a+b}{1+ab} < 1$ .

2. Since  $a \otimes b = \frac{a+b}{1+ab} = \frac{b+a}{1+ba} = b \otimes a$ , commutativity follows trivially. Now

$$\begin{aligned} a \otimes (b \otimes c) &= \frac{a \left( \frac{b+c}{1+bc} \right)}{1 + a \left( \frac{b+c}{1+bc} \right)} \\ &= \frac{a(1+bc) + b+c}{1+bc+a(b+c)} = \frac{a+b+c+abc}{1+ab+bc+ca}. \end{aligned}$$

One the other hand,

$$\begin{aligned} (a \otimes b) \otimes c &= \frac{\left( \frac{a+b}{1+ab} \right) c}{\left( \frac{a+b}{1+ab} \right) + c} \\ &= \frac{1 + \left( \frac{a+b}{1+ab} \right) c}{\frac{(a+b) + c(1+ab)}{1+ab}} \\ &= \frac{1+ab+(a+b)c}{1+ab+bc+ca}, \end{aligned}$$

whence  $\otimes$  is associative.

3. If  $a \otimes e = a$  then  $\frac{a+e}{1+ae} = a$ , which gives  $a+e = a+ea^2$  or  $e(a^2-1) = 0$ . Since  $a \neq \pm 1$ , we must have  $e = 0$ .

4. If  $a \otimes b = 0$ , then  $\frac{a+b}{1+ab} = 0$ , which means that  $b = -a$ , that is,  $a^{-1} = -a$ .

1.4.6 We must shew that  $\forall (a, b) \in G^2$  we have  $ab = ba$ . But

$$\begin{aligned} ab &= e(ab)e \\ &= (b^2)(ab)(a^2) \\ &= b((ba)(ba))a \\ &= b(ba)^2 a \\ &= b(e)a \\ &= ba, \end{aligned}$$

whence the result follows.

1.4.7 We have

$$\begin{aligned} (ab)^3 = a^3 b^3 &\implies ab(ab)ab = a(a^2 b^2)b \\ &\implies baba = a^2 b^2 \\ &\implies (ba)^2 = a^2 b^2. \end{aligned}$$

Similarly

$$(ab)^5 = a^5 b^5 \implies (ba)^4 = a^4 b^4.$$

But we also have

$$(ba)^4 = ((ba)^2)^2 = (a^2 b^2)^2 = a^2 (b^2 a^2) b^2,$$

and so

$$a^2 (b^2 a^2) b^2 = (ba)^4 = a^4 b^4 \implies b^2 a^2 = a^2 b^2.$$

We have shewn that  $\forall (a, b) \in G^2$

$$((ba)^2 = a^2 b^2) \text{ and } (b^2 a^2 = a^2 b^2).$$

Hence

$$\begin{aligned} (ba)^2 = a^2 b^2 = b^2 a^2 &\implies baba = b^2 a^2 \\ &\implies ab = ba, \end{aligned}$$

proving that the group is abelian.

1.4.8 Since

$$(ab)^{i+2} = \underbrace{(ab)(ab) \cdots (ab)}_{i+2 \text{ times}} = a(ba)^{i+1}b,$$

multiplying by  $a^{-1}$  on the left and by  $b^{-1}$  on the right the equality

$$(ab)^{i+2} = a^{i+2}b^{i+2} \tag{A.1}$$

we obtain

$$(ba)^{i+1} = (a)^{i+1}(b)^{i+1}. \tag{A.2}$$

By hypothesis

$$(ab)^{i+1} = (a)^{i+1}(b)^{i+1}. \tag{A.3}$$

Hence (A.2) and (A.3) yield

$$(ab)^{i+1} = (ba)^{i+1}. \tag{A.4}$$

Similarly, from (A.3) we obtain

$$(ab)^i = (ba)^i, \tag{A.5}$$

from which

$$(ab)^{-i} = (ba)^{-i}. \tag{A.6}$$

Multiplying (A.4) and (A.6) together, we deduce

$$ab = ba,$$

which is what we wanted to show.

1.5.1 The first two follow immediately from the Binomial Theorem, the first by putting  $x = y = 1$  and then  $x = -y = 1$ . The third follows by adding the first two and dividing by 2. The fourth follows by subtracting the second from the first and then dividing by 2.

1.5.2 If  $a = 10^3, b = 2$  then

$$1002004008016032 = a^5 + a^4b + a^3b^2 + a^2b^3 + ab^4 + b^5 = \frac{a^6 - b^6}{a - b}.$$

This last expression factorises as

$$\begin{aligned} \frac{a^6 - b^6}{a - b} &= (a + b)(a^2 + ab + b^2)(a^2 - ab + b^2) \\ &= 1002 \cdot 1002004 \cdot 998004 \\ &= 4 \cdot 4 \cdot 1002 \cdot 250501 \cdot k, \end{aligned}$$

where  $k < 250000$ . Therefore  $p = 250501$ .

1.5.4 From the Binomial Theorem,

$$(A + B)^3 = A^3 + 3A^2B + 3AB^2 + B^3 \implies A^3 + B^3 = (A + B)^3 - 3AB(A + B).$$

Then

$$\begin{aligned} a^3 + b^3 + c^3 - 3abc &= (a + b)^3 + c^3 - 3ab(a + b) - 3abc \\ &= (a + b + c)^3 - 3(a + b)c(a + b + c) - 3ab(a + b + c) \\ &= (a + b + c)((a + b + c)^2 - 3ac - 3bc - 3ab) \\ &= (a + b + c)(a^2 + b^2 + c^2 - ab - bc - ca). \end{aligned}$$

1.5.5

$$\binom{n}{k} = \frac{n!}{k!(n-k)!} = \frac{n}{k} \cdot \frac{(n-1)!}{(k-1)!(n-k)!} = \frac{n}{k} \binom{n-1}{k-1}.$$

1.5.6

$$\binom{n}{k} = \frac{n!}{k!(n-k)!} = \frac{n(n-1)}{k(k-1)} \cdot \frac{(n-2)!}{(k-2)!(n-k)!} = \frac{n}{k} \cdot \frac{n-1}{k-1} \binom{n-2}{k-2}.$$

1.5.7 We use the identity  $k \binom{n}{k} = n \binom{n-1}{k-1}$ . Then

$$\begin{aligned} \sum_{k=1}^n k \binom{n}{k} p^k (1-p)^{n-k} &= \sum_{k=1}^n n \binom{n-1}{k-1} p^k (1-p)^{n-k} \\ &= \sum_{k=0}^{n-1} n \binom{n-1}{k} p^{k+1} (1-p)^{n-1-k} \\ &= np \sum_{k=0}^{n-1} \binom{n-1}{k} p^k (1-p)^{n-1-k} \\ &= np(p+1-p)^{n-1} \\ &= np. \end{aligned}$$

1.5.8 We use the identity

$$k(k-1) \binom{n}{k} = n(n-1) \binom{n-2}{k-2}.$$

Then

$$\begin{aligned} \sum_{k=2}^n k(k-1) \binom{n}{k} p^k (1-p)^{n-k} &= \sum_{k=2}^n n(n-1) \binom{n-2}{k-2} p^k (1-p)^{n-k} \\ &= \sum_{k=0}^{n-2} n(n-1) \binom{n-2}{k} p^{k+2} (1-p)^{n-1-k} \\ &= n(n-1)p^2 \sum_{k=0}^{n-2} \binom{n-1}{k} p^k (1-p)^{n-2-k} \\ &= n(n-1)p^2(p+1-p)^{n-2} \\ &= n(n-1)p^2. \end{aligned}$$

1.5.9 We use the identity

$$(k - np)^2 = k^2 - 2knp + n^2 p^2 = k(k-1) + k(1-2np) + n^2 p^2.$$

Then

$$\begin{aligned} \sum_{k=0}^n (k - np)^2 \binom{n}{k} p^k (1-p)^{n-k} &= \sum_{k=0}^n (k(k-1) + k(1-2np) + n^2 p^2) \binom{n}{k} p^k (1-p)^{n-k} \\ &= \sum_{k=0}^n k(k-1) \binom{n}{k} p^k (1-p)^{n-k} \\ &\quad + (1-2np) \sum_{k=0}^n k \binom{n}{k} p^k (1-p)^{n-k} \\ &\quad + n^2 p^2 \sum_{k=0}^n \binom{n}{k} p^k (1-p)^{n-k} \\ &= n(n-1)p^2 + np(1-2np) + n^2 p^2 \\ &= np(1-p). \end{aligned}$$

1.5.11 Observe that the number of  $k$ -tuples with  $\min(a_1, a_2, \dots, a_k) = t$  is  $(n-t+1)^k - (n-t)^k$ .

1.7.2 The given equalities entail

$$\sum_{k=1}^n (x_k^2 - x_k)^2 = 0.$$

A sum of squares is 0 if and only if every term is 0. This gives the result.

1.7.3 The given equality entails that

$$\frac{1}{2} \left( (x_1 - x_2)^2 + (x_2 - x_3)^2 + \dots + (x_{n-1} - x_n)^2 + (x_n - x_1)^2 \right) = 0.$$

A sum of squares is 0 if and only if every term is 0. This gives the result.

1.7.4 Since  $aB < Ab$  one has  $a(b+B) = ab + aB < ab + Ab = (a+A)b$  so  $\frac{a}{b} < \frac{a+A}{b+B}$ . Similarly  $B(a+A) = aB + AB < Ab + AB = A(b+B)$  and so  $\frac{a+A}{b+B} < \frac{A}{B}$ .

We have

$$\frac{7}{10} < \frac{11}{15} \implies \frac{7}{10} < \frac{18}{25} < \frac{11}{15} \implies \frac{7}{10} < \frac{25}{35} < \frac{18}{25} < \frac{11}{15}.$$

Since  $\frac{25}{35} = \frac{5}{7}$ , we have  $q \leq 7$ . Could it be smaller? Observe that  $\frac{5}{6} > \frac{11}{15}$  and that  $\frac{4}{6} < \frac{7}{10}$ . Thus by considering the cases with denominators  $q = 1, 2, 3, 4, 5, 6$ , we see that no such fraction lies in the desired interval. The smallest denominator is thus 7.

1.7.5 We have

$$(r-s+t)^2 - t^2 = (r-s+t-t)(r-s+t+t) = (r-s)(r-s+2t).$$

Since  $t-s \leq 0$ ,  $r-s+2t = r+s+2(t-s) \leq r+s$  and so

$$(r-s+t)^2 - t^2 \leq (r-s)(r+s) = r^2 - s^2$$

which gives

$$(r-s+t)^2 \leq r^2 - s^2 + t^2.$$

1.7.6 Using the CBS Inequality (Theorem 87) on  $\sum_{k=1}^n (a_k b_k c_k)$  once we obtain

$$\sum_{k=1}^n a_k b_k c_k \leq \left( \sum_{k=1}^n a_k^2 b_k^2 \right)^{1/2} \left( \sum_{k=1}^n c_k^2 \right)^{1/2}.$$

Using CBS again on  $\left( \sum_{k=1}^n a_k^2 b_k^2 \right)^{1/2}$  we obtain

$$\begin{aligned} \sum_{k=1}^n a_k b_k c_k &\leq \left( \sum_{k=1}^n a_k^2 b_k^2 \right)^{1/2} \left( \sum_{k=1}^n c_k^2 \right)^{1/2} \\ &\leq \left( \sum_{k=1}^n a_k^4 \right)^{1/4} \left( \sum_{k=1}^n b_k^4 \right)^{1/4} \left( \sum_{k=1}^n c_k^2 \right)^{1/2}, \end{aligned}$$

which gives the required inequality.

1.7.7 This follows directly from the AM-GM Inequality applied to  $1, 2, \dots, n$ :

$$n^{1/n} (1 \cdot 2 \cdot \dots \cdot n)^{1/n} < \frac{1+2+\dots+n}{n} = \frac{n+1}{2},$$

where strict inequality follows since the factors are unequal for  $n > 1$ .

1.7.8 First observe that for integer  $k$ ,  $1 < k < n$ ,  $k(n-k+1) = k(n-k) + k > 1(n-k) + k = n$ . Thus

$$n!^2 = (1 \cdot n)(2 \cdot (n-1))(3 \cdot (n-2)) \dots ((n-1) \cdot 2)(n \cdot 1) > n \cdot n \cdot \dots \cdot n = n^n.$$

1.7.9 From the Binomial Theorem, for  $n \geq 2$ ,

$$2^n = (1+1)^n = \binom{n}{0} + \binom{n}{1} + \binom{n}{2} + \dots + \binom{n}{n} > \binom{n}{2} = \frac{n(n-1)}{2} \implies 2^{n+1} > n(n-1).$$

This establishes the inequality for  $n \geq 2$ . For  $n=0$ ,  $0=0(0-1) < 2^{0+1}$  and for  $n=1$ ,  $0=1(1-1) < 2^{1+1}$ , so the inequality is true for all natural numbers.

1.7.10 Assume without loss of generality that  $a \geq b \geq c$ . Then  $a \geq b \geq c$  is similarly sorted as itself, so by the Rearrangement Inequality

$$a^2 + b^2 + c^2 = aa + bb + cc \geq ab + bc + ca.$$

This also follows directly from the identity

$$a^2 + b^2 + c^2 - ab - bc - ca = \left(a - \frac{b+c}{2}\right)^2 + \frac{3}{4}(b-c)^2.$$

One can also use the AM-GM Inequality thrice:

$$a^2 + b^2 \geq 2ab; \quad b^2 + c^2 \geq 2bc; \quad c^2 + a^2 \geq 2ca,$$

and add.

1.7.11 Assume without loss of generality that  $a \geq b \geq c$ . Then  $a \geq b \geq c$  is similarly sorted as  $a^2 \geq b^2 \geq c^2$ , so by the Rearrangement Inequality

$$a^3 + b^3 + c^3 = aa^2 + bb^2 + cc^2 \geq a^2b + b^2c + c^2a,$$

and

$$a^3 + b^3 + c^3 = aa^2 + bb^2 + cc^2 \geq a^2c + b^2a + c^2b.$$

Upon adding

$$a^3 + b^3 + c^3 = aa^2 + bb^2 + cc^2 \geq \frac{1}{2} (a^2(b+c) + b^2(c+a) + c^2(a+b)).$$

Again, if  $a \geq b \geq c$  then

$$ab \geq ac \geq bc,$$

thus

$$a^3 + b^3 + c^3 \geq a^2b + b^2c + c^2a = (ab)a + (bc)b + (ac)c \geq (ab)c + (bc)a + (ac)b = 3abc.$$

This last inequality also follows directly from the AM-GM Inequality, as

$$(a^3 b^3 c^3)^{1/3} \leq \frac{a^3 + b^3 + c^3}{3},$$

or from the identity

$$a^3 + b^3 + c^3 - 3abc = (a+b+c)(a^2 + b^2 + c^2 - ab - bc - ca),$$

and the inequality of problem 1.7.10.

1.7.12 We apply  $n$  times the Rearrangement Inequality

$$\begin{aligned} \check{a}_1 \hat{b}_1 + \check{a}_2 \hat{b}_2 + \dots + \check{a}_n \hat{b}_n &\leq a_1 b_1 + a_2 b_2 + \dots + a_n b_n &\leq \hat{a}_1 \hat{b}_1 + \hat{a}_2 \hat{b}_2 + \dots + \hat{a}_n \hat{b}_n \\ \check{a}_1 \hat{b}_1 + \check{a}_2 \hat{b}_2 + \dots + \check{a}_n \hat{b}_n &\leq a_1 b_2 + a_2 b_3 + \dots + a_n b_1 &\leq \hat{a}_1 \hat{b}_1 + \hat{a}_2 \hat{b}_2 + \dots + \hat{a}_n \hat{b}_n \\ \check{a}_1 \hat{b}_1 + \check{a}_2 \hat{b}_2 + \dots + \check{a}_n \hat{b}_n &\leq a_1 b_3 + a_2 b_4 + \dots + a_n b_2 &\leq \hat{a}_1 \hat{b}_1 + \hat{a}_2 \hat{b}_2 + \dots + \hat{a}_n \hat{b}_n \\ &\vdots \\ \check{a}_1 \hat{b}_1 + \check{a}_2 \hat{b}_2 + \dots + \check{a}_n \hat{b}_n &\leq a_1 b_n + a_2 b_1 + \dots + a_n b_{n-1} &\leq \hat{a}_1 \hat{b}_1 + \hat{a}_2 \hat{b}_2 + \dots + \hat{a}_n \hat{b}_n \end{aligned}$$

Adding we obtain the desired inequalities.

1.7.14 Use the fact that  $(b-a)^2 = (\sqrt{b}-\sqrt{a})^2(\sqrt{b}+\sqrt{a})^2$ .

1.7.15 Let

$$A = \frac{1}{2} \cdot \frac{3}{4} \cdot \frac{5}{6} \cdots \frac{9999}{10000}$$

and

$$B = \frac{2}{3} \cdot \frac{4}{5} \cdot \frac{6}{7} \cdots \frac{10000}{10001}.$$

Clearly,  $x^2 - 1 < x^2$  for all real numbers  $x$ . This implies that

$$\frac{x-1}{x} < \frac{x}{x+1}$$

whenever these four quantities are positive. Hence

$$\begin{array}{ccc} 1/2 & < & 2/3 \\ 3/4 & < & 4/5 \\ 5/6 & < & 6/7 \\ \vdots & & \vdots \\ \vdots & & \vdots \\ 9999/10000 & < & 10000/10001 \end{array}$$

As all the numbers involved are positive, we multiply both columns to obtain

$$\frac{1}{2} \cdot \frac{3}{4} \cdot \frac{5}{6} \cdots \frac{9999}{10000} < \frac{2}{3} \cdot \frac{4}{5} \cdot \frac{6}{7} \cdots \frac{10000}{10001},$$

or  $A < B$ . This yields  $A^2 = A \cdot A < A \cdot B$ . Now

$$A \cdot B = \frac{1}{2} \cdot \frac{2}{3} \cdot \frac{3}{4} \cdot \frac{4}{5} \cdot \frac{5}{6} \cdot \frac{6}{7} \cdot \frac{7}{8} \cdots \frac{9999}{10000} \cdot \frac{10000}{10001} = \frac{1}{10001},$$

and consequently,  $A^2 < A \cdot B = 1/10001$ . We deduce that  $A < 1/\sqrt{10001} < 1/100$ .

1.7.16 Observe that for  $k \geq 1$ ,  $(x+k)^2 > (x+k)(x+k-1)$  and so

$$\frac{1}{(x+k)^2} < \frac{1}{(x+k)(x+k-1)} = \frac{1}{x+k-1} - \frac{1}{x+k}.$$

Hence

$$\begin{aligned} \frac{1}{(x+1)^2} + \frac{1}{(x+2)^2} + \frac{1}{(x+3)^2} + \cdots + \frac{1}{(x+n-1)^2} + \frac{1}{(x+n)^2} &< \frac{1}{x(x+1)} + \frac{1}{(x+1)(x+2)} + \frac{1}{(x+2)(x+3)} + \cdots + \frac{1}{(x+n-2)(x+n-1)} + \frac{1}{(x+n-1)(x+n)} \\ &= \frac{1}{x} - \frac{1}{x+1} + \frac{1}{x+1} - \frac{1}{x+2} + \frac{1}{x+2} - \frac{1}{x+3} + \cdots + \frac{1}{x+n-2} - \frac{1}{x+n-1} + \frac{1}{x+n-1} - \frac{1}{x+n} \\ &= \frac{1}{x} - \frac{1}{x+n}. \end{aligned}$$

1.7.17 For  $1 \leq i \leq n$ , we have

$$\left| \frac{2}{i} - 1 - \frac{1}{n} \right| \leq 1 - \frac{1}{n} \iff \left( \frac{2}{i} - \left( 1 + \frac{1}{n} \right) \right)^2 \leq \left( 1 - \frac{1}{n} \right)^2 \iff \frac{4}{i^2} - \frac{4}{i} \left( 1 + \frac{1}{n} \right) + \frac{4}{n} \leq 0 \iff \frac{(i-n)(i-1)}{i^2 n} \leq 0.$$

Thus

$$\left| \sum_{i=1}^n \frac{x_i}{i} \right| = \frac{1}{2} \left| \sum_{i=1}^n \left( \frac{2}{i} - \left( 1 + \frac{1}{n} \right) \right) x_i \right|,$$

as  $\sum_{i=1}^n x_i = 0$ . Now

$$\left| \sum_{i=1}^n \left( \frac{2}{i} - \left( 1 + \frac{1}{n} \right) \right) x_i \right| \leq \sum_{i=1}^n \left| \frac{2}{i} - 1 - \frac{1}{n} \right| |x_i| \leq \left( 1 - \frac{1}{n} \right) \sum_{i=1}^n |x_i| = \left( 1 - \frac{1}{n} \right).$$

1.7.18 Expanding the product

$$\prod_{k=1}^n (1+x_k) = 1 + \sum_{k=1}^n x_k + \sum_{1 \leq i < j \leq n} x_i x_j + \cdots \geq 1 + \sum_{k=1}^n x_k,$$

since the  $x_k \geq 0$ . When  $n = 1$  equality is obvious. When  $n > 1$  equality is achieved when  $\sum_{1 \leq i < j \leq n} x_i x_j = 0$ .

1.7.19 Assume  $a \geq b \geq c$ . Put  $s = a + b + c$ . Then

$$-a \leq -b \leq -c \implies s-a \leq s-b \leq s-c \implies \frac{1}{s-a} \geq \frac{1}{s-b} \geq \frac{1}{s-c}$$

and so the sequences  $a, b, c$  and  $\frac{1}{s-a}, \frac{1}{s-b}, \frac{1}{s-c}$  are similarly sorted. Using the Rearrangement Inequality twice:

$$\frac{a}{s-a} + \frac{b}{s-b} + \frac{c}{s-c} \geq \frac{a}{s-c} + \frac{b}{s-a} + \frac{c}{s-b}; \quad \frac{a}{s-a} + \frac{b}{s-b} + \frac{c}{s-c} \geq \frac{a}{s-b} + \frac{b}{s-c} + \frac{c}{s-a}.$$

Adding these two inequalities

$$2 \left( \frac{a}{s-a} + \frac{b}{s-b} + \frac{c}{s-c} \right) \geq \frac{b+c}{s-a} + \frac{c+a}{s-b} + \frac{c+a}{s-c},$$

whence

$$2 \left( \frac{a}{b+c} + \frac{b}{c+a} + \frac{c}{a+b} \right) \geq 3,$$

from where the result follows.

1.7.20 Let

$$P(n): \underbrace{\sqrt{a + \sqrt{a + \sqrt{a + \cdots + \sqrt{a}}}}}_{n \text{ radicands}} < \frac{1 + \sqrt{4a+1}}{2}.$$

Let us prove  $P(1)$ , that is

$$\forall a > 0, \sqrt{a} < \frac{1 + \sqrt{4a+1}}{2}.$$

To get this one, let's work backwards. If  $a > \frac{1}{4}$

$$\begin{aligned} \sqrt{a} < \frac{1 + \sqrt{4a+1}}{2} &\iff 2\sqrt{a} < 1 + \sqrt{4a+1} \\ &\iff 2\sqrt{a} - 1 < \sqrt{4a+1} \\ &\iff (2\sqrt{a} - 1)^2 < (\sqrt{4a+1})^2 \\ &\iff 4a - 4\sqrt{a} + 1 < 4a + 1 \\ &\iff -2\sqrt{a} < 0. \end{aligned}$$

all the steps are reversible and the last inequality is always true. If  $a \leq \frac{1}{4}$  then trivially  $2\sqrt{a} - 1 < \sqrt{4a+1}$ . Thus  $P(1)$  is true. Assume now that  $P(n)$  is true and let's derive  $P(n+1)$ . From

$$\underbrace{\sqrt{a + \sqrt{a + \sqrt{a + \cdots + \sqrt{a}}}}}_{n \text{ radicands}} < \frac{1 + \sqrt{4a+1}}{2} \implies \underbrace{\sqrt{a + \sqrt{a + \sqrt{a + \cdots + \sqrt{a}}}}}_{n+1 \text{ radicands}} < \sqrt{a + \frac{1 + \sqrt{4a+1}}{2}}.$$

we see that it is enough to show that

$$\sqrt{a + \frac{1 + \sqrt{4a+1}}{2}} = \frac{1 + \sqrt{4a+1}}{2}.$$

But observe that

$$(\sqrt{4a+1} + 1)^2 = 4a + 2\sqrt{4a+1} + 2 \implies \frac{1 + \sqrt{4a+1}}{2} = \sqrt{a + \frac{1 + \sqrt{4a+1}}{2}},$$

proving the claim.

1.7.21 From the AM-GM Inequality,

$$a + b \geq 2\sqrt{ab}; \quad b + c \geq 2\sqrt{bc}; \quad c + a \geq 2\sqrt{ca},$$

and the desired inequality follows upon multiplication of these three inequalities.

1.7.22 By the Rearrangement inequality

$$\sum_{k=1}^n \frac{a_k}{k^2} \geq \sum_{k=1}^n \frac{\hat{a}_k}{k^2} \geq \sum_{k=1}^n \frac{1}{k},$$

as  $\hat{a}_k \geq k$ , the  $a$ 's being pairwise distinct positive integers.

1.7.23 By the AM-GM Inequality,

$$\left(\frac{1}{x_1} \frac{1}{x_2} \cdots \frac{1}{x_n}\right)^{1/n} \leq \frac{\frac{1}{x_1} + \frac{1}{x_2} + \cdots + \frac{1}{x_n}}{n},$$

whence the inequality.

1.7.24 By the CBS Inequality,

$$(1 \cdot x_1 + 1 \cdot x_2 + \cdots + 1 \cdot x_n)^2 \leq (1^2 + 1^2 + \cdots + 1^2)(x_1^2 + x_2^2 + \cdots + x_n^2),$$

which gives the desired inequality.

1.7.25 Put

$$T_m = \sum_{1 \leq k \leq m} a_k - \sum_{m < k \leq n} a_k.$$

Clearly  $T_0 = -T_n$ . Since the sequence  $T_0, T_1, \dots, T_n$  changes signs, choose an index  $p$  such that  $T_{p-1}$  and  $T_p$  have different signs. Thus either  $T_{p-1} - T_p = 2|a_p|$  or  $T_p - T_{p-1} = 2|a_p|$ . We claim that

$$\min(|T_{p-1}|, |T_p|) \leq \max_{1 \leq k \leq n} |a_k|.$$

For

For, if contrariwise both  $|T_{p-1}| > \max_{1 \leq k \leq n} |a_k|$  and  $|T_p| > \max_{1 \leq k \leq n} |a_k|$ , then  $2|a_p| = |T_{p-1} - T_p| > 2 \max_{1 \leq k \leq n} |a_k|$ , a contradiction.

1.7.26 It is enough to prove this in the case when  $a, b, c, d$  are all positive. To this end, put  $O = (0, 0)$ ,  $L = (a, b)$  and  $M = (a + c, b + d)$ . By the triangle inequality  $OM \leq OL + LM$ , where equality occurs if and only if the points are collinear. But then

$$\sqrt{(a+c)^2 + (b+d)^2} = OM \leq OL + LM = \sqrt{a^2 + b^2} + \sqrt{c^2 + d^2},$$

and equality occurs if and only if the points are collinear, that is  $\frac{a}{b} = \frac{c}{d}$ .

1.7.31 Use Minkowski's Inequality and the fact that  $17^2 + 144^2 = 145^2$ . The desired value is  $S_{12}$ .

1.8.3 We have

$$\begin{aligned} \sum_{1 \leq i < j \leq n} (x_j - x_i) &= \sum_{1 \leq i < j \leq n} x_j - \sum_{1 \leq i < j \leq n} x_i \\ &= \sum_{j=2}^n (j-1)x_j - \sum_{i=1}^{n-1} (n-1)x_i \\ &= -(n-1)x_1 + \sum_{k=2}^{n-1} ((k-1) - (n-k))x_k + (n-1)x_n \\ &= -(n-1)x_1 - (n-3)x_2 - \cdots + (n-3)x_{n-1} + (n-1)x_n. \end{aligned}$$

This sum is maximal when the negative coefficients of the  $x_i$  are 0 and the positive coefficients of the  $x_i$  are equal to 1. If  $n$  is even the maximum is

$$1 + 3 + \cdots + (n-1).$$

If  $n$  is odd, the maximum coefficient is

$$2 + 4 + \cdots + (n-1).$$

The result follows thus.

1.8.4 We claim that  $3 \lfloor 2t \rfloor - 2 \lfloor 3t \rfloor = 0, \pm 1$  or  $-2$ . We can then take

$$P(x, y) = (3x-2y)(3x-2y-1)(3x-2y+1)(3x-2y+2).$$

In order to prove the claim, we observe that  $\lfloor x \rfloor$  has unit period, so it is enough to prove the claim for  $t \in [0, 1)$ . We divide  $[0, 1[$  as

$$[0, 1[ = [0, 1/3[ \cup [1/3, 1/2[ \cup [1/2, 2/3[ \cup [2/3, 1[.$$

If  $t \in [0, 1/3[$ , then both  $\lfloor 2t \rfloor$  and  $\lfloor 3t \rfloor$  are 0, and so  $3 \lfloor 2t \rfloor - 2 \lfloor 3t \rfloor = 0$ . If  $t \in [1/3, 1/2[$  then  $\lfloor 3t \rfloor = 1$  and  $\lfloor 2t \rfloor = 0$ , and so  $3 \lfloor 2t \rfloor - 2 \lfloor 3t \rfloor = -2$ . If  $t \in [1/2, 2/3[$ , then  $\lfloor 2t \rfloor = 1$ ,  $\lfloor 3t \rfloor = 1$ , and so  $3 \lfloor 2t \rfloor - 2 \lfloor 3t \rfloor = 1$ . If  $t \in [2/3, 1[$ , then  $\lfloor 2t \rfloor = 1$ ,  $\lfloor 3t \rfloor = 2$ , and  $3 \lfloor 2t \rfloor - 2 \lfloor 3t \rfloor = -1$ .

1.8.5 By the Binomial Theorem

$$(1 + \sqrt{2})^n + (1 - \sqrt{2})^n = 2 \sum_{0 \leq k \leq n/2} \binom{n}{2k} := 2N,$$

an even integer. Since  $-1 < 1 - \sqrt{2} < 0$ , it must be the case that  $(1 - \sqrt{2})^n$  is the fractional part of  $(1 + \sqrt{2})^n$  or  $(1 + \sqrt{2})^n + 1$  depending on whether  $n$  is odd or even, respectively. Thus for odd  $n$ ,  $(1 + \sqrt{2})^n - 1 < (1 + \sqrt{2})^n + (1 - \sqrt{2})^n < (1 + \sqrt{2})^n$ , whence  $(1 + \sqrt{2})^n + (1 - \sqrt{2})^n = \lfloor (1 + \sqrt{2})^n \rfloor$ , always even, and for  $n$  even  $2N := (1 + \sqrt{2})^n + (1 - \sqrt{2})^n = \lfloor (1 + \sqrt{2})^n \rfloor + 1$ , and so  $\lfloor (1 + \sqrt{2})^n \rfloor = 2N - 1$ , always odd for even  $n$ .

471 Example Prove that the first thousand digits after the decimal point in

$$(6 + \sqrt{35})^{1980}$$

are all 9's.

Solution: Reasoning as in the preceding problem,

$$(6 + \sqrt{35})^{1980} + (6 - \sqrt{35})^{1980} = 2k,$$

an even integer. But  $0 < 6 - \sqrt{35} < 1/10$ , (for if  $\frac{1}{10} < 6 - \sqrt{35}$ , upon squaring  $3500 < 3481$ , which is clearly nonsense), and hence  $0 < (6 - \sqrt{35})^{1980} < 10^{-1980}$  which yields

$$2k - 1 + \underbrace{0.9 \dots 9}_{1979 \text{ nines}} = 2k - \frac{1}{10^{1980}} < (6 + \sqrt{35})^{1980} < 2k,$$

This proves the assertion of the problem.

1.8.7 By squaring, it is easy to see that

$$\sqrt{4n+1} < \sqrt{n} + \sqrt{n+1} < \sqrt{4n+3}.$$

Neither  $4n+2$  nor  $4n+3$  are squares since squares are either congruent to  $0$  or  $1 \pmod 4$ , so

$$\lfloor \sqrt{4n+2} \rfloor = \lfloor \sqrt{4n+3} \rfloor,$$

and the result follows.

1.8.8 Let  $T_n$  be the  $n$ -th non-square. There is a natural number  $m$  such that  $m^2 < T_n < (m+1)^2$ . As there are  $m$  squares less than  $T_n$  and  $n$  non-squares up to  $T_n$ , we see that  $T_n = n + m$ . We have then  $m^2 < n + m < (m+1)^2$  or  $m^2 - m < n < m^2 + m + 1$ . Since  $n, m^2 - m, m^2 + m + 1$  are all integers, these inequalities imply  $m^2 - m + \frac{1}{4} < n < m^2 + m + \frac{1}{4}$ , that is to say,  $(m-1/2)^2 < n < (m+1/2)^2$ . But then  $m = \lfloor \sqrt{n + \frac{1}{2}} \rfloor$ . Thus the  $n$ -th non-square is  $T_n = n + \lfloor \sqrt{n + 1/2} \rfloor$ .

1.8.9 Assume on the contrary that

$$\frac{(a+2b)^2}{(a+b)^2} \geq 2 \implies a^2 + 4ab + 4b^2 \geq 2(a^2 + 2ab + b^2) \implies 2b^2 \geq a^2 \implies \frac{a^2}{b^2} \geq 2,$$

a contradiction. By adding,

$$\frac{a^2}{b^2} < 2, \quad \frac{(a+2b)^2}{(a+b)^2} < 2 \implies \frac{a^2}{b^2} + \frac{(a+2b)^2}{(a+b)^2} < 4 \implies \frac{(a+2b)^2}{(a+b)^2} - 2 < 2 - \frac{a^2}{b^2}.$$

1.8.10 It needs to be proved that

$$\left| \frac{2x+5}{x+2} - \sqrt{5} \right| < |x - \sqrt{5}|.$$

1.8.11 Consider the set  $E = \{x : x > 0, x^n < a\}$ . Show that  $E$  is bounded above with supremum  $b = \sup E$ . Then show that  $b^n = a$  by arguing by contradiction first against  $b^n < a$  and then against  $b^n > a$ . In the first case it may be advantageous to prove  $\left(b + \frac{a-b^n}{N}\right)^n < a$  for  $N$  large enough and use the Binomial Theorem to establish the inequality. In the second case consider  $b^n \left(1 + \frac{b^n}{Ma}\right)^{-n} > a$ , for integral  $M$  sufficiently large, again using the Binomial Theorem to establish the inequality.

2.2.1  $\{500; 501\}$ .

2.2.2  $\{1; 2\}$ .

2.2.3  $\mathbb{R}$ .

2.2.4  $\{1\}$ .

2.2.5  $\emptyset$ .

2.2.6  $\emptyset$ .

2.2.8 Closure is immediate. Most of the other axioms are inherited from the larger set  $\mathbb{R}$ . Observe  $0_F = 0, 1_F = 1$  and the multiplicative inverse of  $a + \sqrt{2}b, (a, b) \neq (0, 0)$  is

$$(a + \sqrt{2}b)^{-1} = \frac{1}{a + \sqrt{2}b} = \frac{a - \sqrt{2}b}{a^2 - 2b^2} = \frac{a}{a^2 - 2b^2} - \frac{\sqrt{2}b}{a^2 - 2b^2}.$$

Here  $a^2 - 2b^2 \neq 0$  since  $\sqrt{2}$  is irrational.

2.2.9 Assume  $(a, b) \in \mathbb{R}^2$  with  $a < b$ . If  $ab < 0$ , then  $0 \in D$  is between  $a$  and  $b$ . If  $0 < a < b$  then  $\sqrt{a} < \sqrt{b}$ , and since  $\mathbb{Q}$  is dense in  $\mathbb{R}$ , there is a rational number  $r$  such that  $\sqrt{a} < r < \sqrt{b} \implies a < r^2 < b$ . If  $a < b < 0$ , then  $-\sqrt{b} < -\sqrt{a}$ , and since  $\mathbb{Q}$  is dense in  $\mathbb{R}$ , there is a rational number  $s$  such that  $-\sqrt{b} < s < -\sqrt{a} \implies -b < s^2 < -a \implies a < -s^2 < b$ .

2.2.10 Assume  $(a, b) \in \mathbb{R}^2$  with  $a < b$ . There is a strictly positive integer  $n$  such that  $n > \frac{1}{b-a}$ . Thus

$$0 < \frac{1}{2^n} < \frac{1}{n} < b - a.$$

Put  $m = \lfloor 2^n a \rfloor + 1$ , and so by definition  $m - 1 \leq 2^n x < m$ . Hence

$$a < \frac{m}{2^n} \leq a + \frac{1}{2^n} < a + \frac{1}{n} < a + b - a = a.$$

2.6.6 For the proof of this let  $G$  be such a set (so that  $x + y$  is in  $G$  if  $x, y$  are, and  $G$  is closed), and suppose that we are not in cases (i) or (ii). Then it is enough to show that  $G$  contains arbitrarily small positive numbers, for then multiples of these will be dense in  $\mathbb{R}$ , but  $G$  being closed forces  $G = \mathbb{R}$ . To achieve this let  $\mathcal{S} = \inf\{x : x \in G, x > 0\}$ . If  $\mathcal{S} = 0$  we are done; but if  $\mathcal{S} > 0$  there cannot be numbers  $x \in G$  arbitrarily close to and greater than  $\mathcal{S}$ , for then  $x - \mathcal{S}$  would run through small positive members of  $G$ , in particular smaller than  $\mathcal{S}$ , contradicting its definition. This means that  $\mathcal{S}$  belongs itself to  $G$ , and from there it is easy to see that we are in case (ii) contrary to the assumption. Hence indeed  $\mathcal{S} = 0, G = \mathbb{R}$ .

3.2.1 No. Take  $a_n = \frac{1}{n}$ . Then  $a_n > 0$  always, but  $L = 0$ .

3.2.9 We have for  $n > 1$ ,

$$\frac{n^2}{n^2 + n} = \frac{n}{n^2 + n} + \dots + \frac{n}{n^2 + n} < \sum_{i=1}^n \frac{n}{n^2 + i} < \frac{n}{n^2 + 1} + \dots + \frac{n}{n^2 + 1} = \frac{n^2}{n^2 + 1},$$

and the result follows by the Sandwich Theorem since each of the sequences on the extremes converges to  $1$ .

3.2.10 Evidently  $n! \leq n^n$ . By problem 1.7.8, if  $n > 2$  then  $n^{n/2} \leq n!$ . Thus

$$\frac{1}{n} \leq \frac{1}{(n!)^{1/n}} \leq \frac{1}{n^{1/2}}$$

and the result follows by the Sandwich Theorem.

3.2.11 For  $n \geq 2$  we have

$$\frac{2^n}{n!} = \frac{2}{1} \cdot \frac{2}{2} \cdot \frac{2}{3} \cdots \frac{2}{n} \leq 2 \cdot 1 \cdot 1 \cdots 1 \cdot \frac{2}{n} = \frac{4}{n} \rightarrow 0.$$



3.2.12 There is a positive integer  $m$  with  $m^2 \leq n < (m+1)^2$ . Consider

$$\left| \frac{s_m^2}{m^2} - \frac{s_n}{n} \right|.$$

3.2.13 Since  $-1 \leq \sin n \leq 1$ , any possible limit must be finite. By way of contradiction assume that  $\sin n \rightarrow a$  as  $n \rightarrow +\infty$ . Then

$$\lim_{n \rightarrow +\infty} \sin n = a \implies \lim_{n \rightarrow +\infty} \sin(n+2) = a,$$

whence

$$\lim_{n \rightarrow +\infty} (\sin(n+2) - \sin n) = a - a = 0.$$

Now,

$$\sin(n+2) - \sin n = 2(\sin 1) \cos(n+1) \implies \cos(n+1) \rightarrow 0, \text{ as } n \rightarrow +\infty.$$

From

$$\cos(n+1) = \cos n \cos 1 - \sin n \sin 1$$

we obtain

$$\sin n = \frac{1}{\sin 1} (\cos n \cos 1 - \cos(n+1)) \rightarrow \frac{1}{\sin 1} (0 \cdot \cos 1 - 0) = 0,$$

and so  $a = 0$ . But then

$$1 = \sin^2 n + \cos^2 n \rightarrow 0^2 + 0^2 = 0,$$

a contradiction.

3.2.14 By problem 1.7.8,  $(n!)^{1/n} > \sqrt{n}$  for  $n \geq 3$ . Hence, for all  $M > 0$ , as long as  $n > M^2$  we will have

$$(n!)^{1/n} > \sqrt{n} > M,$$

giving the result.

3.2.16 We have

$$\sqrt{n+1} - \sqrt{n} = \frac{n+1-n}{\sqrt{n+1} + \sqrt{n}} = \frac{1}{\sqrt{n+1} + \sqrt{n}} < \frac{1}{2\sqrt{n}}.$$

Hence, as long as  $\frac{1}{2\sqrt{n}} < \epsilon$  that is, as long as  $n > \frac{1}{4\epsilon^2}$  we will have

$$|\sqrt{n+1} - \sqrt{n}| < \frac{1}{2\sqrt{n}} < \epsilon.$$

3.2.17 Write

$$\sum_{n=1}^{2^M} \frac{1}{n} = \sum_{m=1}^M \sum_{n=2^{m-1}+1}^{2^m} \frac{1}{n}.$$

Since  $1/n \geq 1/N$  when  $n \leq N$ , we gather that

$$\sum_{n=2^{m-1}+1}^{2^m} \frac{1}{n} \geq \sum_{n=2^{m-1}+1}^{2^m} 2^{-m} = (2^m - 2^{m-1})2^{-m} = \frac{1}{2}.$$

Thus

$$\sum_{n=1}^{2^M} \frac{1}{n} \geq \frac{M}{2}$$

and the sequence can be made arbitrarily large.

3.2.18 Observe that for  $n \geq 2$ ,

$$\begin{aligned} & \frac{\sqrt{(n-1)!}}{(1+\sqrt{1})(1+\sqrt{2})(1+\sqrt{3}) \cdots (1+\sqrt{n-1})} - \frac{\sqrt{(n)!}}{(1+\sqrt{1})(1+\sqrt{2})(1+\sqrt{3}) \cdots (1+\sqrt{n})} \\ &= \frac{\sqrt{(n-1)!}}{(1+\sqrt{1})(1+\sqrt{2})(1+\sqrt{3}) \cdots (1+\sqrt{n-1})} \left( 1 - \frac{\sqrt{n}}{1+\sqrt{n}} \right) \\ &= \frac{\sqrt{(n-1)!}}{(1+\sqrt{1})(1+\sqrt{2})(1+\sqrt{3}) \cdots (1+\sqrt{n})}. \end{aligned}$$

Therefore

$$\sum_{n=1}^K \frac{\sqrt{(n-1)!}}{(1+\sqrt{1})(1+\sqrt{2})(1+\sqrt{3}) \cdots (1+\sqrt{n})} = 1 - \frac{\sqrt{K!}}{(1+\sqrt{1})(1+\sqrt{2}) \cdots (1+\sqrt{K})}.$$

Now prove that  $u_K = \frac{\sqrt{K!}}{(1+\sqrt{1})(1+\sqrt{2}) \cdots (1+\sqrt{K})}$  decreases to 0.

3.2.19 Put  $x_1 = 1$ ,  $x_{n+1} = \sqrt{1+x_n}$ ,  $n \geq 0$ . We claim that the sequence  $\{x_n\}_{n=1}^{+\infty}$  is increasing and bounded above. By Theorem 165 the sequence must have a limit  $L$ . To prove that the sequence is increasing consider  $x_{n+1} - x_n$  (fill in this gap). To prove that the sequence is bounded, we claim that for all  $n \geq 1$ ,  $x_n < 4$ . For this is clearly true for  $n = 1$ . So assume that  $x_n < 4$ . Then

$$x_{n+1} = \sqrt{1+x_n} < \sqrt{1+4} = \sqrt{5} < 4,$$

and so the assertion follows by induction.

Since we have shown that  $L$  exists we now may compute

$$L = \lim_{n \rightarrow +\infty} x_{n+1} = \lim_{n \rightarrow +\infty} \sqrt{1+x_n} = \sqrt{1+L} \implies L = \sqrt{1+L} \implies L^2 - L - 1 = 0 \implies L = \frac{1+\sqrt{5}}{2},$$

where we have chosen the positive root as the sequence is clearly strictly positive.

3.2.20 By Theorem 56,  $1 + 2 + \cdots + n = \frac{n^2 + n}{2}$ , and the desired result follows.

3.2.21  $\frac{1}{3}; 1; \frac{1}{4}$ .

3.2.22 Put  $x_1 = 1$ ,  $x_{n+1} = \frac{1}{1+x_n}$ ,  $n \geq 0$ . We claim that the sequence  $\{x_n\}_{n=1}^{+\infty}$  is increasing and bounded above. By Theorem 165 the sequence must have a limit  $L$ . To prove that the sequence is increasing consider  $x_{n+1} - x_n$  (fill in this gap). To prove that the sequence is bounded, we claim that for all  $n \geq 1$ , prove by induction that  $x_n < 4$  (fill in this gap).

Since we have shown that  $L$  exists we now may compute

$$L = \lim_{n \rightarrow +\infty} x_{n+1} = \lim_{n \rightarrow +\infty} \frac{1}{1+x_n} = \frac{1}{1+L} \implies L = \frac{1}{1+L} \implies L^2 + L - 1 = 0 \implies L = \frac{\sqrt{5}-1}{2},$$

where we have chosen the positive root as the sequence is clearly strictly positive.

3.2.24 Assume that  $\left\{\frac{a_n}{b_n}\right\}_{n=1}^{+\infty}$  is increasing. Then

$$\frac{a_1}{b_1} \leq \frac{a_2}{b_2} \leq \dots \leq \frac{a_n}{b_n} \leq \frac{a_{n+1}}{b_{n+1}}.$$

Using Theorem 79,

$$\frac{a_1 + a_2 + \dots + a_n}{b_1 + b_2 + \dots + b_n} \leq \frac{a_n}{b_n} \leq \frac{a_{n+1}}{b_{n+1}} \implies \frac{a_1 + a_2 + \dots + a_n}{b_1 + b_2 + \dots + b_n} \leq \frac{a_1 + a_2 + \dots + a_{n+1}}{b_1 + b_2 + \dots + b_{n+1}} \leq \frac{a_{n+1}}{b_{n+1}},$$

proving that  $\left\{\frac{a_1 + a_2 + \dots + a_n}{b_1 + b_2 + \dots + b_n}\right\}_{n=1}^{+\infty}$  is also increasing. If  $\left\{\frac{a_n}{b_n}\right\}_{n=1}^{+\infty}$  were decreasing,  $\left\{-\frac{a_n}{b_n}\right\}_{n=1}^{+\infty}$  is increasing and we apply what we just have proved.

3.2.26 We have

$$\prod_{k=2}^n \frac{k^3 - 1}{k^3 + 1} = \prod_{k=2}^n \frac{k-1}{k+1} \prod_{k=2}^n \frac{k^2 + k + 1}{k^2 - k + 1}.$$

Now

$$\prod_{k=2}^n \frac{k-1}{k+1} = \frac{(n-1)!}{(n+1)!} = \frac{2}{n(n+1)}.$$

By observing that  $(k+1)^2 - (k+1) + 1 = k^2 + k + 1$ , we gather that

$$\prod_{k=2}^n \frac{k^2 - k + 1}{k^2 + k + 1} = \frac{3^2 + 3 + 1}{2^2 - 2 + 1} \cdot \frac{4^2 + 4 + 1}{3^2 + 3 + 1} \cdot \frac{5^2 + 5 + 1}{4^2 + 4 + 1} \cdots \frac{n^2 + n + 1}{(n-1)^2 + (n-1) + 1} = \frac{n^2 + n + 1}{3}.$$

Thus

$$\prod_{k=2}^n \frac{k^3 - 1}{k^3 + 1} = \frac{2}{3} \cdot \frac{n^2 + n + 1}{n(n+1)} \rightarrow \frac{2}{3},$$

as  $n \rightarrow +\infty$ .

3.2.27 Clearly  $x_n < x_n + \frac{1}{(n+1)^2} = x_{n+1}$ , and so the sequence is strictly increasing. By showing that  $x_n < 2 - \frac{1}{n} < 2$  we will be showing that it is bounded above, and hence convergent by Theorem 165. For  $n = 1$ ,  $x_1 = 1 = 2 - \frac{1}{1}$  and so the assertion is true. Assume that  $x_n < 2 - \frac{1}{n}$ . Then

$$x_{n+1} = x_n + \frac{1}{(n+1)^2} < 2 - \frac{1}{n} + \frac{1}{(n+1)^2} = 2 + \frac{n-(n+1)^2}{n(n+1)^2} = 2 - \frac{n^2+n+1}{n(n+1)^2} < 2 - \frac{n^2+n}{n(n+1)^2} = 2 - \frac{1}{n+1},$$

and the claimed inequality follows by induction. We will prove later on a result of Euler:

$$1 + \frac{1}{2^2} + \frac{1}{3^2} + \dots + \frac{1}{n^2} + \dots = \frac{\pi^2}{6}.$$

3.3.1 The product rule for limits only applies to a finite number of factors. Here the number of factors grows with  $n$ .

3.3.3 From Theorem 177, and since  $x \mapsto \log x$  is increasing,

$$\left(1 + \frac{1}{k+1}\right)^{k+1} < e < \left(1 + \frac{1}{k}\right)^{k+1} \implies (k+1) \log\left(1 + \frac{1}{k+1}\right) < 1 < (k+1) \log\left(1 + \frac{1}{k}\right).$$

Rearranging,

$$\log \frac{k+2}{k+1} < \frac{1}{k+1} < \log \frac{k+1}{k}.$$

Summing from  $k = n-1$  to  $k = 2n-1$ ,

$$\begin{aligned} \sum_{k=n-1}^{2n-1} \log \frac{k+2}{k+1} &< \sum_{k=n-1}^{2n-1} \frac{1}{k+1} < \sum_{k=n-1}^{2n-1} \log \frac{k+1}{k} \implies \log \frac{2n+1}{n} < \frac{1}{n} + \frac{1}{n+1} + \dots + \frac{1}{2n} < \log \frac{2n}{n-1} \\ &\implies \log\left(2 + \frac{1}{n}\right) < \frac{1}{n} + \frac{1}{n+1} + \dots + \frac{1}{2n} < \log\left(2 + \frac{2}{n-1}\right) \end{aligned}$$

and the result follows from the Sandwich Theorem.

3.3.4 Observe that

$$\begin{aligned} &\left(1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots + \frac{1}{2n-1} + \frac{1}{2n}\right) \\ &\quad - 2\left(\frac{1}{2} + \frac{1}{4} + \frac{1}{6} + \dots + \frac{1}{2n}\right) \\ &= \left(1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots + \frac{1}{2n-1} + \frac{1}{2n}\right) \\ &\quad - 2 \cdot \frac{1}{2} \left(1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots + \frac{1}{n}\right) \\ &= \left(1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots + \frac{1}{2n-1} + \frac{1}{2n}\right) \\ &\quad - \left(1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots + \frac{1}{n}\right) \\ &= \frac{1}{n+1} + \frac{1}{n+2} + \dots + \frac{1}{2n}, \end{aligned}$$

and use the result of problem 3.3.3.

3.3.7 We begin by looking at the Taylor series for  $e^x$ :

$$e^x = \sum_{k=0}^{\infty} \frac{x^k}{k!}.$$

This converges for every  $x \in \mathbb{R}$ , so  $e = \sum_{k=0}^{\infty} \frac{1}{k!}$  and  $e^{-1} = \sum_{k=0}^{\infty} (-1)^k \frac{1}{k!}$ . Arguing by contradiction, assume  $ae^2 + be + c = 0$  for integers  $a, b$  and  $c$ . That is the same as  $ae + b + ce^{-1} = 0$ . Fix  $n > |a| + |c|$ , then  $a, c \mid n!$  and  $\forall k \leq n, k! \mid n!$ . Consider

$$\begin{aligned} 0 &= n!(ae + b + ce^{-1}) = an! \sum_{k=0}^{\infty} \frac{1}{k!} + b + cn! \sum_{k=0}^{\infty} (-1)^k \frac{1}{k!} \\ &= b + \sum_{k=0}^n (a + c(-1)^k) \frac{n!}{k!} + \sum_{k=n+1}^{\infty} (a + c(-1)^k) \frac{n!}{k!} \end{aligned}$$

Since  $k! \mid n!$  for  $k \leq n$ , the first two terms are integers. So the third term should be an integer. However,

$$\begin{aligned} \left| \sum_{k=n+1}^{\infty} (a + c(-1)^k) \frac{n!}{k!} \right| &\leq (|a| + |c|) \sum_{k=n+1}^{\infty} \frac{n!}{k!} \\ &= (|a| + |c|) \sum_{k=n+1}^{\infty} \frac{1}{(n+1)(n+2) \cdots k} \\ &\leq (|a| + |c|) \sum_{k=n+1}^{\infty} (n+1)^{n-k} \\ &= (|a| + |c|) \sum_{t=1}^{\infty} (n+1)^{-t} \\ &= (|a| + |c|) \frac{1}{n} \end{aligned}$$

is less than 1 by our assumption that  $n > |a| + |c|$ . Since there is only one integer which is less than 1 in absolute value, this means that  $\sum_{k=n+1}^{\infty} (a + c(-1)^k) \frac{1}{k!} = 0$  for every sufficiently large  $n$  which is not the case because

$$\sum_{k=n+1}^{\infty} (a + c(-1)^k) \frac{1}{k!} - \sum_{k=n+2}^{\infty} (a + c(-1)^k) \frac{1}{k!} = (a + c(-1)^{n+1}) \frac{1}{(n+1)!}$$

is not identically zero. The contradiction completes the proof.

3.3.9 Apply Problem 2.6.6 We can apply this to the stated problem by observing that for a fixed  $d$ , a positive integer without square factors, the numbers  $a + b\sqrt{d}$  are quadratic integers if  $a, b$  are rational integers, and that the set of such numbers is an additive group of reals. Clearly the closure of this group (it, together with its set of limit points) is a group too, for if  $x_n \rightarrow x$  and  $y_n \rightarrow y$  then  $x_n + y_n \rightarrow x + y$ . The new group is not of form (i) or (ii), hence must be all reals, and the proof (of a slightly stronger theorem) is complete.

3.5.5  $a_n = o(n^2)$  does, since this says that  $\lim_{n \rightarrow +\infty} \frac{a_n}{n^2} = 0$ , whereas  $a_n = O(n^2)$  says that  $\frac{a_n}{n^2}$  is bounded by some positive constant.

3.5.6 False. Take  $a_n = 2n$ , for example. Then  $a_n \ll n$ ,  $\frac{a_n}{n} = 2$ , and so  $\frac{a_n}{n} \not\rightarrow 0$ .

3.5.7 True.  $\frac{a_n}{n} \rightarrow 0$  and so by Theorem 195,  $a_n \ll n$ .

3.5.8 False. Take  $a_n = n^{3/2}$ . Then  $\frac{a_n}{n^2} \rightarrow 0$  but  $a_n \neq O(n)$ .

3.5.9 True.  $\frac{a_n}{n} \rightarrow 0$  and so by Theorem 195,  $a_n \ll n$ . Since  $n \ll n^2$ , the assertion follows by transitivity.

4.1.1 This is a geometric series with common ratio  $|r| = \frac{2}{e} < 1$ , so it converges. We have

$$\sum_{n=3}^{\infty} \frac{2^n}{e^{n+1}} = \frac{2^3}{e^4} + \frac{2^4}{e^5} + \cdots = \frac{\frac{2^3}{e^4}}{1 - \frac{2}{e}} = \frac{8}{e^4 - 2e^3}.$$

4.1.2 Observe that

$$\frac{1}{4n^2 - 1} = \frac{1}{2(2n-1)} - \frac{1}{2(2n+1)}.$$

Hence

$$\sum_{n=2}^{+\infty} \frac{1}{4n^2 - 1} = \left( \frac{1}{2(1)} - \frac{1}{2(3)} \right) + \left( \frac{1}{2(3)} - \frac{1}{2(5)} \right) + \left( \frac{1}{2(5)} - \frac{1}{2(7)} \right) + \cdots = \frac{1}{2(1)} = \frac{1}{2}.$$

4.1.3 Since  $\tan(x-y) = \frac{\tan x - \tan y}{1 + \tan x \tan y}$ , observe that  $\arctan \frac{1}{n^2 + n + 1} = \arctan(n+1) - \arctan n$ . Hence the series telescopes to  $\lim_{n \rightarrow +\infty} \arctan(n+1) - \arctan 1 = \frac{\pi}{4}$ .

4.1.7 By unique factorisation of the integers, the desired sum is

$$\left( 1 + \frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \cdots \right) \left( 1 + \frac{1}{3} + \frac{1}{3^2} + \frac{1}{3^3} + \cdots \right) = \frac{1}{1 - \frac{1}{2}} \cdot \frac{1}{1 - \frac{1}{3}} = 3.$$

4.1.9 Since the sum of two convergent series is convergent by Theorem 232, if  $\sum_{n \geq 0} (a_n + b_n)$  then from the identity  $b_n = (a_n + b_n) - a_n$  we would deduce that  $\sum_{n \geq 0} b_n$  converges, a contradiction.

4.1.10 Put  $s_N = \sum_{1 \leq n \leq N} a_n$ . There is a positive constant  $M$  such that  $\forall N > 0, s_N \leq M$ . Observe that because the terms are positive

$$s_{N+1} = s_N + a_{N+1} \geq s_N,$$

and so the sequence  $\{s_N\}_{N=1}^{+\infty}$  is a monotonically increasing bounded above sequence and so it converges by Theorem 165.

This is not necessarily true if the series does not have positive terms. For example, the series  $\sum_{n \geq 1} (-1)^{n+1}$  has bounded partial sums, in fact they are either 1 or 0. But the sequence of partial sums then is

$$1, 0, 1, 0, 1, 0, \dots$$

which does not converge.

4.2.2 True. For, we must have  $a_n \rightarrow 0$  and so eventually  $0 < a_n \leq 1$ . This means that eventually  $a_n^2 \leq a_n$  and the series of squares converges by direct comparison to the original series.

4.2.3 True. Since  $a_n \rightarrow 0$ , we must have  $\sin a_n \sim a_n$  and so the series converges by asymptotic comparison to the original series. (Recall that  $\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$ .)

4.2.4 True. Since  $a_n \rightarrow 0$ , we must have  $\tan a_n \sim a_n$  and so the series converges by asymptotic comparison to the original series. (Recall that  $\lim_{x \rightarrow 0} \frac{\tan x}{x} = 1$ .)

4.2.5 False. Since  $a_n \rightarrow 0$ , we must have  $\cos a_n \rightarrow 1$  and so the series diverges by the  $n$ -th Term Test.

4.2.6 Only the fact that  $\frac{a_n}{n} \leq a_n$  is needed here.

4.2.7 Take  $a_n = \frac{1}{2^n}$ . Then  $a_n < \frac{1}{n^2}$  and  $\sum_{n=1}^{+\infty} \frac{1}{2^n} = 1$ .

4.2.8 Take  $a_n = \frac{1}{2n^2}$  or  $a_n = \frac{1}{n2^n}$ .

4.2.9 Take  $a_n = \frac{1}{2^n}$  or  $a_n = \frac{1}{n^n}$ .

4.2.10 For even  $n \geq 0$  take  $a_n = \frac{1}{2^n}$  and for odd  $n \geq 1$  take  $a_n = \frac{1}{3^n}$ . Then

$$\sum_{n=0}^{+\infty} a_n = \sum_{n=0}^{+\infty} \frac{1}{2^{2n}} + \sum_{n=1}^{+\infty} \frac{1}{3^{2n-1}},$$

and both series on the right are geometric convergent series. However if  $n$  is even,  $(a_n)^{1/n} = \frac{1}{2}$  and if  $n$  is odd  $(a_n)^{1/n} = \frac{1}{3}$  meaning that  $\lim_{n \rightarrow +\infty} (a_n)^{1/n}$  does not exist.

4.2.11 By the root test

$$a_n^{1/n} = \left( \frac{3^n}{n2^n} \right)^{1/n} = \frac{3}{n} \rightarrow 0 < 1,$$

and the series converges. By direct comparison, for  $n \geq 3$  we have

$$\frac{3^n}{n2^n} = \frac{3^n}{n^n} \cdot \frac{1}{n^n} \leq \frac{1}{n^n} \leq \frac{1}{n^3},$$

and the series converges by direct comparison to  $\sum_{n=1}^{\infty} \frac{1}{n^3}$ .

4.2.12 We divide the sum into decimal blocks. There are  $9^k$   $k$ -digit integers in the interval  $[10^k; 10^{k+1}[$  that do not have a 0 in their decimal representation. Thus

$$\sum_{n \in \mathcal{S}} \frac{1}{n} = \sum_{k=0}^{+\infty} \sum_{n \in [10^k; 10^{k+1}[ \cap \mathcal{S}} \frac{1}{n} \leq \sum_{k=0}^{+\infty} 9^k \left( \frac{1}{10^k} \right) = 10.$$

4.2.15

1.  $a_n \sim -\frac{e}{2n} \Rightarrow$  diverges.

6. converges iff  $|a| \neq 1$ .

2.  $a_n \sim \frac{\alpha}{2\alpha-1} e^{n(\alpha-2)} \Rightarrow$  converges iff  $\alpha < 2$ .

7. Converges.

3.  $a_n \sim -\frac{3}{n^2} \Rightarrow$  converges.

8.  $a_n \leq \frac{(n-1)(n-1)! + n!}{(n+2)!} \leq \frac{2}{(n+1)(n+2)} \Rightarrow$  converges.

4.  $a_n \sim \frac{1}{n^2} \Rightarrow$  converges.

9. Converges.

5.  $a_n \sim \sqrt{\frac{2}{n^3}} \Rightarrow$  converges.

10.  $a_n \neq 0 \Rightarrow$  diverges.

11.  $a_n = \frac{1}{n \log \log n} \Rightarrow$  converges.

5.1.1 Put  $a_n = \frac{1}{(2n - \frac{1}{2})\pi}$ ,  $b_n = \frac{1}{(2n + \frac{1}{2})\pi}$  for integer  $n \geq 1$ . Then  $a_n \rightarrow 0$  and  $b_n \rightarrow 0$ , but  $\sin \frac{1}{a_n} \rightarrow -1$  and  $\sin \frac{1}{b_n} \rightarrow +1$ , so the limit does not exist in view of Proposition 267.

5.2.10  $f(0) = 0$ , but for  $x > 0$ ,  $f(x) = \frac{1 + \sqrt{1+4x}}{2}$ , so  $f$  is not right-continuous at  $x = 0$ .

5.6.2 Consider a unit circle and take any point  $P$  on the circumference of the circle. Drop the perpendicular from  $P$  to the horizontal line,  $M$  being the foot of the perpendicular and  $Q$  the reflection of  $P$  at  $M$ . (refer to figure)

Let  $x = \angle POM$ .

For  $x$  to be in  $[0, \frac{\pi}{2}]$ , the point  $P$  lies in the first quadrant, as shown.

The length of line segment  $PM$  is  $\sin(x)$ . Construct a circle of radius  $MP$ , with  $M$  as the center.

Length of line segment  $PQ$  is  $2\sin(x)$ .

Length of arc  $PAQ$  is  $2x$ .

Length of arc  $PBQ$  is  $\pi \sin(x)$ .

Since  $PQ \leq$  length of arc  $PAQ$  (equality holds when  $x = 0$ ) we have  $2\sin(x) \leq 2x$ . This implies

$$\sin(x) \leq x$$

Since length of arc  $PAQ$  is  $\leq$  length of arc  $PBQ$  (equality holds true when  $x = 0$  or  $x = \frac{\pi}{2}$ ), we have  $2x \leq \pi \sin(x)$ . This implies

$$\frac{2}{\pi} x \leq \sin(x)$$

Thus we have

$$\frac{2}{\pi} x \leq \sin(x) \leq x, \forall x \in [0, \frac{\pi}{2}]$$

5.9.1 If  $p$  had odd degree, then, by the Intermediate Value Theorem it would have a real root. Let  $\alpha$  be its largest real root. Then

$$0 = p(\alpha)q(\alpha) = p(\alpha^2 + \alpha + 1)$$

meaning that  $\alpha^2 + \alpha + 1 > \alpha$  is a real root larger than the supposedly largest real root  $\alpha$ , a contradiction.

5.9.2 Observe that  $f(1000)f(f(1000)) = 1 \implies f(999) = \frac{1}{999}$ . So the range of  $f$  include all numbers from  $\frac{1}{999}$  to 999. By the intermediate value theorem, there is a real number  $a$  such that  $f(a) = 500$ . Thus

$$f(a)f(f(a)) = 1 \implies f(500) = \frac{1}{500}.$$

5.9.5

5.9.10 If either  $f(0) = 1$  or  $f(1) = 0$ , we are done. So assume that  $0 \geq f(0) < 1$  and  $0 < f(1) \leq 1$ . Put  $g(x) = f(x) + x - 1$ . Then  $g(0) = f(0) - 1 < 0$  and  $g(1) = f(1) > 0$ . By Bolzano's Theorem there is a  $c \in ]0; 1[$  such that  $g(c) = 0$ , that is,  $f(c) + c - 1 = 0$ , as required.

5.9.11 Consider  $g(x) = f(x) - f(x + 1/n)$ , which is clearly continuous. If  $g$  is never 0 in  $]0; 1[$  then by Corollary 321  $g$  must be either strictly positive or strictly negative. But then

$$0 = f(0) - f(1) = \left(f(0) - f\left(\frac{1}{n}\right)\right) + \left(f\left(\frac{1}{n}\right) - f\left(\frac{2}{n}\right)\right) + \left(f\left(\frac{2}{n}\right) - f\left(\frac{3}{n}\right)\right) + \dots + \left(f\left(\frac{n-1}{n}\right) - f\left(\frac{n}{n}\right)\right).$$

The sum of each parenthesis on the right is strictly positive or strictly negative and hence never 0, a contradiction.

5.9.12 Consider the function  $f: ]0; 1[ \rightarrow ]0; 1[$ ,  $x \mapsto \frac{\sin \frac{2\pi x}{a}}{\sin \frac{2\pi}{a}} - x$ .

6.2.1 Observe that that

$$\frac{1}{x-1} - \frac{1}{x+1} = \frac{(x+1) - (x-1)}{(x-1)(x+1)} = \frac{2}{x^2 - 1}.$$

If  $f(x) = (x-1)^{-1}$  then

$$f'(x) = -1(x-1)^{-2}; f''(x) = (-1)(-2)(x-1)^{-3}; (-1)(-2)(-3)(x-1)^{-4}; \dots; f^{(100)}(x) = 100!(x-1)^{-101}.$$

Similarly, if  $g(x) = (x+1)^{-1}$  then

$$g'(x) = -1(x+1)^{-2}; g''(x) = (-1)(-2)(x+1)^{-3}; (-1)(-2)(-3)(x+1)^{-4}; \dots; g^{(100)}(x) = 100!(x+1)^{-101}.$$

Hence

$$\frac{d^{100}}{dx^{100}} \frac{2}{x^2 - 1} = f^{(100)}(x) - g^{(100)}(x) = 100!(x-1)^{-101} - 100!(x+1)^{-101}.$$

6.2.2 Use Leibniz's Rule and the observation that the third derivative of  $x \mapsto x^2$  is 0. Also  $(\sin x)^{(4n)} = \sin x$ ,  $(\sin x)^{(4n+2)} = -\sin x$ ,  $(\sin x)^{(4n+1)} = \cos x$ , and  $(\sin x)^{(4n+3)} = -\cos x$ . Then

$$\frac{d^{100}}{dx^{100}} x^2 \sin x = \binom{100}{0} x^2 (\sin x)^{(100)} + \binom{100}{1} (x^2)' (\sin x)^{(99)} + \binom{100}{2} (x^2)'' (\sin x)^{(98)} = x^2 \sin x - 200x \cos x - 9900 \sin x.$$

6.3.1 Put  $f(x) = x^5 - 2x^2 + x$ . Then  $f(0) = f(1) = 0$  and by Rolle's Theorem there is  $c \in ]0; 1[$  such that  $f'(c) = 5c^4 - 4c + 1 = 0$ .

6.3.2 Set

$$f(x) = a_0 x + \frac{a_1 x^2}{2} + \frac{a_2 x^3}{3} + \dots + \frac{a_n x^{n+1}}{n+1},$$

and use Rolle's Theorem.

6.3.4 Set  $g(x) = f(x)^2 f(1-x)$ . Since  $g(0) = g(1) = 0$ ,  $g$  satisfies the hypotheses of Rolle's Theorem. There is a  $c \in ]0; 1[$  such that

$$g'(c) = 0 \implies 2f'(c)f(c)f(1-c) - f(c)^2 f'(1-c) = 0.$$

Since by assumption  $f(c)f(1-c) \neq 0$  we must have, upon dividing by every term by  $f(c)^2 f(1-c)$ , the assertion.

6.3.5 For  $0 \leq k \leq n-1$ , consider the interval  $\left[\frac{k}{n}, \frac{k+1}{n}\right]$ . By the Mean Theorem, there are  $a_k \in \left[\frac{k}{n}, \frac{k+1}{n}\right]$  such that

$$f'(a_k) = \frac{f\left(\frac{k+1}{n}\right) - f\left(\frac{k}{n}\right)}{\frac{1}{n}} = n \left( f\left(\frac{k+1}{n}\right) - f\left(\frac{k}{n}\right) \right).$$

Summing from  $k=0$  to  $k=n-1$  and noting that the dextral side telescopes,

$$\sum_{k=0}^{n-1} f'(a_k) = n \sum_{k=0}^{n-1} \left( f\left(\frac{k+1}{n}\right) - f\left(\frac{k}{n}\right) \right) = n(f(1) - f(0)) = n.$$

6.3.6 Let  $k_i \in ]0; 1[$  be the smallest number such that  $f(k_i) = \frac{i}{n}$ ,  $1 \leq i \leq n-1$ . Put  $k_0 = 0, k_n = 1$ . The existence of the  $k_i$  is guaranteed by the Intermediate Value Theorem. Moreover, since the  $k_i$  are chosen to be the first time  $f$  is  $\frac{i}{n}$ , once again, by the Intermediate Value Theorem we must have

$$0 < k_1 < k_2 < \dots < k_{n-1} < 1.$$

Hence, by the Mean Value Theorem, there exists  $a_i \in ]k_i; k_{i+1}[$ ,  $0 \leq i \leq n-1$ , such that

$$f'(a_i) = \frac{f(k_{i+1}) - f(k_i)}{k_{i+1} - k_i} = \frac{1}{n(k_{i+1} - k_i)} \implies \frac{1}{f'(a_i)} = n(k_{i+1} - k_i).$$

Summing,

$$\sum_{k=0}^{n-1} \frac{1}{f'(a_k)} = n \sum_{k=0}^{n-1} (k_{i+1} - k_i) = n(k_n - k_0) = n.$$

6.4.2 We have  $f'(x) = x^x(\log x + 1)$  whence  $f'(x) = 0 \implies x = e^{-1}$ . Since  $f'(x) < 0$  for  $0 < x < e^{-1}$  and  $f'(x) > 0$  for  $x > e^{-1}$ ,  $x = e^{-1}$  is a local (relative) minimum. Thus  $f(x) \geq f(e^{-1}) = \left(\frac{1}{e}\right)^{1/e}$ .

6.5.3 Let  $0 < k < 1$ , and consider the function

$$f: \begin{array}{ccc} [0; +\infty[ & \rightarrow & \mathbb{R} \\ x & \mapsto & x^k - k(x-1) \end{array} .$$

Then  $0 = f'(x) = kx^{k-1} - k \Leftrightarrow x = 1$ . Since  $f''(x) = k(k-1)x^{k-2} < 0$  for  $0 < k < 1, x \geq 0, x = 1$  is a maximum point. Hence  $f(x) \leq f(1)$  for  $x \geq 0$ , that is  $x^k \leq 1 + k(x-1)$ . Letting  $k = \frac{1}{p}$  and  $x = \frac{a^p}{b^q}$  we deduce

$$\frac{a}{b^{q/p}} \leq 1 + \frac{1}{p} \left( \frac{a^p}{b^q} - 1 \right).$$

Rearranging gives

$$ab \leq b^{1+p/q} + \frac{a^p b^{1+p/q-p}}{p} - \frac{b^{1+p/q}}{p}$$

from where we obtain the inequality.

6.6.1 We have:

- Put  $f: \mathbb{R} \rightarrow \mathbb{R}, f(x) = e^{x-1} - x$ . Clearly  $f(1) = e^0 - 1 = 0$ . Now,

$$f'(x) = e^{x-1} - 1,$$

$$f''(x) = e^{x-1}.$$

If  $f'(x) = 0$  then  $e^{x-1} = 1$  implying that  $x = 1$ . Thus  $f$  has a single minimum point at  $x = 1$ . Thus for all real numbers  $x$

$$0 = f(1) \leq f(x) = e^{x-1} - x,$$

which gives the desired result.

- Easy Algebra!
- Easy Algebra!
- By the preceding results, we have

$$A_1 \leq \exp(A_1 - 1),$$

$$A_2 \leq \exp(A_2 - 1),$$

⋮

$$A_n \leq \exp(A_n - 1).$$

Since all the quantities involved are positive, we may multiply all these inequalities together, to obtain,

$$A_1 A_2 \cdots A_n \leq \exp(A_1 + A_2 + \cdots + A_n - n).$$

In view of the observations above, the preceding inequality is equivalent to

$$\frac{n^n G_n}{(a_1 + a_2 + \cdots + a_n)^n} \leq \exp(n - n) = e^0 = 1.$$

We deduce that

$$G_n \leq \left( \frac{a_1 + a_2 + \cdots + a_n}{n} \right)^n,$$

which is equivalent to

$$(a_1 a_2 \cdots a_n)^{1/n} \leq \frac{a_1 + a_2 + \cdots + a_n}{n}.$$

Now, for equality to occur, we need each of the inequalities  $A_k \leq \exp(A_k - 1)$  to hold. This occurs, in view of the preceding lemma, if and only if  $A_k = 1, \forall k$ , which translates into  $a_1 = a_2 = \cdots = a_n$ . This completes the proof.

6.7.1  $(\log \log x)^{\log x} = \exp((\log x)(\log \log \log x))$  and  $(\log x)^{\log \log x} = \exp((\log \log x)^2)$ . Now, lexicographically,

$$(\log \log x)^2 \ll (\log x)(\log \log \log x) \implies \exp((\log \log x)^2) \ll \exp((\log x)(\log \log \log x))$$

and thus  $(\log \log x)^{\log x}$  is faster.

7.1.1  $\Leftarrow$  This follows directly from Theorem 437.

$\implies$  If  $f$  is Riemann integrable, let  $\epsilon > 0$  and let  $\mathcal{P}' = \{a = y_0 < y_1 < \cdots < y_m = b\}$  be a partition with  $m + 1$  points such that

$$U(f, \mathcal{P}') - L(f, \mathcal{P}') < \frac{\epsilon}{2}.$$

As  $f$  is bounded, there is  $M > 0$  such that  $\forall x \in [a; b], |f(x)| \leq M$ . Take  $\delta = \frac{\epsilon}{8mM}$  and consider now an arbitrary partition  $\mathcal{P} = \{a = x_0 < x_1 < \cdots < x_n = b\}$  with norm  $|\mathcal{P}| < \delta$ . Put  $\mathcal{P}'' = \mathcal{P} \cup \mathcal{P}'$ . Arguing as in Theorem 434, we obtain

$$L(f, \mathcal{P}'') - L(f, \mathcal{P}) < 2mM|\mathcal{P}'| < 2mM\delta = \frac{\epsilon}{4}.$$

Since by Theorem 435  $L(f, \mathcal{P}') \leq L(f, \mathcal{P}'')$  we gather

$$L(f, \mathcal{P}') - L(f, \mathcal{P}) < \frac{\epsilon}{4}.$$

In a similar fashion we establish that

$$U(f, \mathcal{P}) - U(f, \mathcal{P}') < \frac{\epsilon}{4},$$

and upon assembling the inequalities,

$$U(f, \mathcal{P}) - L(f, \mathcal{P}) < U(f, \mathcal{P}') - L(f, \mathcal{P}') + \frac{\epsilon}{2} < \epsilon,$$

since we had assumed that  $U(f, \mathcal{P}') - L(f, \mathcal{P}') < \frac{\epsilon}{2}$ .

7.1.2  $\implies$  Assume  $f$  is Riemann-integrable. For  $\epsilon > 0$  let  $\delta > 0$  be chosen so that the conditions of Theorem ?? be fulfilled. By definition of a Riemann sum,

$$L(f, \mathcal{P}) \leq S(f, \mathcal{P}) \leq U(f, \mathcal{P}),$$

and therefore

$$U(f, \mathcal{P}) < L(f, \mathcal{P}) + \epsilon \leq \int_a^b f(x) dx + \epsilon = \int_a^b f(x) dx + \epsilon$$

and

$$L(f, \mathcal{P}) > U(f, \mathcal{P}) - \epsilon \geq \int_a^b f(x) dx - \epsilon = \int_a^b f(x) dx - \epsilon.$$

These inequalities give

$$\left| S(f, \mathcal{P}) - \int_a^b f(x) dx \right| < \epsilon,$$

whence  $\lim_{\|\mathcal{P}\| \rightarrow 0} S(f, \mathcal{P}) = \int_a^b f(x) dx.$

$\Leftarrow$  Suppose that  $\lim_{\|\mathcal{P}\| \rightarrow 0} S(f, \mathcal{P}) = L$ , existing and finite. Given  $\epsilon > 0$  there is  $\delta > 0$  such that  $\|\mathcal{P}\| < \delta$  implies

$$L - \frac{\epsilon}{3} < S(f, \mathcal{P}) < A + \frac{\epsilon}{3}. \tag{A.7}$$

Now, choose  $\mathcal{P} = \{a = x_0 < x_1 < \dots < x_n = b\}$ . By letting  $t_k$  range over  $[x_{k-1}; x_k]$  we gather, from (A.7)

$$L - \frac{\epsilon}{3} \leq L(f, \mathcal{P}) \leq U(f, \mathcal{P}) \leq L + \frac{\epsilon}{3},$$

whence

$$U(f, \mathcal{P}) - L(f, \mathcal{P}) \leq \frac{2}{3}\epsilon < \epsilon,$$

meaning that  $f$  is Riemann-integrable over  $[a; b]$  by Theorem 437. Thus

$$L(f, \mathcal{P}) \leq \int_a^b f(x) dx \leq U(f, \mathcal{P}),$$

and so  $\lim_{\|\mathcal{P}\| \rightarrow 0} S(f, \mathcal{P}) = \int_a^b f(x) dx.$

7.1.3  $\implies$  Let  $\mathcal{P} = \{a = x_0 < x_1 < \dots < x_n = b\}$  be a partition of  $[a; b]$ . Set

$$Z(f, \mathcal{P}) = \sum_{k=1}^n \omega(f, [x_{k-1}; x_k]) (x_k - x_{k-1}) = U(f, \mathcal{P}) - L(f, \mathcal{P}), \quad \Omega = \sup_{x \in [a; b]} f(x) - \inf_{x \in [a; b]} f(x).$$

Let

$$\delta = \sum_{k=1}^n (x_k - x_{k-1}) \chi_{\{x \in [a; b]; \omega(f, [x_{k-1}; x_k]) \geq \epsilon'\}}.$$

Then  $Z(f, \mathcal{P}) \geq \delta \epsilon'$ . Since we are assuming that  $f$  is Riemann-integrable, there exists a partition  $\mathcal{P}$  (by Theorem 437) such that

$$Z(f, \mathcal{P}) \leq \epsilon' \epsilon.$$

Thus we have  $\delta \epsilon' < \epsilon' \epsilon$  from where  $\delta < \epsilon$ .

$\Leftarrow$  Assume there is a partition  $\mathcal{P}$  for which  $\delta < \epsilon$ . In the intervals  $I = [x_{k-1}; x_k]$  where  $\omega(f, I) \geq \epsilon'$  the oscillation of  $f$  is at most  $\Omega$ , and in the remaining intervals (the sum of which is  $b - a - \delta$ , the oscillation is less than  $\epsilon'$ . Hence

$$Z(f, \mathcal{P}) \leq \delta \Omega + (b - a - \delta) \epsilon'.$$

Choose now

$$\epsilon' = \frac{\epsilon''}{2(b-a)}, \quad \delta = \frac{\epsilon''}{2\Omega}.$$

Since  $b - a - \delta \leq b - a$ ,

$$Z(f, \mathcal{P}) \leq \delta \Omega + (b - a - \delta) \epsilon' \leq \frac{\epsilon''}{2} + \frac{\epsilon''}{2} = \epsilon'',$$

whence  $f$  is Riemann-integrable by Theorem 437.

7.2.1  $\frac{8}{5}$

7.2.2

$$\begin{aligned} \int_0^3 x \lfloor x \rfloor dx &= \int_0^1 x \lfloor x \rfloor dx + \int_1^2 x \lfloor x \rfloor dx + \int_2^3 x \lfloor x \rfloor dx \\ &= 0 \int_0^1 x dx + 1 \int_1^2 x dx + 2 \int_2^3 x dx \\ &= \frac{x^2}{2} \Big|_0^1 + x^2 \Big|_1^2 \\ &= (2 - \frac{1}{2}) + (9 - 4) \\ &= \frac{13}{2}. \end{aligned}$$

7.2.3 We have

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{e^{x+h} - h - e^x}{h} = \lim_{h \rightarrow 0} \frac{e^{x+h} - e^x}{h} - \lim_{h \rightarrow 0} \frac{h}{h} = e^x - 1,$$

whence  $f(x) = e^x - x + C$ . Since  $3 = f(0) = e^0 - 0 + C \implies C = 2$ , we deduce that  $f(x) = e^x - x + 2$ .

7.2.4 Put  $I = \int_0^a \frac{1}{f(x)+1} dx$ . We have

$$I = \int_0^a \frac{1}{f(u)+1} du = \int_0^a \frac{f(u)f(a-u)}{f(u)+f(u)f(a-u)} du = \int_0^a \frac{f(a-u)}{1+f(a-u)} du = - \int_a^0 \frac{f(v)}{1+f(v)} dv = \int_0^a \frac{f(u)}{1+f(u)} du,$$

whence

$$2I = \int_0^a \frac{f(u)}{1+f(u)} du + \int_0^a \frac{f(a-u)}{1+f(a-u)} du = \int_0^a \frac{2+f(u)+f(a-u)}{2+f(u)+f(a-u)} du = a,$$

and so  $I = \frac{a}{2}$ .

7.2.5 Observe first that  $f(\mathbf{0} + \mathbf{0}) = f(\mathbf{0}) + f(\mathbf{0})$  and so  $f(\mathbf{0}) = \mathbf{0}$ . Integrate  $f(\mathbf{u} + \mathbf{y}) = f(\mathbf{u}) + f(\mathbf{y})$  for  $\mathbf{u} \in [0; \mathbf{x}]$  keeping  $\mathbf{y}$  constant, getting

$$\int_0^{\mathbf{x}} f(\mathbf{u} + \mathbf{y}) d\mathbf{u} = \int_0^{\mathbf{x}} f(\mathbf{u}) d\mathbf{u} + \int_0^{\mathbf{x}} f(\mathbf{y}) d\mathbf{u} = \int_0^{\mathbf{x}} f(\mathbf{u}) d\mathbf{u} + \mathbf{x}f(\mathbf{y}).$$

Also, by substitution,

$$\int_0^{\mathbf{x}} f(\mathbf{u} + \mathbf{y}) d\mathbf{u} = \int_{\mathbf{y}}^{\mathbf{y}+\mathbf{x}} f(\mathbf{u}) d\mathbf{u} = \int_0^{\mathbf{y}+\mathbf{x}} f(\mathbf{u}) d\mathbf{u} - \int_0^{\mathbf{y}} f(\mathbf{u}) d\mathbf{u}.$$

Hence

$$\mathbf{x}f(\mathbf{y}) = \int_0^{\mathbf{y}+\mathbf{x}} f(\mathbf{u}) d\mathbf{u} - \int_0^{\mathbf{y}} f(\mathbf{u}) d\mathbf{u} - \int_0^{\mathbf{x}} f(\mathbf{u}) d\mathbf{u}. \tag{A.8}$$

Exchanging  $\mathbf{x}$  and  $\mathbf{y}$ :

$$\mathbf{y}f(\mathbf{x}) = \int_0^{\mathbf{y}+\mathbf{x}} f(\mathbf{u}) d\mathbf{u} - \int_0^{\mathbf{x}} f(\mathbf{u}) d\mathbf{u} - \int_0^{\mathbf{y}} f(\mathbf{u}) d\mathbf{u}. \tag{A.9}$$

From (A.8) and (A.9) we gather that  $\mathbf{x}f(\mathbf{y}) = \mathbf{y}f(\mathbf{x})$ . If  $\mathbf{x}\mathbf{y} \neq 0$  then  $\frac{f(\mathbf{x})}{\mathbf{x}} = \frac{f(\mathbf{y})}{\mathbf{y}}$ . This means that for  $\frac{f(\mathbf{x})}{\mathbf{x}}$  is constant, and so for  $\mathbf{x} \neq 0$ ,  $f(\mathbf{x}) = c\mathbf{x}$  for some constant  $c$ . Since  $f(\mathbf{0}) = \mathbf{0}$ ,  $f(\mathbf{x}) = c\mathbf{x}$  for all  $\mathbf{x}$ . Taking  $\mathbf{x} = \mathbf{1}$ ,  $f(\mathbf{1}) = c$ .

7.2.7 We have

$$\begin{aligned} \int_{-1}^2 |x^2 - 1| dx &= \int_{-1}^1 (1 - x^2) dx + \int_1^2 (x^2 - 1) dx \\ &= \left(x - \frac{x^3}{3}\right) \Big|_{-1}^1 + \left(\frac{x^3}{3} - x\right) \Big|_1^2 \\ &= 2\left(1 - \frac{1}{3}\right) + \left(\frac{8}{3} - 2\right) - \left(\frac{1}{3} - 1\right) \\ &= \frac{4}{3} + \frac{2}{3} - \frac{2}{3} \\ &= \frac{8}{3} \end{aligned}$$

7.2.16 Put  $u = \sqrt{x^2 - 1}$ ;  $u^2 = x^2 - 1$  so that  $2u du = 2x dx$  and  $\frac{dx}{x} = \frac{x dx}{x^2} = \frac{u du}{u^2 + 1}$ . Thus

$$\int \frac{1}{x\sqrt{x^2 - 1}} dx = \int \frac{u}{(u^2 + 1)u} du = \int \frac{1}{u^2 + 1} du = \arctan u + C = \arctan \sqrt{x^2 - 1} + C.$$

7.2.17 Put  $u = \sqrt{x+1}$ ;  $u^2 = x+1$ ; from where  $dx = 2u du$ . Whence

$$\int \frac{1}{1 + \sqrt{x+1}} dx = \int \frac{2u}{1+u} du = \int \left(2 - \frac{2}{1+u}\right) du = 2u - 2\log|1+u| + C = 2\sqrt{1+x} - 2\log|1 + \sqrt{1+x}| + C.$$

7.2.18 Put  $x = u^6$ ;  $dx = 6u^5 du$ , giving

$$\begin{aligned} \int \frac{x^{1/2}}{x^{1/2} - x^{1/3}} dx &= \int \frac{(u^3)(6u^5)}{u^3 - u^2} du \\ &= \int \frac{6u^6}{u-1} du \\ &= 6 \int \left(u^5 + u^4 + u^3 + u^2 + u + 1 + \frac{1}{u-1}\right) du \\ &= 6 \left(\frac{u^6}{6} + \frac{u^5}{5} + \frac{u^4}{4} + \frac{u^3}{3} + \frac{u^2}{2} + u + \log|u-1|\right) + C \\ &= x + \frac{6x^{5/6}}{5} + \frac{3x^{2/3}}{2} + 2x^{1/2} + 3x^{1/3} + 6x^{1/6} + 6\log|x^{1/6} - 1| + C. \end{aligned}$$

7.2.19 Put  $u^2 = a^x + 1$ ;  $2u du = (\log a) a^x dx$  and so

$$\int \frac{a^{2x}}{\sqrt{a^x + 1}} dx = \int \frac{2u(u^2 - 1)}{u \log a} du = \int \frac{2u^2 - 2}{\log a} du = \frac{2u^3}{3 \log a} - \frac{2u}{\log a} + C = \frac{2(a^x + 1)^{3/2}}{3 \log a} - \frac{2(a^x + 1)^{1/2}}{\log a} + C.$$

7.2.20 Observe that  $(e^x - e^{-x})^2 = (e^{-x}(e^{2x} - 1))^2 = e^{-2x}(e^{2x} - 1)^2$ , and so

$$\int \frac{1}{(e^x - e^{-x})^2} dx = \int \frac{e^{2x}}{(e^{2x} - 1)^2} dx = \int \frac{1}{2u^2} du = -\frac{1}{2u} + C = -\frac{1}{2(e^{2x} - 1)} + C,$$

on putting  $u = e^{2x} - 1$ .

7.2.21 We have

$$\begin{aligned} \int_1^5 \frac{[x]}{x} dx &= \int_1^2 \frac{[x]}{x} dx + \int_2^3 \frac{[x]}{x} dx + \int_3^4 \frac{[x]}{x} dx + \int_4^5 \frac{[x]}{x} dx \\ &= \int_1^2 \frac{1}{x} dx + \int_2^3 \frac{2}{x} dx + \int_3^4 \frac{3}{x} dx + \int_4^5 \frac{4}{x} dx \\ &= (\log 2 - \log 1) + 2(\log 3 - \log 2) + 3(\log 4 - \log 3) + 4(\log 5 - \log 4) \\ &= 4 \log(5) - 3 \log(2) - \log(3). \end{aligned}$$

7.2.22 Put  $u = e^x$ , etc.

$$\int e^{e^x + x} dx = \int e^x e^{e^x} dx = \int e^{e^x} de^x = e^{e^x} + C$$



7.2.23 Put  $u = \log(\cos x)$ , etc.

$$\int \tan x \log(\cos x) dx = \int (\log(\cos x)) d(-\log(\cos x)) = -\frac{(\log(\cos x))^2}{2} + C$$

7.2.24 Put  $u = \log \log x$ , etc.

$$\int \frac{\log \log x}{x \log x} dx = \int \log \log x d(\log \log x) = \frac{\log \log x}{2} + C$$

7.2.25 Carry out the long division.

$$\int \frac{x^{18}-1}{x^3-1} dx = \int (x^{15} + x^{12} + x^9 + x^6 + x^3 + 1) dx = \frac{x^{16}}{16} + \frac{x^{13}}{13} + \frac{x^{10}}{10} + \frac{x^7}{7} + \frac{x^4}{4} + x + C$$

7.2.26 After an algebraic trick, put  $u = 1 + x^{-7}$ , etc.

$$\int \frac{1}{x^8+x} dx = \int \frac{x^{-8}}{1+x^{-7}} dx = -\frac{1}{7} \int \frac{d(1+x^{-7})}{1+x^{-7}} = -\frac{1}{7} \log|1+x^{-7}| + C$$

7.2.27 Put  $u = 2^x + 1$ 

$$\int \frac{2^x 2^x}{2^x + 1} dx = \frac{1}{\log 2} \int \frac{2^x}{2^x + 1} d(2^x + 1) = \frac{1}{\log 2} \int \frac{u-1}{u} du = \frac{1}{\log 2} (u - \log|u|) + C = \frac{1}{\log 2} (2^x + 1 - \log|2^x + 1|) + C$$

7.2.28 Put  $u = x + 1$ . Then  $x^2 = (u-1)^2 = u^2 - 2u + 1$ , and hence

$$\begin{aligned} \int \frac{x^2}{(x+1)^{10}} dx &= \int \frac{u^2 - 2u + 1}{u^{10}} du \\ &= \int u^{-8} - 2u^{-9} + u^{-10} du \\ &= \frac{u^{-7}}{7} + \frac{u^{-8}}{8} - \frac{u^{-9}}{9} + C \\ &= \frac{(x+1)^{-7}}{7} + \frac{(x+1)^{-8}}{8} - \frac{(x+1)^{-9}}{9} + C \end{aligned}$$

7.2.29 Algebraic trick, and then  $u = e^{-x} + 1$ , etc.

$$\int \frac{1}{1+e^x} dx = \int \frac{e^{-x}}{e^{-x}+1} dx = -\int \frac{1}{e^{-x}+1} d(e^{-x}+1) = -\log|e^{-x}+1| + C$$

7.2.30

$$\int \frac{1}{1-\sin x} dx = \int \frac{1+\sin x}{1-\sin^2 x} dx = \int \frac{1+\sin x}{\cos^2 x} dx = \int \sec^2 x + \sec x \tan x dx = \tan x + \sec x + C$$

7.2.31

$$\begin{aligned} \int \sqrt{1+\sin 2x} dx &= \int \sqrt{\sin^2 x + 2 \sin x \cos x + \cos^2 x} dx \\ &= \int \sqrt{(\sin x + \cos x)^2} dx \\ &= \int |\sin x + \cos x| dx \\ &= \mp \cos x \pm \sin x + C \end{aligned}$$

7.2.32 Put  $u = x^2$ , etc.

$$\int \frac{x}{\sqrt{1-(x^2)^2}} dx = \frac{1}{2} \int \frac{1}{\sqrt{1-u^2}} du = \frac{1}{2} \arcsin u + C = \frac{1}{2} \arcsin x^2 + C$$

7.2.33 We have

$$\begin{aligned} \int \sec^4 x dx &= \int \sec^2 x (\tan^2 x + 1) dx \\ &= \int \sec^2 x \tan^2 x dx + \int \sec^2 x dx \\ &= \int (\tan x)^2 d(\tan x) + \int \sec^2 x dx \\ &= \frac{\tan^3 x}{3} + \tan x + C. \end{aligned}$$

7.2.34 We have

$$\begin{aligned} \int \sec^5 x dx &= \int \sec^3 x \sec^2 x dx \\ &= \int \sec^3 x d(\tan x) \\ &= \sec^3 x \tan x - \int \tan x d(\sec^3 x) \\ &= \sec^3 x \tan x - 3 \int \tan^2 x \sec^2 x dx \\ &= \sec^3 x \tan x - 3 \int (\sec^2 x - 1) \sec^2 x dx \\ &= \sec^3 x \tan x - 3 \int \sec^4 x dx + 3 \int \sec^2 x dx \end{aligned}$$

The above implies that

$$\begin{aligned} \int \sec^5 x dx &= \frac{\tan x \sec^3 x}{4} + \frac{3}{4} \int \sec^3 x dx \\ &= \frac{\tan x \sec^3 x}{4} + \frac{3 \tan x \sec x}{8} + \frac{3}{8} \log|\sec x + \tan x| + C, \end{aligned}$$

upon recalling from class that

$$\int \sec^3 x dx = \frac{\tan x \sec x}{2} + \frac{1}{2} \log|\sec x + \tan x| + C$$

7.2.35 First put  $t = x^{1/3}$ , then  $t^3 = x \implies 3t^2 dt = dx$ . Thus

$$\begin{aligned} \int e^{x^{1/3}} dx &= \int 3t^2 e^t dt \\ &= 3t^2 e^t - 6t e^t - 6e^t + C \\ &= 3x^{2/3} e^{x^{1/3}} - 6x^{1/3} e^{x^{1/3}} - 6e^{x^{1/3}} + C, \end{aligned}$$

where the penultimate step results from tabular integration by parts.

7.2.36 We have

$$\begin{aligned} \int \log(x^2 + 1) dx &= x \log(x^2 + 1) - \int x d(\log(x^2 + 1)) \\ &= x \log(x^2 + 1) - 2 \int \frac{x^2}{x^2 + 1} dx \\ &= x \log(x^2 + 1) - 2 \int \frac{x^2 + 1 - 1}{x^2 + 1} dx \\ &= x \log(x^2 + 1) - 2 \int \left(1 - \frac{1}{x^2 + 1}\right) dx \\ &= x \log(x^2 + 1) - 2(x - \arctan x) + C \end{aligned}$$

7.2.37 Put

$$I = \int x e^x \cos x := (Ax + B)e^x \cos x + (Cx + D)e^x \sin x + K.$$

Differentiating both sides,

$$x e^x \cos x = A e^x \cos x + (Ax + B)e^x \cos x - (Ax + B)e^x \sin x + C e^x \sin x + (Cx + D)e^x \sin x + (Cx + D)e^x \cos x.$$

Equating coefficients,

$$\begin{aligned} x e^x \cos x &: 1 = A + C \\ x e^x \sin x &: 0 = -A + C \\ e^x \cos x &: 0 = A + B + D \\ e^x \sin x &: 0 = -B + C + D \end{aligned}$$

From the first two equations  $C = \frac{1}{2}, A = \frac{1}{2}$ . Then the third and fourth equations become  $-\frac{1}{2} = B + D; -\frac{1}{2} = -B + D$ , whence  $D = -\frac{1}{2}$ , and  $B = 0$ . We conclude that

$$\int x e^x \cos x = \frac{x}{2} e^x \cos x + \left(\frac{x-1}{2}\right) e^x \sin x + K.$$

7.2.38 We will do this one two ways: first, by making the substitution

$$t = \log x \Rightarrow e^t = x \Rightarrow e^t dt = dx.$$

Observe also that  $x^{2/3} = e^{2t/3}$ . Then

$$\begin{aligned} \int x^{2/3} \log x dx &= \int t e^{2t/3} e^t dt \\ &= \frac{3t}{5} e^{5t/3} - \frac{9}{25} e^{5t/3} + C \\ &= \frac{3(\log x)}{5} x^{5/3} - \frac{9}{25} x^{5/3} + C. \end{aligned}$$

Alter: By directly integrating by parts,

$$\begin{aligned} \int x^{2/3} \log x dx &= \int \log x d\left(\frac{3x^{5/3}}{5}\right) \\ &= \frac{3x^{5/3}}{5} \log x - \frac{3}{5} \int x^{5/3} d(\log x) \\ &= \frac{3(\log x)}{5} x^{5/3} - \frac{3}{5} \int x^{2/3} dx \\ &= \frac{3(\log x)}{5} x^{5/3} - \frac{9}{25} x^{5/3} + C, \end{aligned}$$

as before.

7.2.39 This integral can be done multiple ways. For example, you may integrate by parts directly and then "solve" for the integral. Another way is the following. Start by putting

$$t = \log x \Rightarrow e^t = x \Rightarrow e^t dt = dx.$$

Then

$$\int \sin(\log x) dx = \int e^t \sin t dt,$$

an integral that we found in class. We will find it again, using a method similar of problem 7.2.37. Put

$$I = \int e^t \cos t dt := A e^t \cos t + B e^t \sin t + K.$$

Differentiating both sides

$$e^t \cos t = A e^t \cos t - A e^t \sin t + B e^t \sin t + B e^t \cos t.$$

Equating coefficients,

$$\begin{aligned} e^t \cos t &: 1 = A + B \\ e^t \sin t &: 0 = -A + B \end{aligned}$$

and so  $A = B = \frac{1}{2}$ . We have thus

$$\begin{aligned} \int \sin(\log x) dx &= \int e^t \sin t dt \\ &= \frac{1}{2} e^t \cos t + \frac{1}{2} e^t \sin t + K \\ &= \frac{1}{2} x \cos \log x + \frac{1}{2} x \sin \log x + K. \end{aligned}$$

7.2.40 Put  $t = \log \log x \Rightarrow e^{e^t} = x \Rightarrow e^t e^{e^t} dt = dx$ . Hence

$$\begin{aligned} \int \frac{\log \log x}{x} dx &= \int \frac{t e^t e^{e^t}}{e^{e^t}} dt \\ &= t e^t - e^t + C \\ &= (\log x)(\log \log x) - (\log x) + C, \end{aligned}$$

where the penultimate equality follows from a tabular integration by parts.

7.2.41 Observe that

$$\int \sec x dx = \int \frac{\sec x \tan x + \sec^2 x}{\tan x + \sec x} dx = \int d(\log(\tan x + \sec x)) = \log(\tan x + \sec x) + C,$$

For the second way, simple algebra will yield the identity. We have

$$\begin{aligned} \int \sec x dx &= \int \frac{\cos x}{2(1 + \sin x)} dx + \int \frac{\cos x}{2(1 - \sin x)} dx \\ &= \frac{1}{2} \log|1 + \sin x| - \frac{1}{2} \log|1 - \sin x| + C \\ &= \frac{1}{2} \log \left| \frac{1 + \sin x}{1 - \sin x} \right| + C \end{aligned}$$

For the third way, we have

$$\begin{aligned} \int \csc x dx &= \int \frac{1}{\sin x} dx \\ &= \int \frac{1}{2 \sin \frac{x}{2} \cos \frac{x}{2}} dx \\ &= \int \frac{\cos \frac{x}{2}}{2 \sin \frac{x}{2} \cos^2 \frac{x}{2}} dx \\ &= \int \frac{\sec^2 \frac{x}{2}}{2 \tan \frac{x}{2}} dx \\ &\stackrel{u = \tan \frac{x}{2}}{=} \int \frac{du}{u} \\ &= \log \left| \tan \frac{x}{2} \right| + C. \end{aligned}$$

Thus

$$\int \sec x dx = \int \csc\left(\frac{\pi}{2} + x\right) dx = \int \csc\left(\frac{\pi}{2} + x\right) d\left(\frac{\pi}{2} + x\right) = \log \left| \tan\left(\frac{\pi}{4} + \frac{x}{2}\right) \right| + C.$$

7.2.42 Putting  $t = \arcsin x$  we have

$$\sin t = x \implies \cos t dt = dx,$$

whence

$$\begin{aligned} \int (\arcsin x)^2 dx &= \int t^2 \cos t dt \\ &= t^2 \sin t + 2t \cos t - 2 \sin t + C \\ &= (\arcsin x)^2 x + 2(\arcsin x) \cos(\arcsin x) - 2x + C \\ &= (\arcsin x)^2 x + 2(\arcsin x) \sqrt{1 - x^2} - 2x + C \end{aligned}$$

7.2.43 We have

$$\begin{aligned} \int \frac{dx}{\sqrt{x+1} + \sqrt{x-1}} &= \int \frac{(\sqrt{x+1} - \sqrt{x-1}) dx}{2} \\ &= \frac{1}{3} (x+1)^{3/2} - \frac{1}{3} (x-1)^{3/2} + C \end{aligned}$$

7.2.44 We have

$$\begin{aligned} \int x \arctan x dx &= \int \arctan x d\left(\frac{x^2}{2}\right) \\ &= \frac{x^2}{2} \arctan x - \int \frac{x^2}{2} d(\arctan x) \\ &= \frac{x^2}{2} \arctan x - \int \frac{1}{2} \frac{x^2}{1+x^2} dx \\ &= \frac{x^2}{2} \arctan x - \int \frac{1}{2} \frac{x^2 + 1 - 1}{1+x^2} dx \\ &= \frac{x^2}{2} \arctan x - \frac{x}{2} + \frac{1}{2} \arctan x + C \end{aligned}$$

7.2.45 Put  $u = \sqrt{\tan x}$  and so  $u^2 = \tan x$ ,  $2u du = \sec^2 x dx = (\tan^2 x + 1) dx = (u^4 + 1) dx$ . Hence the integral becomes

$$\int \sqrt{\tan x} dx = 2 \int \frac{u^2}{u^4 + 1} du.$$

To decompose the above fraction into partial fractions observe (Sophie Germain's trick) that  $u^4 + 1 = u^4 + 2u^2 + 1 - 2u^2 = (u^2 + u\sqrt{2} + 1)(u^2 - u\sqrt{2} + 1)$  and hence

$$\begin{aligned} \int \sqrt{\tan x} dx &= 2 \int \frac{u^2}{u^4 + 1} du \\ &= -\frac{\sqrt{2}}{2} \int \frac{u}{u^2 + u\sqrt{2} + 1} du + \frac{\sqrt{2}}{2} \int \frac{u}{u^2 - u\sqrt{2} + 1} du \\ &= -\frac{\sqrt{2}}{4} \log(u^2 + u\sqrt{2} + 1) + \frac{\sqrt{2}}{4} \log(u^2 - u\sqrt{2} + 1) + \frac{\sqrt{2}}{2} \arctan(\sqrt{2}u + 1) - \frac{\sqrt{2}}{2} \arctan(-\sqrt{2}u + 1) + C \\ &= -\frac{\sqrt{2}}{4} \log(\tan x + \sqrt{2}\tan x + 1) + \frac{\sqrt{2}}{4} \log(\tan x - \sqrt{2}\tan x + 1) \\ &\quad + \frac{\sqrt{2}}{2} \arctan(\sqrt{2}\tan x + 1) - \frac{\sqrt{2}}{2} \arctan(-\sqrt{2}\tan x + 1) + C \end{aligned}$$

7.2.46 Put

$$\frac{2x+1}{x^2(x-1)} = \frac{A}{x} + \frac{B}{x^2} + \frac{C}{x-1} \implies 2x+1 = Ax(x-1) + B(x-1) + Cx^2.$$

Letting  $x = 1$  we get  $3 = C$ . Letting  $x = 0$  we get  $1 = -B \implies B = -1$ . To get  $A$  observe that equating the coefficients of  $x^2$  on both sides we get  $0 = A + C$ , whence  $A = -3$ . Thus

$$\begin{aligned} \int \frac{2x+1}{x^2(x-1)} dx &= -3 \int \frac{1}{x} dx - \int \frac{1}{x^2} dx + 3 \int \frac{1}{x-1} dx \\ &= -3 \log|x| + \frac{1}{x} + 3 \log|x-1| + C \\ &= 3 \log \left| \frac{x-1}{x} \right| + \frac{1}{x} + C. \end{aligned}$$

7.2.47 Integrating by parts,

$$\begin{aligned}
 \int \log(x + \sqrt{x}) dx &= x \log(x + \sqrt{x}) - \int x d \log(x + \sqrt{x}) \\
 &= x \log(x + \sqrt{x}) - \int \frac{x(1 + \frac{1}{2\sqrt{x}})}{x + \sqrt{x}} dx \\
 &= x \log(x + \sqrt{x}) - \int \left(1 - \frac{1}{2} \frac{\sqrt{x}}{x + \sqrt{x}}\right) dx \\
 &= x \log(x + \sqrt{x}) - x + \frac{1}{2} \int \frac{\sqrt{x}}{x + \sqrt{x}} dx \\
 &\stackrel{u=\sqrt{x}}{=} x \log(x + \sqrt{x}) - x + \int \frac{u^2}{u^2 + u} du \\
 &\stackrel{u=\sqrt{x}}{=} x \log(x + \sqrt{x}) - x + \int \left(1 - \frac{1}{u+1}\right) du \\
 &= x \log(x + \sqrt{x}) - x + u - \log(u+1) + C \\
 &= x \log(x + \sqrt{x}) - x + \sqrt{x} - \log(\sqrt{x} + 1) + C
 \end{aligned}$$

7.2.48 We use Sophie Germain's trick to factor

$$x^4 + 1 = x^4 + 2x^2 + 1 - 2x^2 = (x^2 + 1)^2 - 2x^2 = (x^2 - \sqrt{2}x + 1)(x^2 + \sqrt{2}x + 1),$$

and seek the partial fraction decomposition

$$\frac{1}{x^4 + 1} = \frac{Ax + B}{x^2 - \sqrt{2}x + 1} + \frac{Cx + D}{x^2 + \sqrt{2}x + 1} \implies 1 = (Ax + B)(x^2 + \sqrt{2}x + 1) + (Cx + D)(x^2 - \sqrt{2}x + 1).$$

Equating coefficients

$$\begin{aligned}
 x^3 &: 0 = A + C \\
 x^2 &: 0 = B + D + \sqrt{2}(A - C) \\
 x &: 0 = A + C + \sqrt{2}(B - D) \\
 x^0 &: 1 = B + D
 \end{aligned}$$

From the first and third equation it follows that  $A = -C$  and that  $B = D$ . From the fourth equation  $B = D = \frac{1}{2}$  and from the second equation  $A = -\frac{1}{2\sqrt{2}} = -C$ . Hence we must integrate

$$\begin{aligned}
 \int \frac{1}{x^4 + 1} dx &= \int \frac{\sqrt{2}x + 2}{4(x^2 + \sqrt{2}x + 1)} dx - \int \frac{\sqrt{2}x - 2}{4(x^2 - \sqrt{2}x + 1)} dx \\
 &= \frac{\sqrt{2}}{8} \int \frac{2x + \sqrt{2}}{x^2 + \sqrt{2}x + 1} dx + \frac{1}{4} \int \frac{1}{x^2 + \sqrt{2}x + 1} dx - \frac{\sqrt{2}}{8} \int \frac{2x + \sqrt{2}}{x^2 - \sqrt{2}x + 1} dx + \frac{1}{4} \int \frac{1}{x^2 - \sqrt{2}x + 1} dx \\
 &= \frac{\sqrt{2}}{8} \log(x^2 + x\sqrt{2} + 1) - \frac{\sqrt{2}}{8} \log(x^2 - x\sqrt{2} + 1) + \frac{1}{2} \int \frac{dx}{(x\sqrt{2} + 1)^2 + 1} + \frac{1}{2} \int \frac{dx}{(-x\sqrt{2} + 1)^2 + 1} \\
 &= \frac{\sqrt{2}}{8} \log(x^2 + x\sqrt{2} + 1) - \frac{\sqrt{2}}{8} \log(x^2 - x\sqrt{2} + 1) + \frac{\sqrt{2}}{4} \arctan(x\sqrt{2} + 1) - \frac{\sqrt{2}}{4} \arctan(-x\sqrt{2} + 1) + C
 \end{aligned}$$

7.2.49 We begin by observing that

$$\frac{1}{x^3 + 1} = \frac{A}{x + 1} + \frac{Bx + C}{x^2 - x + 1} \implies 1 = A(x^2 - x + 1) + (Bx + C)(x + 1).$$

Letting  $x = -1$  we obtain  $1 = 3A \implies A = \frac{1}{3}$ . Letting  $x = 0$  we obtain  $1 = A + C \implies C = 1 - A = \frac{2}{3}$ . Finally, we must have  $A + B = 0$ , since the coefficient of  $x^2$  must be zero, thus  $B = -\frac{1}{3}$ . We must then integrate

$$\begin{aligned}
 \int \frac{dx}{3(x+1)} - \int \frac{x-2}{3(x^2-x+1)} dx &= \frac{1}{3} \log|x+1| - \int \frac{x-\frac{1}{2}}{3(x-\frac{1}{2})^2 + \frac{3}{4}} + \frac{1}{2} \int \frac{1}{(x-\frac{1}{2})^2 + \frac{3}{4}} \\
 &= \frac{1}{3} \log|x+1| - \frac{1}{6} \log|(x-\frac{1}{2})^2 + \frac{3}{4}| + \frac{3}{4} + \frac{2}{3} \int \frac{1}{\frac{4}{3}(x-\frac{1}{2})^2 + 1} \\
 &= \frac{1}{3} \log|x+1| - \frac{1}{6} \log|(x-\frac{1}{2})^2 + \frac{3}{4}| + \frac{3}{4} + \frac{2}{3} \cdot \frac{\sqrt{3}}{2} \arctan(x - \frac{1}{2}) \\
 &= \frac{1}{3} \log|x+1| - \frac{1}{6} \log|x^2 - x + 1| + \frac{\sqrt{3}}{3} \arctan \frac{2}{\sqrt{3}}(x - \frac{1}{2})
 \end{aligned}$$

8.8.1

8.8.2

8.8.3

8.8.4

# Bibliography

[Apo] Apostol, T. M., **Calculus**, Vol 1 & 2, 2nd ed., Waltham: Xerox, 1967.

[Har] Hardy, G. H., **Pure Mathematics**, 10th ed., New York: Cambridge University Press, 1952.

[Kla] Klambauer, Gabriel, **Aspects of Calculus**, New York: Springer-Verlag, 1986.

[Lan] Landau, E., **Differential and Integral Calculus**, New York, Chelsea Publishing Company, 1950.

[Olm] Olmstead, J. M. H., **Calculus with Analytic Geometry**, Vol 1 & 2, New York: Appleton-Century-Crofts, 1966.

[Spi] Spivak, Michael **Calculus**, 3rd ed., Houston, Texas: Publish or Perish, Inc., 1994.