


Linear Algebra Notes

David A. SANTOS
dsantos@ccp.edu



Contents

Preface	iii	4.5 Vectors in \mathbb{R}^3	86
To the Student	v	4.6 Planes and Lines in \mathbb{R}^3	91
1 Preliminaries	1	4.7 \mathbb{R}^n	95
1.1 Sets and Notation	1	5 Vector Spaces	100
1.2 Partitions and Equivalence Relations	4	5.1 Vector Spaces	100
1.3 Binary Operations	6	5.2 Vector Subspaces	103
1.4 \mathbb{Z}_n	9	5.3 Linear Independence	105
1.5 Fields	12	5.4 Spanning Sets	107
1.6 Functions	14	5.5 Bases	110
5.6 Coordinates	114		
2 Matrices and Matrix Operations	17	6 Linear Transformations	120
2.1 The Algebra of Matrices	17	6.1 Linear Transformations	120
2.2 Matrix Multiplication	20	6.2 Kernel and Image of a Linear Transformation	122
2.3 Trace and Transpose	24	6.3 Matrix Representation	127
2.4 Special Matrices	26		
2.5 Matrix Inversion	32	7 Determinants	133
2.6 Block Matrices	40	7.1 Permutations	133
2.7 Rank of a Matrix	41	7.2 Cycle Notation	136
2.8 Rank and Invertibility	50	7.3 Determinants	141
		7.4 Laplace Expansion	150
3 Linear Equations	57	7.5 Determinants and Linear Systems	156
3.1 Definitions	57		
3.2 Existence of Solutions	61	8 Eigenvalues and Eigenvectors	157
3.3 Examples of Linear Systems	63	8.1 Similar Matrices	157
4 \mathbb{R}^2, \mathbb{R}^3 and \mathbb{R}^n	67	8.2 Eigenvalues and Eigenvectors	158
4.1 Points and Bi-points in \mathbb{R}^2	67	8.3 Diagonalisability	162
4.2 Vectors in \mathbb{R}^2	70	8.4 The Minimal Polynomial	165
4.3 Dot Product in \mathbb{R}^2	76		
4.4 Lines on the Plane	81	A Answers and Hints	167
		Answers and Hints	167

Preface

These notes started during the Spring of 2002, when John MAJEWICZ and I each taught a section of Linear Algebra. I would like to thank him for numerous suggestions on the written notes.

The students of my class were: Craig BARIBAULT, Chun CAO, Jacky CHAN, Pho DO, Keith HARMON, Nicholas SELVAGGI, Sanda SHWE, and Huong VU.

John's students were David HERNÁNDEZ, Adel JAILILI, Andrew KIM, Jong KIM, Abdelmounaim LAAYOUNI, Aju MATHEW, Nikita MORIN, Thomas NEGRÓN, Latoya ROBINSON, and Saem SOEURN.

Linear Algebra is often a student's first introduction to abstract mathematics. Linear Algebra is well suited for this, as it has a number of beautiful but elementary and easy to prove theorems. My purpose with these notes is to introduce students to the concept of proof in a gentle manner.

David A. Santos dsantos@ccp.edu

Legal Notice

This material may be distributed only subject to the terms and conditions set forth in the Open Publication License, version 1.0 or later (the latest version is presently available at

<http://www.opencontent.org/openpub/>.

THIS WORK IS LICENSED AND PROVIDED "AS IS" WITHOUT WARRANTY OF ANY KIND, EXPRESS OR IMPLIED, INCLUDING, BUT NOT LIMITED TO, THE IMPLIED WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE OR A WARRANTY OF NON-INFRINGEMENT.

THIS DOCUMENT MAY NOT BE SOLD FOR PROFIT OR INCORPORATED INTO COMMERCIAL DOCUMENTS WITHOUT EXPRESS PERMISSION FROM THE AUTHOR(S). THIS DOCUMENT MAY BE FREELY DISTRIBUTED PROVIDED THE NAME OF THE ORIGINAL AUTHOR(S) IS(ARE) KEPT AND ANY CHANGES TO IT NOTED.

To the Student

These notes are provided for your benefit as an attempt to organise the salient points of the course. They are a *very terse* account of the main ideas of the course, and are to be used mostly to refer to central definitions and theorems. The number of examples is minimal, and here you will find few exercises. The *motivation* or informal ideas of looking at a certain topic, the ideas linking a topic with another, the worked-out examples, etc., are given in class. Hence these notes are not a substitute to lectures: **you must always attend to lectures**. The order of the notes may not necessarily be the order followed in the class.

There is a certain algebraic fluency that is necessary for a course at this level. These algebraic prerequisites would be difficult to codify here, as they vary depending on class response and the topic lectured. If at any stage you stumble in Algebra, seek help! I am here to help you!

Tutoring can sometimes help, but bear in mind that whoever tutors you may not be familiar with my conventions. Again, I am here to help! On the same vein, other books may help, but the approach presented here is at times unorthodox and finding alternative sources might be difficult.

Here are more recommendations:

- Read a section before class discussion, in particular, read the definitions.
- Class provides the informal discussion, and you will profit from the comments of your classmates, as well as gain confidence by providing your insights and interpretations of a topic. **Don't be absent!**
- Once the lecture of a particular topic has been given, take a fresh look at the notes of the lecture topic.
- Try to understand a single example well, rather than ill-digest multiple examples.
- Start working on the distributed homework ahead of time.
- **Ask questions during the lecture.** There are two main types of questions that you are likely to ask.
 1. *Questions of Correction:* *Is that a minus sign there?* If you think that, for example, I have missed out a minus sign or wrote **P** where it should have been **Q**,¹ then by all means, ask. No one likes to carry an error till line XLV because the audience failed to point out an error on line I. Don't wait till the end of the class to point out an error. Do it when there is still time to correct it!
 2. *Questions of Understanding:* *I don't get it!* Admitting that you do not understand something is an act requiring utmost courage. But if you don't, it is likely that many others in the audience also don't. On the same vein, if you feel you can explain a point to an inquiring classmate, I will allow you time in the lecture to do so. The best way to ask a question is something like: "How did you get from the second step to the third step?" or "What does it mean to complete the square?" Asseverations like "I don't understand" do not help me answer your queries. If I consider that you are asking the same questions too many times, it may be that you need extra help, in which case we will settle what to do outside the lecture.
- Don't fall behind! The sequence of topics is closely interrelated, with one topic leading to another.
- The use of calculators is allowed, especially in the occasional lengthy calculations. However, when graphing, you will need to provide algebraic/analytic/geometric support of your arguments. The questions on assignments and exams will be posed in such a way that it will be of no advantage to have a graphing calculator.
- Presentation is critical. Clearly outline your ideas. When writing solutions, outline major steps and write in complete sentences. As a guide, you may try to emulate the style presented in the scant examples furnished in these notes.

¹My doctoral adviser used to say "I said **A**, I wrote **B**, I meant **C** and it should have been **D**!"

Chapter 1

Preliminaries

1.1 Sets and Notation

1 Definition We will mean by a *set* a collection of well defined members or *elements*.

2 Definition The following sets have special symbols.

$\mathbb{N} = \{0, 1, 2, 3, \dots\}$ denotes the set of natural numbers.

$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$ denotes the set of integers.

\mathbb{Q} denotes the set of rational numbers.

\mathbb{R} denotes the set of real numbers.

\mathbb{C} denotes the set of complex numbers.

\emptyset denotes the empty set.

3 Definition (Implications) The symbol \implies is read “implies”, and the symbol \iff is read “if and only if.”

4 Example Prove that between any two rational numbers there is always a rational number.

Solution: Let $(a, c) \in \mathbb{Z}^2$, $(b, d) \in (\mathbb{N} \setminus \{0\})^2$, $\frac{a}{b} < \frac{c}{d}$. Then $da < bc$. Now

$$ab + ad < ab + bc \implies a(b + d) < b(a + c) \implies \frac{a}{b} < \frac{a + c}{b + d},$$

$$da + dc < cb + cd \implies d(a + c) < c(b + d) \implies \frac{a + c}{b + d} < \frac{c}{d},$$

whence the rational number $\frac{a + c}{b + d}$ lies between $\frac{a}{b}$ and $\frac{c}{d}$.

 We can also argue that the average of two distinct numbers lies between the numbers and so if $r_1 < r_2$ are rational numbers, then $\frac{r_1 + r_2}{2}$ lies between them.

5 Definition Let A be a set. If a belongs to the set A , then we write $a \in A$, read “ a is an element of A .” If a does not belong to the set A , we write $a \notin A$, read “ a is not an element of A .”

6 Definition (Conjunction, Disjunction, and Negation) The symbol \vee is read “or” (*disjunction*), the symbol \wedge is read “and” (*conjunction*), and the symbol \neg is read “not.”

7 Definition (Quantifiers) The symbol \forall is read “for all” (the *universal quantifier*), and the symbol \exists is read “there exists” (the *existential quantifier*).

We have

$$\neg(\forall x \in A, P(x)) \iff (\exists x \in A, \neg P(x)) \quad (1.1)$$

$$\neg(\exists x \in A, P(x)) \iff (\forall x \in A, \neg P(x)) \quad (1.2)$$

8 Definition (Subset) If $\forall a \in A$ we have $a \in B$, then we write $A \subseteq B$, which we read “ A is a subset of B .”

In particular, notice that for any set A , $\emptyset \subseteq A$ and $A \subseteq A$. Also

$$\mathbb{N} \subseteq \mathbb{Z} \subseteq \mathbb{Q} \subseteq \mathbb{R} \subseteq \mathbb{C}.$$

 $A = B \iff (A \subseteq B) \wedge (B \subseteq A).$

9 Definition The *union* of two sets A and B , is the set

$$A \cup B = \{x : (x \in A) \vee (x \in B)\}.$$

This is read “ A union B .” See figure 1.1.

10 Definition The *intersection* of two sets A and B , is

$$A \cap B = \{x : (x \in A) \wedge (x \in B)\}.$$

This is read “ A intersection B .” See figure 1.2.

11 Definition The *difference* of two sets A and B , is

$$A \setminus B = \{x : (x \in A) \wedge (x \notin B)\}.$$

This is read “ A set minus B .” See figure 1.3.




Figure 1.1: $A \cup B$




Figure 1.2: $A \cap B$




Figure 1.3: $A \setminus B$

12 Example Prove by means of set inclusion that

$$(A \cup B) \cap C = (A \cap C) \cup (B \cap C).$$

Solution: We have,

$$\begin{aligned}
 x \in (\mathbf{A} \cup \mathbf{B}) \cap \mathbf{C} &\iff x \in (\mathbf{A} \cup \mathbf{B}) \wedge x \in \mathbf{C} \\
 &\iff (x \in \mathbf{A} \vee x \in \mathbf{B}) \wedge x \in \mathbf{C} \\
 &\iff (x \in \mathbf{A} \wedge x \in \mathbf{C}) \vee (x \in \mathbf{B} \wedge x \in \mathbf{C}) \\
 &\iff (x \in \mathbf{A} \cap \mathbf{C}) \vee (x \in \mathbf{B} \cap \mathbf{C}) \\
 &\iff x \in (\mathbf{A} \cap \mathbf{C}) \cup (\mathbf{B} \cap \mathbf{C}),
 \end{aligned}$$

which establishes the equality.

13 Definition Let $\mathbf{A}_1, \mathbf{A}_2, \dots, \mathbf{A}_n$, be sets. The *Cartesian Product* of these n sets is defined and denoted by

$$\mathbf{A}_1 \times \mathbf{A}_2 \times \cdots \times \mathbf{A}_n = \{(a_1, a_2, \dots, a_n) : a_k \in \mathbf{A}_k\},$$

that is, the set of all ordered n -tuples whose elements belong to the given sets.

 In the particular case when all the \mathbf{A}_k are equal to a set \mathbf{A} , we write

$$\mathbf{A}_1 \times \mathbf{A}_2 \times \cdots \times \mathbf{A}_n = \mathbf{A}^n.$$

If $a \in \mathbf{A}$ and $b \in \mathbf{A}$ we write $(a, b) \in \mathbf{A}^2$.

14 Definition Let $x \in \mathbb{R}$. The *absolute value* of x —denoted by $|x|$ —is defined by

$$|x| = \begin{cases} -x & \text{if } x < 0, \\ x & \text{if } x \geq 0. \end{cases}$$

It follows from the definition that for $x \in \mathbb{R}$,

$$-|x| \leq x \leq |x|. \quad (1.3)$$

$$t \geq 0 \implies |x| \leq t \iff -t \leq x \leq t. \quad (1.4)$$

$$\forall a \in \mathbb{R} \implies \sqrt{a^2} = |a|. \quad (1.5)$$

15 Theorem (Triangle Inequality) Let $(a, b) \in \mathbb{R}^2$. Then

$$|a + b| \leq |a| + |b|. \quad (1.6)$$

Proof: From 1.3, by addition,

$$-|a| \leq a \leq |a|$$

to

$$-|b| \leq b \leq |b|$$

we obtain

$$-(|a| + |b|) \leq a + b \leq (|a| + |b|),$$

whence the theorem follows by 1.4. \square

16 Problem Prove that between any two rational numbers there is an irrational number.

17 Problem Prove that $X \setminus (X \setminus A) = X \cap A$.

18 Problem Prove that $X \setminus (A \cup B) = (X \setminus A) \cap (X \setminus B)$.

19 Problem Prove that $X \setminus (A \cap B) = (X \setminus A) \cup (X \setminus B)$.

20 Problem Prove that $(A \cup B) \setminus (A \cap B) = (A \setminus B) \cup (B \setminus A)$.

21 Problem Shew how to write the union $A \cup B \cup C$ as a disjoint union of sets.

22 Problem Prove that a set with $n \geq 0$ elements has 2^n subsets.

23 Problem Let $(a, b) \in \mathbb{R}^2$. Prove that $\|a\| - \|b\| \leq |a - b|$.

1.2 Partitions and Equivalence Relations

24 Definition Let $S \neq \emptyset$ be a set. A *partition* of S is a collection of non-empty, pairwise disjoint subsets of S whose union is S .

25 Example Let

$$2\mathbb{Z} = \{\dots, -6, -4, -2, 0, 2, 4, 6, \dots\} = \overline{0}$$

be the set of even integers and let

$$2\mathbb{Z} + 1 = \{\dots, -5, -3, -1, 1, 3, 5, \dots\} = \overline{1}$$

be the set of odd integers. Then

$$(2\mathbb{Z}) \cup (2\mathbb{Z} + 1) = \mathbb{Z}, \quad (2\mathbb{Z}) \cap (2\mathbb{Z} + 1) = \emptyset,$$

and so $\{2\mathbb{Z}, 2\mathbb{Z} + 1\}$ is a partition of \mathbb{Z} .

26 Example Let

$$3\mathbb{Z} = \{\dots, -9, -6, -3, 0, 3, 6, 9, \dots\} = \overline{0}$$

be the integral multiples of 3, let

$$3\mathbb{Z} + 1 = \{\dots, -8, -5, -2, 1, 4, 7, \dots\} = \overline{1}$$

be the integers leaving remainder 1 upon division by 3, and let

$$3\mathbb{Z} + 2 = \{\dots, -7, -4, -1, 2, 5, 8, \dots\} = \overline{2}$$

be integers leaving remainder 2 upon division by 3. Then

$$(3\mathbb{Z}) \cup (3\mathbb{Z} + 1) \cup (3\mathbb{Z} + 2) = \mathbb{Z},$$

$$(3\mathbb{Z}) \cap (3\mathbb{Z} + 1) = \emptyset, \quad (3\mathbb{Z}) \cap (3\mathbb{Z} + 2) = \emptyset, \quad (3\mathbb{Z} + 1) \cap (3\mathbb{Z} + 2) = \emptyset,$$

and so $\{3\mathbb{Z}, 3\mathbb{Z} + 1, 3\mathbb{Z} + 2\}$ is a partition of \mathbb{Z} .

 Notice that $\overline{0}$ and $\overline{1}$ do not mean the same in examples 25 and 26. Whenever we make use of this notation, the integral divisor must be made explicit.

27 Example Observe

$$\mathbb{R} = (\mathbb{Q}) \cup (\mathbb{R} \setminus \mathbb{Q}), \quad \emptyset = (\mathbb{Q}) \cap (\mathbb{R} \setminus \mathbb{Q}),$$

which means that the real numbers can be partitioned into the rational and irrational numbers.

28 Definition Let A, B be sets. A *relation* R is a subset of the Cartesian product $A \times B$. We write the fact that $(x, y) \in R$ as $x \sim y$.

29 Definition Let A be a set and R be a relation on $A \times A$. Then R is said to be

- reflexive if $(\forall x \in A), x \sim x$,
- symmetric if $(\forall (x, y) \in A^2), x \sim y \implies y \sim x$,
- anti-symmetric if $(\forall (x, y) \in A^2), (x \sim y) \wedge (y \sim x) \implies x = y$,
- transitive if $(\forall (x, y, z) \in A^3), (x \sim y) \wedge (y \sim z) \implies (x \sim z)$.

A relation R which is reflexive, symmetric and transitive is called an *equivalence relation* on A . A relation R which is reflexive, anti-symmetric and transitive is called a *partial order* on A .

30 Example Let $S = \{\text{All Human Beings}\}$, and define \sim on S as $a \sim b$ if and only if a and b have the same mother. Then $a \sim a$ since any human a has the same mother as himself. Similarly, $a \sim b \implies b \sim a$ and $(a \sim b) \wedge (b \sim c) \implies (a \sim c)$. Therefore \sim is an equivalence relation.

31 Example Let L be the set of all lines on the plane and write $l_1 \sim l_2$ if $l_1 \parallel l_2$ (the line l_1 is parallel to the line l_2). Then \sim is an equivalence relation on L .

32 Example In \mathbb{Q} define the relation $\frac{a}{b} \sim \frac{x}{y} \iff ay = bx$, where we will always assume that the denominators are non-zero. Then \sim is an equivalence relation. For $\frac{a}{b} \sim \frac{a}{b}$ since $ab = ab$. Clearly

$$\frac{a}{b} \sim \frac{x}{y} \implies ay = bx \implies xb = ya \implies \frac{x}{y} \sim \frac{a}{b}.$$

Finally, if $\frac{a}{b} \sim \frac{x}{y}$ and $\frac{x}{y} \sim \frac{s}{t}$ then we have $ay = bx$ and $xt = sy$. Multiplying these two equalities $ayxt = bxsy$. This gives

$$ayxt - bxsy = 0 \implies xy(at - bs) = 0.$$

Now if $x = 0$, we will have $a = s = 0$, in which case trivially $at = bs$. Otherwise we must have $at - bs = 0$ and so $\frac{a}{b} \sim \frac{s}{t}$.

33 Example Let X be a collection of sets. Write $A \sim B$ if $A \subseteq B$. Then \sim is a partial order on X .

34 Example For $(a, b) \in \mathbb{R}^2$ define

$$a \sim b \Leftrightarrow a^2 + b^2 > 2.$$

Determine, with proof, whether \sim is reflexive, symmetric, and/or transitive. Is \sim an equivalence relation?

Solution: Since $0^2 + 0^2 \not> 2$, we have $0 \not\sim 0$ and so \sim is not reflexive. Now,

$$\begin{aligned} a \sim b &\Leftrightarrow a^2 + b^2 \\ &\Leftrightarrow b^2 + a^2 \\ &\Leftrightarrow b \sim a, \end{aligned}$$

so \sim is symmetric. Also $0 \sim 3$ since $0^2 + 3^2 > 2$ and $3 \sim 1$ since $3^2 + 1^2 > 2$. But $0 \not\sim 1$ since $0^2 + 1^2 \not> 2$. Thus the relation is not transitive. The relation, therefore, is not an equivalence relation.

35 Definition Let \sim be an equivalence relation on a set S . Then the *equivalence class* of a is defined and denoted by

$$[a] = \{x \in S : x \sim a\}.$$

36 Lemma Let \sim be an equivalence relation on a set S . Then two equivalence classes are either identical or disjoint.

Proof: We prove that if $(a, b) \in S^2$, and $[a] \cap [b] \neq \emptyset$ then $[a] = [b]$. Suppose that $x \in [a] \cap [b]$. Now $x \in [a] \implies x \sim a \implies a \sim x$, by symmetry. Similarly, $x \in [b] \implies x \sim b$. By transitivity

$$(a \sim x) \wedge (x \sim b) \implies a \sim b.$$

Now, if $y \in [b]$ then $b \sim y$. Again by transitivity, $a \sim y$. This means that $y \in [a]$. We have shewn that $y \in [b] \implies y \in [a]$ and so $[b] \subseteq [a]$. In a similar fashion, we may prove that $[a] \subseteq [b]$. This establishes the result. \square

37 Theorem Let $S \neq \emptyset$ be a set. Any equivalence relation on S induces a partition of S . Conversely, given a partition of S into disjoint, non-empty subsets, we can define an equivalence relation on S whose equivalence classes are precisely these subsets.

Proof: By Lemma 36, if \sim is an equivalence relation on S then

$$S = \bigcup_{a \in S} [a],$$

and $[a] \cap [b] = \emptyset$ if $a \not\sim b$. This proves the first half of the theorem.

Conversely, let

$$S = \bigcup_{\alpha} S_{\alpha}, \quad S_{\alpha} \cap S_{\beta} = \emptyset \text{ if } \alpha \neq \beta,$$

be a partition of S . We define the relation \approx on S by letting $a \approx b$ if and only if they belong to the same S_{α} . Since the S_{α} are mutually disjoint, it is clear that \approx is an equivalence relation on S and that for $a \in S_{\alpha}$, we have $[a] = S_{\alpha}$. \square

38 Problem For $(a, b) \in (\mathbb{Q} \setminus \{0\})^2$ define the relation \sim as follows: $a \sim b \iff \frac{a}{b} \in \mathbb{Z}$. Determine whether this relation is reflexive, symmetric, and/or transitive.

39 Problem Give an example of a relation on $\mathbb{Z} \setminus \{0\}$ which is reflexive, but is neither symmetric nor transitive.

40 Problem Define the relation \sim in \mathbb{R} by $x \sim y \iff xe^y = ye^x$. Prove that \sim is an equivalence relation.

41 Problem Define the relation \sim in \mathbb{Q} by $x \sim y \iff \exists h \in \mathbb{Z}$ such that $x = \frac{3y + h}{3}$. [A] Prove that \sim is an equivalence relation. [B] Determine $[x]$, the equivalence of $x \in \mathbb{Q}$. [C] Is $\frac{2}{3} \sim \frac{4}{5}$?

1.3 Binary Operations

42 Definition Let S, T be sets. A *binary operation* is a function

$$\begin{aligned} \otimes : \quad & S \times S \rightarrow T \\ & (a, b) \mapsto (a, b) \end{aligned} .$$

We usually use the “infix” notation $a \otimes b$ rather than the “prefix” notation $\otimes(a, b)$. If $S = T$ then we say that the binary operation is *internal* or *closed* and if $S \neq T$ then we say that it is *external*. If

$$a \otimes b = b \otimes a$$

then we say that the operation \otimes is *commutative* and if

$$a \otimes (b \otimes c) = (a \otimes b) \otimes c,$$

we say that it is *associative*. If \otimes is associative, then we can write

$$a \otimes (b \otimes c) = (a \otimes b) \otimes c = a \otimes b \otimes c,$$

without ambiguity.

 We usually omit the sign \otimes and use juxtaposition to indicate the operation \otimes . Thus we write ab instead of $a \otimes b$.

43 Example The operation $+$ (ordinary addition) on the set $\mathbb{Z} \times \mathbb{Z}$ is a commutative and associative closed binary operation.

44 Example The operation $-$ (ordinary subtraction) on the set $\mathbb{N} \times \mathbb{N}$ is a non-commutative, non-associative non-closed binary operation.

45 Example The operation \otimes defined by $a \otimes b = 1 + ab$ on the set $\mathbb{Z} \times \mathbb{Z}$ is a commutative but non-associative internal binary operation. For

$$a \otimes b = 1 + ab = 1 + ba = ba,$$

proving commutativity. Also, $1 \otimes (2 \otimes 3) = 1 \otimes (7) = 8$ and $(1 \otimes 2) \otimes 3 = (3) \otimes 3 = 10$, evincing non-associativity.

46 Definition Let S be a set and $\otimes : S \times S \rightarrow S$ be a closed binary operation. The couple $\langle S, \otimes \rangle$ is called an *algebra*.

 When we desire to drop the sign \otimes and indicate the binary operation by juxtaposition, we simply speak of the “algebra S .”

47 Example Both $\langle \mathbb{Z}, + \rangle$ and $\langle \mathbb{Q}, \cdot \rangle$ are algebras. Here $+$ is the standard addition of real numbers and \cdot is the standard multiplication.

48 Example $\langle \mathbb{Z}, - \rangle$ is a non-commutative, non-associative algebra. Here $-$ is the standard subtraction operation on the real numbers

49 Example (Putnam Exam, 1972) Let S be a set and let $*$ be a binary operation of S satisfying the laws $\forall (x, y) \in S^2$

$$x * (x * y) = y, \tag{1.7}$$

$$(y * x) * x = y. \tag{1.8}$$

Shew that $*$ is commutative, but not necessarily associative.

Solution: By (1.8)

$$x * y = ((x * y) * x) * x.$$

By (1.8) again

$$((x * y) * x) * x = ((x * y) * ((x * y) * y)) * x.$$

By (1.7)

$$((x * y) * ((x * y) * y)) * x = (y) * x = y * x,$$

which is what we wanted to prove.

To shew that the operation is not necessarily associative, specialise $S = \mathbb{Z}$ and $x * y = -x - y$ (the opposite of x minus y). Then clearly in this case $*$ is commutative, and satisfies (1.7) and (1.8) but

$$0 * (0 * 1) = 0 * (-0 - 1) = 0 * (-1) = -0 - (-1) = 1,$$

and

$$(0 * 0) * 1 = (-0 - 0) * 1 = (0) * 1 = -0 - 1 = -1,$$

evincing that the operation is not associative.

50 Definition Let S be an algebra. Then $l \in S$ is called a *left identity* if $\forall s \in S$ we have $ls = s$. Similarly $r \in S$ is called a *right identity* if $\forall s \in S$ we have $sr = s$.

51 Theorem If an algebra S possesses a left identity l and a right identity r then $l = r$.

Proof: Since l is a left identity

$$r = lr.$$

Since r is a right identity

$$l = lr.$$

Combining these two, we gather

$$r = lr = l,$$

whence the theorem follows. \square

52 Example In $\langle \mathbb{Z}, + \rangle$ the element $0 \in \mathbb{Z}$ acts as an identity, and in $\langle \mathbb{Q}, \cdot \rangle$ the element $1 \in \mathbb{Q}$ acts as an identity.

53 Definition Let S be an algebra. An element $a \in S$ is said to be *left-cancellable* or *left-regular* if $\forall (x, y) \in S^2$

$$ax = ay \implies x = y.$$

Similarly, element $b \in S$ is said to be *right-cancellable* or *right-regular* if $\forall (x, y) \in S^2$

$$xb = yb \implies x = y.$$

Finally, we say an element $c \in S$ is *cancellable* or *regular* if it is both left and right cancellable.

54 Definition Let $\langle S, \otimes \rangle$ and $\langle S, \top \rangle$ be algebras. We say that \top is *left-distributive* with respect to \otimes if

$$\forall (x, y, z) \in S^3, x\top(y \otimes z) = (x\top y) \otimes (x\top z).$$

Similarly, we say that \top is *right-distributive* with respect to \otimes if

$$\forall (x, y, z) \in S^3, (y \otimes z)\top x = (y\top x) \otimes (z\top x).$$

We say that \top is *distributive* with respect to \otimes if it is both left and right distributive with respect to \otimes .

55 Problem Let

$$S = \{x \in \mathbb{Z} : \exists (a, b) \in \mathbb{Z}^2, x = a^3 + b^3 + c^3 - 3abc\}.$$

Prove that S is closed under multiplication, that is, if $x \in S$ and $y \in S$ then $xy \in S$.

56 Problem Let $\langle S, \otimes \rangle$ be an associative algebra, let $a \in S$ be a fixed element and define the closed binary operation \top by

$$x\top y = x \otimes a \otimes y.$$

Prove that \top is also associative over $S \times S$.

57 Problem On $\mathbb{Q} \cap]-1; 1[$ define the a binary operation \otimes

$$a \otimes b = \frac{a + b}{1 + ab},$$

where juxtaposition means ordinary multiplication and $+$ is the ordinary addition of real numbers. Prove that

① Prove that \otimes is a closed binary operation on $\mathbb{Q} \cap]-1; 1[$.

② Prove that \otimes is both commutative and associative.

③ Find an element $e \in \mathbb{R}$ such that $(\forall a \in \mathbb{Q} \cap]-1; 1[)$ ($e \otimes a = a$).

④ Given e as above and an arbitrary element $a \in \mathbb{Q} \cap]-1; 1[$, solve the equation $a \otimes b = e$ for b .

58 Problem On $\mathbb{R} \setminus \{1\}$ define the a binary operation \otimes

$$a \otimes b = a + b - ab,$$

where juxtaposition means ordinary multiplication and $+$ is the ordinary addition of real numbers. Clearly \otimes is a closed binary operation. Prove that

① Prove that \otimes is both commutative and associative.

② Find an element $e \in \mathbb{R} \setminus \{1\}$ such that $(\forall a \in \mathbb{R} \setminus \{1\})$ ($e \otimes a = a$).

③ Given e as above and an arbitrary element $a \in \mathbb{R} \setminus \{1\}$, solve the equation $a \otimes b = e$ for b .

59 Problem (Putnam Exam, 1971) Let S be a set and let \circ be a binary operation on S satisfying the two laws

$$(\forall x \in S)(x \circ x = x),$$

and

$$(\forall (x, y, z) \in S^3)((x \circ y) \circ z = (y \circ z) \circ x).$$

Show that \circ is commutative.

60 Problem Define the *symmetric difference* of the sets A, B as $A \Delta B = (A \setminus B) \cup (B \setminus A)$. Prove that Δ is commutative and associative.

1.4 \mathbb{Z}_n

61 Theorem (Division Algorithm) Let $n > 0$ be an integer. Then for any integer a there exist unique integers q (called the *quotient*) and r (called the *remainder*) such that $a = qn + r$ and $0 \leq r < n$.

Proof: In the proof of this theorem, we use the following property of the integers, called the well-ordering principle: any non-empty set of non-negative integers has a smallest element.

Consider the set

$$S = \{a - bn : b \in \mathbb{Z} \wedge a \geq bn\}.$$

Then S is a collection of nonnegative integers and $S \neq \emptyset$ as $\pm a - 0 \cdot n \in S$ and this is non-negative for one choice of sign. By the Well-Ordering Principle, S has a least element, say r . Now, there must be some $q \in \mathbb{Z}$ such that $r = a - qn$ since $r \in S$. By construction, $r \geq 0$. Let us prove that $r < n$. For assume that $r \geq n$. Then $r > r - n = a - qn - n = a - (q + 1)n \geq 0$, since $r - n \geq 0$. But then $a - (q + 1)n \in S$ and $a - (q + 1)n < r$ which contradicts the fact that r is the smallest member of S . Thus we must have $0 \leq r < n$. To prove that r and q are unique, assume that $q_1 n + r_1 = a = q_2 n + r_2$, $0 \leq r_1 < n$, $0 \leq r_2 < n$. Then $r_2 - r_1 = n(q_1 - q_2)$, that is, n divides $(r_2 - r_1)$. But $|r_2 - r_1| < n$, whence $r_2 = r_1$. From this it also follows that $q_1 = q_2$. This completes the proof. \square

62 Example If $n = 5$ the Division Algorithm says that we can arrange all the integers in five columns as follows:

⋮	⋮	⋮	⋮	⋮
-10	-9	-8	-7	-6
-5	-4	-3	-2	-1
0	1	2	3	4
5	6	7	8	9
⋮	⋮	⋮	⋮	⋮

The arrangement above shews that any integer comes in one of 5 flavours: those leaving remainder 0 upon division by 5, those leaving remainder 1 upon division by 5, etc. We let

$$5\mathbb{Z} = \{\dots, -15, -10, -5, 0, 5, 10, 15, \dots\} = \overline{0},$$

$$5\mathbb{Z} + 1 = \{\dots, -14, -9, -4, 1, 6, 11, 16, \dots\} = \overline{1},$$

$$5\mathbb{Z} + 2 = \{\dots, -13, -8, -3, 2, 7, 12, 17, \dots\} = \overline{2},$$

$$5\mathbb{Z} + 3 = \{\dots, -12, -7, -2, 3, 8, 13, 18, \dots\} = \overline{3},$$

$$5\mathbb{Z} + 4 = \{\dots, -11, -6, -1, 4, 9, 14, 19, \dots\} = \overline{4},$$

and

$$\mathbb{Z}_5 = \{\overline{0}, \overline{1}, \overline{2}, \overline{3}, \overline{4}\}.$$

Let n be a fixed positive integer. Define the relation \equiv by $x \equiv y$ if and only if they leave the same remainder upon division by n . Then clearly \equiv is an equivalence relation. As such it partitions the set of integers \mathbb{Z} into disjoint equivalence classes by Theorem 37. This motivates the following definition.

63 Definition Let n be a positive integer. The n residue classes upon division by n are

$$\overline{0} = n\mathbb{Z}, \quad \overline{1} = n\mathbb{Z} + 1, \quad \overline{2} = n\mathbb{Z} + 2, \quad \dots, \quad \overline{n-1} = n\mathbb{Z} + n - 1.$$

The set of residue classes modulo n is

$$\mathbb{Z}_n = \{\overline{0}, \overline{1}, \dots, \overline{n-1}\}.$$

Our interest is now to define some sort of “addition” and some sort of “multiplication” in \mathbb{Z}_n .

64 Theorem (Addition and Multiplication Modulo n) Let n be a positive integer. For $(\overline{a}, \overline{b}) \in (\mathbb{Z}_n)^2$ define $\overline{a} + \overline{b} = \overline{r}$, where r is the remainder of $a + b$ upon division by n . and $\overline{a} \cdot \overline{b} = \overline{t}$, where t is the remainder of ab upon division by n . Then these operations are well defined.

Proof: We need to prove that given arbitrary representatives of the residue classes, we always obtain the same result from our operations. That is, if $\overline{a} = \overline{a}'$ and $\overline{b} = \overline{b}'$ then we have $\overline{a} + \overline{b} = \overline{a}' + \overline{b}'$ and $\overline{a} \cdot \overline{b} = \overline{a}' \cdot \overline{b}'$.

Now

$$\overline{a} = \overline{a}' \implies \exists (q, q') \in \mathbb{Z}^2, r \in \mathbb{N}, a = qn + r, a' = q'n + r, 0 \leq r < n,$$

$$\overline{b} = \overline{b}' \implies \exists (q_1, q'_1) \in \mathbb{Z}^2, r_1 \in \mathbb{N}, b = q_1n + r_1, b' = q'_1n + r_1, 0 \leq r_1 < n.$$

Hence

$$a + b = (q + q_1)n + r + r_1, \quad a' + b' = (q' + q'_1)n + r + r_1,$$

meaning that both $a + b$ and $a' + b'$ leave the same remainder upon division by n , and therefore

$$\overline{a} + \overline{b} = \overline{a + b} = \overline{a' + b'} = \overline{a'} + \overline{b'}.$$

Similarly

$$ab = (qq_1n + qr_1 + rq_1)n + rr_1, \quad a'b' = (q'q'_1n + q'r_1 + rq'_1)n + rr_1,$$

and so both ab and $a'b'$ leave the same remainder upon division by n , and therefore

$$\overline{a} \cdot \overline{b} = \overline{ab} = \overline{a'b'} = \overline{a'} \cdot \overline{b'}.$$

This proves the theorem. \square

65 Example Let

$$\mathbb{Z}_6 = \{\overline{0}, \overline{1}, \overline{2}, \overline{3}, \overline{4}, \overline{5}\}$$

be the residue classes modulo 6. Construct the natural addition $+$ table for \mathbb{Z}_6 . Also, construct the natural multiplication \cdot table for \mathbb{Z}_6 .

Solution: The required tables are given in tables 1.1 and 1.2.

We notice that even though $\overline{2} \neq \overline{0}$ and $\overline{3} \neq \overline{0}$ we have $\overline{2} \cdot \overline{3} = \overline{0}$. This prompts the following definition.

66 Definition (Zero Divisor) An element $a \neq \overline{0}$ of \mathbb{Z}_n is called a zero divisor if $ab = \overline{0}$ for some $b \in \mathbb{Z}_n$.

$+$	$\bar{0}$	$\bar{1}$	$\bar{2}$	$\bar{3}$	$\bar{4}$	$\bar{5}$
$\bar{0}$	$\bar{0}$	$\bar{1}$	$\bar{2}$	$\bar{3}$	$\bar{4}$	$\bar{5}$
$\bar{1}$	$\bar{1}$	$\bar{2}$	$\bar{3}$	$\bar{4}$	$\bar{5}$	$\bar{0}$
$\bar{2}$	$\bar{2}$	$\bar{3}$	$\bar{4}$	$\bar{5}$	$\bar{0}$	$\bar{1}$
$\bar{3}$	$\bar{3}$	$\bar{4}$	$\bar{5}$	$\bar{0}$	$\bar{1}$	$\bar{2}$
$\bar{4}$	$\bar{4}$	$\bar{5}$	$\bar{0}$	$\bar{1}$	$\bar{2}$	$\bar{3}$
$\bar{5}$	$\bar{5}$	$\bar{0}$	$\bar{1}$	$\bar{2}$	$\bar{3}$	$\bar{4}$

 Table 1.1: Addition table for \mathbb{Z}_6 .

\cdot	$\bar{0}$	$\bar{1}$	$\bar{2}$	$\bar{3}$	$\bar{4}$	$\bar{5}$
$\bar{0}$						
$\bar{1}$	$\bar{0}$	$\bar{1}$	$\bar{2}$	$\bar{3}$	$\bar{4}$	$\bar{5}$
$\bar{2}$	$\bar{0}$	$\bar{2}$	$\bar{4}$	$\bar{0}$	$\bar{2}$	$\bar{4}$
$\bar{3}$	$\bar{0}$	$\bar{3}$	$\bar{0}$	$\bar{3}$	$\bar{0}$	$\bar{3}$
$\bar{4}$	$\bar{0}$	$\bar{4}$	$\bar{2}$	$\bar{0}$	$\bar{4}$	$\bar{2}$
$\bar{5}$	$\bar{0}$	$\bar{5}$	$\bar{4}$	$\bar{3}$	$\bar{2}$	$\bar{1}$

 Table 1.2: Multiplication table for \mathbb{Z}_6 .

We will extend the concept of zero divisor later on to various algebras.

67 Example

Let

$$\mathbb{Z}_7 = \{\bar{0}, \bar{1}, \bar{2}, \bar{3}, \bar{4}, \bar{5}, \bar{6}\}$$

be the residue classes modulo 7. Construct the natural addition $+$ table for \mathbb{Z}_7 . Also, construct the natural multiplication \cdot table for \mathbb{Z}_7 .

Solution: The required tables are given in tables 1.3 and 1.4.

$+$	$\bar{0}$	$\bar{1}$	$\bar{2}$	$\bar{3}$	$\bar{4}$	$\bar{5}$	$\bar{6}$
$\bar{0}$	$\bar{0}$	$\bar{1}$	$\bar{2}$	$\bar{3}$	$\bar{4}$	$\bar{5}$	$\bar{6}$
$\bar{1}$	$\bar{1}$	$\bar{2}$	$\bar{3}$	$\bar{4}$	$\bar{5}$	$\bar{6}$	$\bar{0}$
$\bar{2}$	$\bar{2}$	$\bar{3}$	$\bar{4}$	$\bar{5}$	$\bar{6}$	$\bar{0}$	$\bar{1}$
$\bar{3}$	$\bar{3}$	$\bar{4}$	$\bar{5}$	$\bar{6}$	$\bar{0}$	$\bar{1}$	$\bar{2}$
$\bar{4}$	$\bar{4}$	$\bar{5}$	$\bar{6}$	$\bar{0}$	$\bar{1}$	$\bar{2}$	$\bar{3}$
$\bar{5}$	$\bar{5}$	$\bar{6}$	$\bar{0}$	$\bar{1}$	$\bar{2}$	$\bar{3}$	$\bar{4}$
$\bar{6}$	$\bar{6}$	$\bar{0}$	$\bar{1}$	$\bar{2}$	$\bar{3}$	$\bar{4}$	$\bar{5}$

 Table 1.3: Addition table for \mathbb{Z}_7 .

\cdot	$\bar{0}$	$\bar{1}$	$\bar{2}$	$\bar{3}$	$\bar{4}$	$\bar{5}$	$\bar{6}$
$\bar{0}$							
$\bar{1}$	$\bar{0}$	$\bar{1}$	$\bar{2}$	$\bar{3}$	$\bar{4}$	$\bar{5}$	$\bar{6}$
$\bar{2}$	$\bar{0}$	$\bar{2}$	$\bar{4}$	$\bar{6}$	$\bar{1}$	$\bar{3}$	$\bar{5}$
$\bar{3}$	$\bar{0}$	$\bar{3}$	$\bar{6}$	$\bar{2}$	$\bar{5}$	$\bar{1}$	$\bar{4}$
$\bar{4}$	$\bar{0}$	$\bar{4}$	$\bar{1}$	$\bar{5}$	$\bar{2}$	$\bar{6}$	$\bar{3}$
$\bar{5}$	$\bar{0}$	$\bar{5}$	$\bar{3}$	$\bar{1}$	$\bar{6}$	$\bar{4}$	$\bar{2}$
$\bar{6}$	$\bar{0}$	$\bar{6}$	$\bar{5}$	$\bar{4}$	$\bar{3}$	$\bar{2}$	$\bar{1}$

 Table 1.4: Multiplication table for \mathbb{Z}_7 .

68 Example

Solve the equation

$$\bar{5}x = \bar{3}$$

in \mathbb{Z}_{11} .

Solution: Multiplying by $\bar{9}$ on both sides

$$\overline{45x} = \overline{27},$$

that is,

$$x = \bar{5}.$$

We will use the following result in the next section.

69 Definition Let a, b be integers with one of them different from 0. The greatest common divisor d of a, b , denoted by $d = \gcd(a, b)$ is the largest positive integer that divides both a and b .

70 Theorem (Bachet-Bezout Theorem) The greatest common divisor of any two integers a, b can be written as a linear combination of a and b , i.e., there are integers x, y with

$$\gcd(a, b) = ax + by.$$

Proof: Let $A = \{ax + by : ax + by > 0, x, y \in \mathbb{Z}\}$. Clearly one of $\pm a, \pm b$ is in A , as one of a, b is not zero. By the Well Ordering Principle, A has a smallest element, say d . Therefore, there are x_0, y_0 such that $d = ax_0 + by_0$. We prove that $d = \gcd(a, b)$. To do this we prove that d divides a and b and that if t divides a and b , then t must also divide then d .

We first prove that d divides a . By the Division Algorithm, we can find integers $q, r, 0 \leq r < d$ such that $a = dq + r$. Then

$$r = a - dq = a(1 - qx_0) - by_0.$$

If $r > 0$, then $r \in A$ is smaller than the smaller element of A , namely d , a contradiction. Thus $r = 0$. This entails $dq = a$, i.e. d divides a . We can similarly prove that d divides b .

Assume that t divides a and b . Then $a = tm, b = tn$ for integers m, n . Hence $d = ax_0 + bx_0 = t(mx_0 + ny_0)$, that is, t divides d . The theorem is thus proved. \square

71 Problem Write the addition and multiplication tables of \mathbb{Z}_{11} under natural addition and multiplication modulo 11.

72 Problem Solve the equation

$$\overline{5x^2} = \overline{3}$$

73 Problem Prove that if $n > 0$ is a composite integer, \mathbb{Z}_n has zero divisors.

1.5 Fields

74 Definition Let \mathbb{F} be a set having at least two elements $0_{\mathbb{F}}$ and $1_{\mathbb{F}}$ ($0_{\mathbb{F}} \neq 1_{\mathbb{F}}$) together with two operations \cdot (multiplication, which we usually represent via juxtaposition) and $+$ (addition). A field $\langle \mathbb{F}, \cdot, + \rangle$ is a triplet satisfying the following axioms $\forall (a, b, c) \in \mathbb{F}^3$:

F1 Addition and multiplication are associative:

$$(a + b) + c = a + (b + c), \quad (ab)c = a(bc) \quad (1.9)$$

F2 Addition and multiplication are commutative:

$$a + b = b + a, \quad ab = ba \quad (1.10)$$

F3 The multiplicative operation distributes over addition:

$$a(b + c) = ab + ac \quad (1.11)$$

F4 $0_{\mathbb{F}}$ is the additive identity:

$$0_{\mathbb{F}} + \mathbf{a} = \mathbf{a} + 0_{\mathbb{F}} = \mathbf{a} \quad (1.12)$$

F5 $1_{\mathbb{F}}$ is the multiplicative identity:

$$1_{\mathbb{F}}\mathbf{a} = \mathbf{a}1_{\mathbb{F}} = \mathbf{a} \quad (1.13)$$

F6 Every element has an additive inverse:

$$\exists -\mathbf{a} \in \mathbb{F}, \quad \mathbf{a} + (-\mathbf{a}) = (-\mathbf{a}) + \mathbf{a} = 0_{\mathbb{F}} \quad (1.14)$$

F7 Every non-zero element has a multiplicative inverse: if $\mathbf{a} \neq 0_{\mathbb{F}}$

$$\exists \mathbf{a}^{-1} \in \mathbb{F}, \quad \mathbf{a}\mathbf{a}^{-1} = \mathbf{a}^{-1}\mathbf{a} = 1_{\mathbb{F}} \quad (1.15)$$

The elements of a field are called *scalars*.

An important property of fields is the following.

75 Theorem A field does not have zero divisors.

Proof: Assume that $\mathbf{a}\mathbf{b} = 0_{\mathbb{F}}$. If $\mathbf{a} \neq 0_{\mathbb{F}}$ then it has a multiplicative inverse \mathbf{a}^{-1} . We deduce

$$\mathbf{a}^{-1}\mathbf{a}\mathbf{b} = \mathbf{a}^{-1}0_{\mathbb{F}} \implies \mathbf{b} = 0_{\mathbb{F}}.$$

This means that the only way of obtaining a zero product is if one of the factors is $0_{\mathbb{F}}$. \square

76 Example $\langle \mathbb{Q}, \cdot, + \rangle$, $\langle \mathbb{R}, \cdot, + \rangle$, and $\langle \mathbb{C}, \cdot, + \rangle$ are all fields. The multiplicative identity in each case is 1 and the additive identity is 0.

77 Example Let

$$\mathbb{Q}(\sqrt{2}) = \{\mathbf{a} + \sqrt{2}\mathbf{b} : (\mathbf{a}, \mathbf{b}) \in \mathbb{Q}^2\}$$

and define addition on this set as

$$(\mathbf{a} + \sqrt{2}\mathbf{b}) + (\mathbf{c} + \sqrt{2}\mathbf{d}) = (\mathbf{a} + \mathbf{c}) + \sqrt{2}(\mathbf{b} + \mathbf{d}),$$

and multiplication as

$$(\mathbf{a} + \sqrt{2}\mathbf{b})(\mathbf{c} + \sqrt{2}\mathbf{d}) = (\mathbf{a}\mathbf{c} + 2\mathbf{b}\mathbf{d}) + \sqrt{2}(\mathbf{ad} + \mathbf{bc}).$$

Then $\langle \mathbb{Q} + \sqrt{2}\mathbb{Q}, \cdot, + \rangle$ is a field. Observe $0_{\mathbb{F}} = 0$, $1_{\mathbb{F}} = 1$, that the additive inverse of $\mathbf{a} + \sqrt{2}\mathbf{b}$ is $-\mathbf{a} - \sqrt{2}\mathbf{b}$, and the multiplicative inverse of $\mathbf{a} + \sqrt{2}\mathbf{b}$, $(\mathbf{a}, \mathbf{b}) \neq (0, 0)$ is

$$(\mathbf{a} + \sqrt{2}\mathbf{b})^{-1} = \frac{1}{\mathbf{a} + \sqrt{2}\mathbf{b}} = \frac{\mathbf{a} - \sqrt{2}\mathbf{b}}{\mathbf{a}^2 - 2\mathbf{b}^2} = \frac{\mathbf{a}}{\mathbf{a}^2 - 2\mathbf{b}^2} - \frac{\sqrt{2}\mathbf{b}}{\mathbf{a}^2 - 2\mathbf{b}^2}.$$

Here $\mathbf{a}^2 - 2\mathbf{b}^2 \neq 0$ since $\sqrt{2}$ is irrational.

78 Theorem If p is a prime, $\langle \mathbb{Z}_p, \cdot, + \rangle$ is a field under \cdot multiplication modulo p and $+$ addition modulo p .

Proof: Clearly the additive identity is $\bar{0}$ and the multiplicative identity is $\bar{1}$. The additive inverse of \bar{a} is $\bar{p-a}$. We must prove that every $\bar{a} \in \mathbb{Z}_p \setminus \{\bar{0}\}$ has a multiplicative inverse. Such an \mathbf{a} satisfies $\gcd(a, p) = 1$ and by the Bachet-Bezout Theorem 70, there exist integers x, y with $px + ay = 1$. In such case we have

$$\bar{1} = \overline{px + ay} = \overline{ay} = \bar{a} \cdot \bar{y},$$

whence $(\bar{a})^{-1} = \bar{y}$. \square

79 Definition A field is said to be of *characteristic $p \neq 0$* if for some positive integer p we have $\forall a \in F, pa = 0_F$, and no positive integer smaller than p enjoys this property.

If the field does not have characteristic $p \neq 0$ then we say that it is of *characteristic 0*. Clearly \mathbb{Q}, \mathbb{R} and \mathbb{C} are of characteristic 0, while \mathbb{Z}_p for prime p , is of characteristic p .

80 Theorem The characteristic of a field is either 0 or a prime.

Proof: If the characteristic of the field is 0, there is nothing to prove. Let p be the least positive integer for which $\forall a \in F, pa = 0_F$. Let us prove that p must be a prime. Assume that instead we had $p = st$ with integers $s > 1, t > 1$. Take $a = 1_F$. Then we must have $(st)1_F = 0_F$, which entails $(s1_F)(t1_F) = 0_F$. But in a field there are no zero-divisors by Theorem 75, hence either $s1_F = 0_F$ or $t1_F = 0_F$. But either of these equalities contradicts the minimality of p . Hence p is a prime. \square

81 Problem Consider the set of numbers

$$\mathbb{Q}(\sqrt{2}, \sqrt{3}, \sqrt{6}) = \{a + b\sqrt{2} + c\sqrt{3} + d\sqrt{6} : (a, b, c, d) \in \mathbb{Q}^4\}$$

Assume that $\mathbb{Q}(\sqrt{2}, \sqrt{3}, \sqrt{6})$ is a field under ordinary addition and multiplication. What is the multiplicative inverse of the element $\sqrt{2} + 2\sqrt{3} + 3\sqrt{6}$?

82 Problem Let F be a field and a, b two non-zero elements of F . Prove that

$$-(ab^{-1}) = (-a)b^{-1} = a(-b^{-1}).$$

83 Problem Let F be a field and $a \neq 0_F$. Prove that

$$(-a)^{-1} = -(a^{-1}).$$

84 Problem Let F be a field and a, b two non-zero elements of F . Prove that

$$ab^{-1} = (-a)(-b^{-1}).$$

1.6 Functions

85 Definition By a *function* or a *mapping* from one set to another, we mean a rule or mechanism that assigns to every input element of the first set a unique output element of the second set. We shall call the set of inputs the *domain* of the function, the set of possible outputs the *target set* of the function, and the set of actual outputs the *image* of the function.

We will generally refer to a function with the following notation:

$$\begin{array}{ccc} D & \rightarrow & T \\ f: & & . \\ x & \mapsto & f(x) \end{array}$$

Here f is the *name of the function*, D is its domain, T is its target set, x is the name of a typical input and $f(x)$ is the output or *image of x under f* . We call the assignment $x \mapsto f(x)$ the *assignment rule* of the function. Sometimes x is also called the *independent variable*. The set $f(D) = \{f(a) | a \in D\}$ is called the *image* of f . Observe that $f(D) \subseteq T$.




Figure 1.4: An injection.




Figure 1.5: Not an injection

86 Definition A function $f : X \rightarrow Y$ is said to be *injective* or *one-to-one* if $\forall (a, b) \in X^2$, we have
 $x \mapsto f(x)$

$$a \neq b \implies f(a) \neq f(b).$$

This is equivalent to saying that

$$f(a) = f(b) \implies a = b.$$

87 Example The function α in the diagram 1.4 is an injective function. The function β represented by the diagram 1.5, however, is not injective, $\beta(3) = \beta(1) = 4$, but $3 \neq 1$.

88 Example Prove that

$$\begin{aligned} t : \mathbb{R} \setminus \{1\} &\rightarrow \mathbb{R} \setminus \{1\} \\ x &\mapsto \frac{x+1}{x-1} \end{aligned}$$

is an injection.

Solution: Assume $t(a) = t(b)$. Then

$$\begin{aligned} t(a) = t(b) &\implies \frac{a+1}{a-1} = \frac{b+1}{b-1} \\ &\implies (a+1)(b-1) = (b+1)(a-1) \\ &\implies ab - a + b - 1 = ab - b + a - 1 \\ &\implies 2a = 2b \\ &\implies a = b \end{aligned}$$

We have proved that $t(a) = t(b) \implies a = b$, which shews that t is injective.




Figure 1.6: A surjection




Figure 1.7: Not a surjection

89 Definition A function $f : A \rightarrow B$ is said to be *surjective* or *onto* if $(\forall b \in B) (\exists a \in A) : f(a) = b$. That is, each element of B has a pre-image in A .

A function is surjective if its image coincides with its target set. It is easy to see that a graphical criterion for a function to be surjective is that every horizontal line passing through a point of the target set (a subset of the y-axis) of the function must also meet the curve.

90 Example The function β represented by diagram 1.6 is surjective. The function γ represented by diagram 1.7 is not surjective as 8 does not have a preimage.

91 Example Prove that $t : \mathbb{R} \rightarrow \mathbb{R}$ is a surjection.

$$\begin{array}{rcl} \mathbb{R} & \rightarrow & \mathbb{R} \\ x & \mapsto & x^3 \end{array}$$

Solution: Since the graph of t is that of a cubic polynomial with only one zero, every horizontal line passing through a point in \mathbb{R} will eventually meet the graph of t , whence t is surjective. To prove this analytically, proceed as follows. We must prove that $(\forall b \in \mathbb{R}) (\exists a)$ such that $t(a) = b$. We choose a so that $a = b^{1/3}$. Then

$$t(a) = t(b^{1/3}) = (b^{1/3})^3 = b.$$

Our choice of a works and hence the function is surjective.

92 Definition A function is *bijection* if it is both injective and surjective.

93 Problem Prove that

$$\begin{array}{rcl} \mathbb{R} & \rightarrow & \mathbb{R} \\ h : & & \\ x & \mapsto & x^3 \end{array}$$

is an injection.

94 Problem Show that

$$\begin{array}{rcl} \mathbb{R} \setminus \left\{\frac{3}{2}\right\} & \rightarrow & \mathbb{R} \setminus \{3\} \\ f : & & \\ x & \mapsto & \frac{6x}{2x-3} \end{array}$$

is a bijection.

Chapter 2

Matrices and Matrix Operations

2.1 The Algebra of Matrices

95 Definition Let $\langle \mathbb{F}, \cdot, + \rangle$ be a field. An $m \times n$ (m by n) *matrix A* with m rows and n columns with entries over \mathbb{F} is a rectangular array of the form

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix},$$

where $\forall (i, j) \in \{1, 2, \dots, m\} \times \{1, 2, \dots, n\}$, $a_{ij} \in \mathbb{F}$.

 As a shortcut, we often use the notation $A = [a_{ij}]$ to denote the matrix A with entries a_{ij} . Notice that when we refer to the matrix we put parentheses—as in “[a_{ij}]”, and when we refer to a specific entry we do not use the surrounding parentheses—as in “ a_{ij} .”

96 Example

$$A = \begin{bmatrix} 0 & -1 & 1 \\ 1 & 2 & 3 \end{bmatrix}$$

is a 2×3 matrix and

$$B = \begin{bmatrix} -2 & 1 \\ 1 & 2 \\ 0 & 3 \end{bmatrix}$$

is a 3×2 matrix.

97 Example Write out explicitly the 4×4 matrix $A = [a_{ij}]$ where $a_{ij} = i^2 - j^2$.

Solution: This is

$$\mathbf{A} = \begin{bmatrix} 1^2 - 1^1 & 1^2 - 2^2 & 1^2 - 3^2 & 1^2 - 4^2 \\ 2^2 - 1^2 & 2^2 - 2^2 & 2^2 - 3^2 & 2^2 - 4^2 \\ 3^2 - 1^2 & 3^2 - 2^2 & 3^2 - 3^2 & 3^2 - 4^2 \\ 4^2 - 1^2 & 4^2 - 2^2 & 4^2 - 3^2 & 4^2 - 4^2 \end{bmatrix} = \begin{bmatrix} 0 & -3 & -8 & -15 \\ 3 & 0 & -5 & -12 \\ 8 & 5 & 0 & -7 \\ 15 & 12 & 7 & 0 \end{bmatrix}.$$

98 Definition Let $\langle \mathbb{F}, \cdot, + \rangle$ be a field. We denote by $M_{m \times n}(\mathbb{F})$ the set of all $m \times n$ matrices with entries over \mathbb{F} . If $m = n$ we use the abbreviated notation $M_n(\mathbb{F}) = M_{n \times n}(\mathbb{F})$. $M_n(\mathbb{F})$ is thus the set of all square matrices of size n with entries over \mathbb{F} .

99 Definition The $m \times n$ zero matrix $0_{m \times n} \in M_{m \times n}(\mathbb{F})$ is the matrix with $0_{\mathbb{F}}$'s everywhere,

$$0_{m \times n} = \begin{bmatrix} 0_{\mathbb{F}} & 0_{\mathbb{F}} & 0_{\mathbb{F}} & \cdots & 0_{\mathbb{F}} \\ 0_{\mathbb{F}} & 0_{\mathbb{F}} & 0_{\mathbb{F}} & \cdots & 0_{\mathbb{F}} \\ 0_{\mathbb{F}} & 0_{\mathbb{F}} & 0_{\mathbb{F}} & \cdots & 0_{\mathbb{F}} \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ 0_{\mathbb{F}} & 0_{\mathbb{F}} & 0_{\mathbb{F}} & \cdots & 0_{\mathbb{F}} \end{bmatrix}.$$

When $m = n$ we write 0_n as a shortcut for $0_{n \times n}$.

100 Definition The $n \times n$ identity matrix $I_n \in M_n(\mathbb{F})$ is the matrix with $1_{\mathbb{F}}$'s on the main diagonal and $0_{\mathbb{F}}$'s everywhere else,

$$I_n = \begin{bmatrix} 1_{\mathbb{F}} & 0_{\mathbb{F}} & 0_{\mathbb{F}} & \cdots & 0_{\mathbb{F}} \\ 0_{\mathbb{F}} & 1_{\mathbb{F}} & 0_{\mathbb{F}} & \cdots & 0_{\mathbb{F}} \\ 0_{\mathbb{F}} & 0_{\mathbb{F}} & 1_{\mathbb{F}} & \cdots & 0_{\mathbb{F}} \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ 0_{\mathbb{F}} & 0_{\mathbb{F}} & 0_{\mathbb{F}} & \cdots & 1_{\mathbb{F}} \end{bmatrix}.$$

101 Definition (Matrix Addition and Multiplication of a Matrix by a Scalar) Let $\mathbf{A} = [a_{ij}] \in M_{m \times n}(\mathbb{F})$, $\mathbf{B} = [b_{ij}] \in M_{m \times n}(\mathbb{F})$ and $\alpha \in \mathbb{F}$. The matrix $\mathbf{A} + \alpha\mathbf{B}$ is the matrix $\mathbf{C} \in M_{m \times n}(\mathbb{F})$ with entries $\mathbf{C} = [c_{ij}]$ where $c_{ij} = a_{ij} + \alpha b_{ij}$.

102 Example For $\mathbf{A} = \begin{bmatrix} 1 & 1 \\ -1 & 1 \\ 0 & 2 \end{bmatrix}$ and $\mathbf{B} = \begin{bmatrix} -1 & 1 \\ 2 & 1 \\ 0 & -1 \end{bmatrix}$ we have

$$\mathbf{A} + 2\mathbf{B} = \begin{bmatrix} -1 & 3 \\ 3 & 3 \\ 0 & 0 \end{bmatrix}.$$

103 Theorem Let $(\mathbf{A}, \mathbf{B}, \mathbf{C}) \in (\mathbf{M}_{m \times n}(\mathbb{F}))^3$ and $(\alpha, \beta) \in \mathbb{F}^2$. Then

M1 $\mathbf{M}_{m \times n}(\mathbb{F})$ is close under matrix addition and scalar multiplication

$$\mathbf{A} + \mathbf{B} \in \mathbf{M}_{m \times n}(\mathbb{F}), \quad \alpha\mathbf{A} \in \mathbf{M}_{m \times n}(\mathbb{F}) \quad (2.1)$$

M2 Addition of matrices is commutative

$$\mathbf{A} + \mathbf{B} = \mathbf{B} + \mathbf{A} \quad (2.2)$$

M3 Addition of matrices is associative

$$\mathbf{A} + (\mathbf{B} + \mathbf{C}) = (\mathbf{A} + \mathbf{B}) + \mathbf{C} \quad (2.3)$$

M4 There is a matrix $\mathbf{0}_{m \times n}$ such that

$$\mathbf{A} + \mathbf{0}_{m \times n} \quad (2.4)$$

M5 There is a matrix $-\mathbf{A}$ such that

$$\mathbf{A} + (-\mathbf{A}) = (-\mathbf{A}) + \mathbf{A} = \mathbf{0}_{m \times n} \quad (2.5)$$

M6 Distributive law

$$\alpha(\mathbf{A} + \mathbf{B}) = \alpha\mathbf{A} + \alpha\mathbf{B} \quad (2.6)$$

M7 Distributive law

$$(\alpha + \beta)\mathbf{A} = \alpha\mathbf{A} + \beta\mathbf{B} \quad (2.7)$$

M8

$$1_{\mathbb{F}}\mathbf{A} = \mathbf{A} \quad (2.8)$$

M9

$$\alpha(\beta\mathbf{A}) = (\alpha\beta)\mathbf{A} \quad (2.9)$$

Proof: The theorem follows at once by reducing each statement to an entry-wise and appealing to the field axioms. \square

104 Problem Write out explicitly the 3×3 matrix $\mathbf{A} = [a_{ij}]$ where $a_{ij} = i^j$.

105 Problem Write out explicitly the 3×3 matrix $\mathbf{A} = [a_{ij}]$ where $a_{ij} = ij$.

106 Problem Let

$$\mathbf{M} = \begin{bmatrix} a & -2a & c \\ 0 & -a & b \\ a+b & 0 & -1 \end{bmatrix}, \quad \mathbf{N} = \begin{bmatrix} 1 & 2a & c \\ a & b-a & -b \\ a-b & 0 & -1 \end{bmatrix}$$

be square matrices with entries over \mathbb{R} . Find $\mathbf{M} + \mathbf{N}$ and $2\mathbf{M}$.

107 Problem Determine x and y such that

$$\begin{bmatrix} 3 & x & 1 \\ 1 & 2 & 0 \end{bmatrix} + 2 \begin{bmatrix} 2 & 1 & 3 \\ 5 & x & 4 \end{bmatrix} = \begin{bmatrix} 7 & 3 & 7 \\ 11 & y & 8 \end{bmatrix}.$$

108 Problem Determine 2×2 matrices \mathbf{A} and \mathbf{B} such that

$$2\mathbf{A} - 5\mathbf{B} = \begin{bmatrix} 1 & -2 \\ 0 & 1 \end{bmatrix}, \quad -2\mathbf{A} + 6\mathbf{B} = \begin{bmatrix} 4 & 2 \\ 6 & 0 \end{bmatrix}.$$

109 Problem Let $\mathbf{A} = [a_{ij}] \in M_n(\mathbb{R})$. Prove that

$$\min_j \max_i a_{ij} \geq \max_i \min_j a_{ij}.$$

110 Problem A person goes along the rows of a movie theater and asks the tallest person of each row to stand up. Then he selects the shortest of these people, who we will call the

shortest giant. Another person goes along the rows and asks the shortest person to stand up and from these he selects the tallest, which we will call the tallest midget. Who is taller, the tallest midget or the shortest giant?

111 Problem (Putnam Exam, 1959) Choose five elements from the matrix

$$\begin{bmatrix} 11 & 17 & 25 & 19 & 16 \\ 24 & 10 & 13 & 15 & 3 \\ 12 & 5 & 14 & 2 & 18 \\ 23 & 4 & 1 & 8 & 22 \\ 6 & 20 & 7 & 21 & 9 \end{bmatrix},$$

no two coming from the same row or column, so that the minimum of these five elements is as large as possible.

2.2 Matrix Multiplication

112 Definition Let $\mathbf{A} = [a_{ij}] \in M_{m \times n}(\mathbb{F})$ and $\mathbf{B} = [b_{ij}] \in M_{n \times p}(\mathbb{F})$. Then the matrix product \mathbf{AB} is defined as the matrix $\mathbf{C} = [c_{ij}] \in M_{m \times p}(\mathbb{F})$ with entries $c_{ij} = \sum_{l=1}^n a_{il} b_{lj}$:

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{i1} & a_{i2} & \cdots & a_{in} \\ \vdots & \vdots & \cdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \begin{bmatrix} b_{11} & \cdots & b_{1j} & \cdots & b_{1p} \\ b_{21} & \cdots & b_{2j} & \cdots & b_{2p} \\ \vdots & \cdots & \vdots & \cdots & \vdots \\ b_{n1} & \cdots & b_{nj} & \cdots & b_{np} \end{bmatrix} = \begin{bmatrix} c_{11} & \cdots & c_{1p} \\ c_{21} & \cdots & c_{2p} \\ \vdots & \cdots & \vdots \\ \cdots & c_{ij} & \cdots \\ \vdots & \cdots & \vdots \\ c_{m1} & \cdots & c_{mp} \end{bmatrix}.$$



Observe that we use juxtaposition rather than a special symbol to denote matrix multiplication. This will simplify notation. In order to obtain the ij -th entry of the matrix \mathbf{AB} we multiply elementwise the i -th row of \mathbf{A} by the j -th column of \mathbf{B} . Observe that \mathbf{AB} is a $m \times p$ matrix.

113 Example Let $\mathbf{M} = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ and $\mathbf{N} = \begin{bmatrix} 5 & 6 \\ 7 & 8 \end{bmatrix}$ be matrices over \mathbb{R} . Then

$$\mathbf{MN} = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \begin{bmatrix} 5 & 6 \\ 7 & 8 \end{bmatrix} = \begin{bmatrix} 1 \cdot 5 + 2 \cdot 7 & 1 \cdot 6 + 2 \cdot 8 \\ 3 \cdot 5 + 4 \cdot 7 & 3 \cdot 6 + 4 \cdot 8 \end{bmatrix} = \begin{bmatrix} 19 & 22 \\ 43 & 50 \end{bmatrix},$$

and

$$\mathbf{NM} = \begin{bmatrix} 5 & 6 \\ 7 & 8 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} 5 \cdot 1 + 6 \cdot 3 & 5 \cdot 2 + 6 \cdot 4 \\ 7 \cdot 1 + 8 \cdot 3 & 7 \cdot 2 + 8 \cdot 4 \end{bmatrix} = \begin{bmatrix} 23 & 34 \\ 31 & 46 \end{bmatrix}.$$

Hence, in particular, matrix multiplication is not necessarily commutative.

114 Example We have

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix},$$

over \mathbb{R} . Observe then that the product of two non-zero matrices may be the zero matrix.

115 Example Consider the matrix

$$\mathbf{A} = \begin{bmatrix} \bar{2} & \bar{1} & \bar{3} \\ \bar{0} & \bar{1} & \bar{1} \\ \bar{4} & \bar{4} & \bar{0} \end{bmatrix}$$

with entries over \mathbb{Z}_5 . Then

$$\begin{aligned} \mathbf{A}^2 &= \begin{bmatrix} \bar{2} & \bar{1} & \bar{3} \\ \bar{0} & \bar{1} & \bar{1} \\ \bar{4} & \bar{4} & \bar{0} \end{bmatrix} \begin{bmatrix} \bar{2} & \bar{1} & \bar{3} \\ \bar{0} & \bar{1} & \bar{1} \\ \bar{4} & \bar{4} & \bar{0} \end{bmatrix} \\ &= \begin{bmatrix} \bar{1} & \bar{0} & \bar{2} \\ \bar{4} & \bar{0} & \bar{1} \\ \bar{3} & \bar{3} & \bar{1} \end{bmatrix}. \end{aligned}$$



Even though matrix multiplication is not necessarily commutative, it is associative.

116 Theorem If $(\mathbf{A}, \mathbf{B}, \mathbf{C}) \in M_{m \times n}(\mathbb{F}) \times M_{n \times r}(\mathbb{F}) \times M_{r \times s}(\mathbb{F})$ we have

$$(\mathbf{AB})\mathbf{C} = \mathbf{A}(\mathbf{BC}),$$

i.e., matrix multiplication is associative.

Proof: To shew this we only need to consider the ij -th entry of each side, appeal to the associativity of the underlying field \mathbb{F} and verify that both sides are indeed equal to

$$\sum_{k=1}^n \sum_{k'=1}^r a_{ik} b_{kk'} c_{k'j}.$$



 By virtue of associativity, a square matrix commutes with its powers, that is, if $A \in M_n(\mathbb{F})$, and $(r, s) \in \mathbb{N}^2$, then $(A^r)(A^s) = (A^s)(A^r) = A^{r+s}$.

117 Example Let $A \in M_3(\mathbb{R})$ be given by

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}.$$

Demonstrate, using induction, that $A^n = 3^{n-1}A$ for $n \in \mathbb{N}, n \geq 1$.

Solution: The assertion is trivial for $n = 1$. Assume its truth for $n - 1$, that is, assume $A^{n-1} = 3^{n-2}A$. Observe that

$$A^2 = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 3 & 3 \\ 3 & 3 & 3 \\ 3 & 3 & 3 \end{bmatrix} = 3A.$$

Now

$$A^n = AA^{n-1} = A(3^{n-2}A) = 3^{n-2}A^2 = 3^{n-2}3A = 3^{n-1}A,$$

and so the assertion is proved by induction.

118 Theorem Let $A \in M_n(\mathbb{F})$. Then there is a unique identity matrix. That is, if $E \in M_n(\mathbb{F})$ is such that $AE = EA = A$, then $E = I_n$.

Proof: It is clear that for any $A \in M_n(\mathbb{F})$, $AI_n = I_nA = A$. Now because E is an identity, $EI_n = I_n$. Because I_n is an identity, $EI_n = E$. Whence

$$I_n = EI_n = E,$$

demonstrating uniqueness. \square

119 Example Let $A = [a_{ij}] \in M_n(\mathbb{R})$ be such that $a_{ij} = 0$ for $i > j$ and $a_{ij} = 1$ if $i \leq j$. Find A^2 .

Solution: Let $A^2 = B = [b_{ij}]$. Then

$$b_{ij} = \sum_{k=1}^n a_{ik}a_{kj}.$$

Observe that the i -th row of A has $i - 1$ 0's followed by $n - i + 1$ 1's, and the j -th column of A has j 1's followed by $n - j$ 0's. Therefore if $i - 1 > j$, then $b_{ij} = 0$. If $i \leq j + 1$, then

$$b_{ij} = \sum_{k=i}^j a_{ik}a_{kj} = j - i + 1.$$

This means that

$$\mathbf{A}^2 = \begin{bmatrix} 1 & 2 & 3 & 4 & \cdots & n-1 & n \\ 0 & 1 & 2 & 3 & \cdots & n-2 & n-1 \\ 0 & 0 & 1 & 2 & \cdots & n-3 & n-2 \\ \vdots & \vdots & \vdots & \vdots & \cdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & 1 & 2 \\ 0 & 0 & 0 & 0 & \cdots & 0 & 1 \end{bmatrix}.$$

120 Problem Determine the product

$$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} -2 & 1 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix}$$

121 Problem Let $\mathbf{A} = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 1 & 1 \end{bmatrix}$, $\mathbf{B} = \begin{bmatrix} a & b & c \\ c & a & b \\ b & c & a \end{bmatrix}$. Find \mathbf{AB} and \mathbf{BA} .

122 Problem Let

$$\mathbf{A} = \begin{bmatrix} \bar{2} & \bar{3} & \bar{4} & \bar{1} \\ \bar{1} & \bar{2} & \bar{3} & \bar{4} \\ \bar{4} & \bar{1} & \bar{2} & \bar{3} \\ \bar{3} & \bar{4} & \bar{1} & \bar{2} \end{bmatrix}, \quad \mathbf{B} = \begin{bmatrix} \bar{1} & \bar{1} & \bar{1} & \bar{1} \\ \bar{1} & \bar{1} & \bar{1} & \bar{1} \\ \bar{1} & \bar{1} & \bar{1} & \bar{1} \\ \bar{1} & \bar{1} & \bar{1} & \bar{1} \end{bmatrix}$$

be matrices in $M_4(\mathbb{Z}_5)$. Find the products \mathbf{AB} and \mathbf{BA} .

123 Problem Solve the equation

$$\begin{bmatrix} -4 & x \\ -x & 4 \end{bmatrix}^2 = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$$

over \mathbb{R} .

124 Problem Prove or disprove! If $(\mathbf{A}, \mathbf{B}) \in (M_n(\mathbb{F}))^2$ are such that $\mathbf{AB} = \mathbf{0}_n$, then also $\mathbf{BA} = \mathbf{0}_n$.

125 Problem Prove or disprove! For all matrices $(\mathbf{A}, \mathbf{B}) \in (M_n(\mathbb{F}))^2$,

$$(\mathbf{A} + \mathbf{B})(\mathbf{A} - \mathbf{B}) = \mathbf{A}^2 - \mathbf{B}^2.$$

126 Problem Prove, using mathematical induction, that

$$\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}^n = \begin{bmatrix} 1 & n \\ 0 & 1 \end{bmatrix}.$$

127 Problem Let $\mathbf{M} = \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$. Find \mathbf{M}^6 .

128 Problem Let $\mathbf{A} = \begin{bmatrix} 0 & 3 \\ 2 & 0 \end{bmatrix}$. Find, with proof, \mathbf{A}^{2003} .

129 Problem Let $(\mathbf{A}, \mathbf{B}, \mathbf{C}) \in M_{l \times m}(\mathbb{F}) \times M_{m \times n}(\mathbb{F}) \times M_{n \times l}(\mathbb{F})$ and $\alpha \in \mathbb{F}$. Prove that

$$\mathbf{A}(\mathbf{B} + \mathbf{C}) = \mathbf{AB} + \mathbf{AC},$$

$$(\mathbf{A} + \mathbf{B})\mathbf{C} = \mathbf{AC} + \mathbf{BC},$$

$$\alpha(\mathbf{AB}) = (\alpha\mathbf{A})\mathbf{B} = \mathbf{A}(\alpha\mathbf{B}).$$

130 Problem Let $\mathbf{A} \in M_2(\mathbb{R})$ be given by

$$\mathbf{A} = \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}.$$

Demonstrate, using induction, that for $n \in \mathbb{N}$, $n \geq 1$,

$$\mathbf{A}^n = \begin{bmatrix} \cos n\alpha & -\sin n\alpha \\ \sin n\alpha & \cos n\alpha \end{bmatrix}.$$

131 Problem A matrix $\mathbf{A} = [a_{ij}] \in M_n(\mathbb{R})$ is said to be *checkered* if $a_{ij} = 0$ when $(j - i)$ is odd. Prove that the sum and the product of two checkered matrices is checkered.

132 Problem Let $A \in M_3(\mathbb{R})$,

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}.$$

Prove that

$$A^n = \begin{bmatrix} 1 & n & \frac{n(n+1)}{2} \\ 0 & 1 & n \\ 0 & 0 & 1 \end{bmatrix}.$$

133 Problem Prove, by means of induction that for the following $n \times n$ we have

$$\begin{bmatrix} 1 & 1 & 1 & \dots & 1 \\ 0 & 1 & 1 & \dots & 1 \\ 0 & 0 & 1 & \dots & 1 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & 0 & 0 & \dots & 1 \end{bmatrix}^3 = \begin{bmatrix} 1 & 3 & 6 & \dots & \frac{n(n+1)}{2} \\ 0 & 1 & 3 & \dots & \frac{(n-1)n}{2} \\ 0 & 0 & 1 & \dots & \frac{(n-2)(n-1)}{2} \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & 0 & 0 & \dots & 1 \end{bmatrix}.$$

134 Problem Let $(A, B) \in (M_n(\mathbb{F}))^2$ and k be a positive integer such that $A^k = 0_n$. If $AB = B$ prove that $B = 0_n$.

135 Problem Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Demonstrate that

$$A^2 - (a+d)A + (ad - bc)I_2 = 0_2$$

136 Problem Let $A \in M_2(\mathbb{F})$ and let $k \in \mathbb{Z}, k > 2$. Prove that $A^k = 0_2$ if and only if $A^2 = 0_2$.

137 Problem Find all matrices $A \in M_2(\mathbb{R})$ such that $A^2 = 0_2$

138 Problem Find all matrices $A \in M_2(\mathbb{R})$ such that $A^2 = I_2$

139 Problem Find a solution $X \in M_2(\mathbb{R})$ for

$$X^2 - 2X = \begin{bmatrix} -1 & 0 \\ 6 & 3 \end{bmatrix}.$$

2.3 Trace and Transpose

140 Definition Let $A = [a_{ij}] \in M_n(\mathbb{F})$. Then the *trace* of A , denoted by $\text{tr}(A)$ is the sum of the diagonal elements of A , that is

$$\text{tr}(A) = \sum_{k=1}^n a_{kk}.$$

141 Theorem Let $A = [a_{ij}] \in M_n(\mathbb{F}), B = [b_{ij}] \in M_n(\mathbb{F})$. Then

$$\text{tr}(A + B) = \text{tr}(A) + \text{tr}(B), \quad (2.10)$$

$$\text{tr}(AB) = \text{tr}(BA). \quad (2.11)$$

Proof: The first assertion is trivial. To prove the second, observe that $AB = (\sum_{k=1}^n a_{ik}b_{kj})$ and $BA = (\sum_{k=1}^n b_{ik}a_{kj})$. Then

$$\text{tr}(AB) = \sum_{i=1}^n \sum_{k=1}^n a_{ik}b_{ki} = \sum_{k=1}^n \sum_{i=1}^n b_{ki}a_{ik} = \text{tr}(BA),$$

whence the theorem follows. \square

142 Example Let $A \in M_n(\mathbb{R})$. Shew that A can be written as the sum of two matrices whose trace is different from 0.

Solution: Write

$$\mathbf{A} = (\mathbf{A} - \alpha \mathbf{I}_n) + \alpha \mathbf{I}_n.$$

Now, $\text{tr}(\mathbf{A} - \alpha \mathbf{I}_n) = \text{tr}(\mathbf{A}) - n\alpha$ and $\text{tr}(\alpha \mathbf{I}_n) = n\alpha$. Thus it suffices to take $\alpha \neq \frac{\text{tr}(\mathbf{A})}{n}$, $\alpha \neq 0$. Since \mathbb{R} has infinitely many elements, we can find such an α .

143 Example Let \mathbf{A}, \mathbf{B} be square matrices of the same size and over the same field of characteristic 0. Is it possible that $\mathbf{AB} - \mathbf{BA} = \mathbf{I}_n$? Prove or disprove!

Solution: This is impossible. For if, taking traces on both sides

$$0 = \text{tr}(\mathbf{AB}) - \text{tr}(\mathbf{BA}) = \text{tr}(\mathbf{AB} - \mathbf{BA}) = \text{tr}(\mathbf{I}_n) = n$$

a contradiction, since $n > 0$.

144 Definition The *transpose* of a matrix of a matrix $\mathbf{A} = [a_{ij}] \in M_{m \times n}(\mathbb{F})$ is the matrix $\mathbf{A}^T = \mathbf{B} = [b_{ij}] \in M_{n \times m}(\mathbb{F})$, where $b_{ij} = a_{ji}$.

145 Example If

$$\mathbf{M} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix},$$

with entries in \mathbb{R} , then

$$\mathbf{M}^T = \begin{bmatrix} a & d & g \\ b & e & h \\ c & f & i \end{bmatrix}.$$

146 Theorem Let

$$\mathbf{A} = [a_{ij}] \in M_{m \times n}(\mathbb{F}), \mathbf{B} = [b_{ij}] \in M_{m \times n}(\mathbb{F}), \mathbf{C} = [c_{ij}] \in M_{n \times r}(\mathbb{F}), \alpha \in \mathbb{F}, u \in \mathbb{N}.$$

Then

$$\mathbf{A}^{TT} = \mathbf{A}, \tag{2.12}$$

$$(\mathbf{A} + \alpha \mathbf{B})^T = \mathbf{A}^T + \alpha \mathbf{B}^T, \tag{2.13}$$

$$(\mathbf{AC})^T = \mathbf{C}^T \mathbf{A}^T, \tag{2.14}$$

$$(\mathbf{A}^u)^T = (\mathbf{A}^T)^u. \tag{2.15}$$

Proof: The first two assertions are obvious, and the fourth follows from the third by using induction. To prove the third put $\mathbf{A}^T = (\alpha_{ij})$, $\alpha_{ij} = a_{ji}$, $\mathbf{C}^T = (\gamma_{ij})$, $\gamma_{ij} = c_{ji}$, $\mathbf{AC} = (\mathbf{u}_{ij})$ and $\mathbf{C}^T \mathbf{A}^T = (\mathbf{v}_{ij})$. Then

$$\mathbf{u}_{ij} = \sum_{k=1}^n \alpha_{ik} c_{kj} = \sum_{k=1}^n \alpha_{ki} \gamma_{jk} = \sum_{k=1}^n \gamma_{jk} \alpha_{ki} = \mathbf{v}_{ji},$$

whence the theorem follows. \square

147 Definition A square matrix $\mathbf{A} \in M_n(\mathbb{F})$ is *symmetric* if $\mathbf{A}^T = \mathbf{A}$. A matrix $\mathbf{B} \in M_n(\mathbb{F})$ is *skew-symmetric* if $\mathbf{B}^T = -\mathbf{B}$.

148 Example Let \mathbf{A}, \mathbf{B} be square matrices of the same size, with \mathbf{A} symmetric and \mathbf{B} skew-symmetric. Prove that the matrix $\mathbf{A}^2 \mathbf{B} \mathbf{A}^2$ is skew-symmetric.

Solution: We have

$$(\mathbf{A}^2 \mathbf{B} \mathbf{A}^2)^T = (\mathbf{A}^2)^T (\mathbf{B})^T (\mathbf{A}^2)^T = \mathbf{A}^2 (-\mathbf{B}) \mathbf{A}^2 = -\mathbf{A}^2 \mathbf{B} \mathbf{A}^2.$$

149 Theorem Let \mathbb{F} be a field of characteristic different from 2. Then any square matrix \mathbf{A} can be written as the sum of a symmetric and a skew-symmetric matrix.

Proof: Observe that

$$(\mathbf{A} + \mathbf{A}^T)^T = \mathbf{A}^T + \mathbf{A}^{TT} = \mathbf{A}^T + \mathbf{A},$$

and so $\mathbf{A} + \mathbf{A}^T$ is symmetric. Also,

$$(\mathbf{A} - \mathbf{A}^T)^T = \mathbf{A}^T - \mathbf{A}^{TT} = -(\mathbf{A} - \mathbf{A}^T),$$

and so $\mathbf{A} - \mathbf{A}^T$ is skew-symmetric. We only need to write \mathbf{A} as

$$\mathbf{A} = (2^{-1})(\mathbf{A} + \mathbf{A}^T) + (2^{-1})(\mathbf{A} - \mathbf{A}^T)$$

to prove the assertion. \square

150 Problem Write

$$\mathbf{A} = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 3 & 1 & 2 \end{bmatrix} \in M_3(\mathbb{R})$$

as the sum of two 3×3 matrices $\mathbf{E}_1, \mathbf{E}_2$, with $\text{tr}(\mathbf{E}_2) = 10$.

151 Problem Shew that there are no matrices $(\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}) \in (M_n(\mathbb{R}))^4$ such that

$$\mathbf{AC} + \mathbf{DB} = \mathbf{I}_n,$$

$$\mathbf{CA} + \mathbf{BD} = \mathbf{0}_n.$$

152 Problem Let $(\mathbf{A}, \mathbf{B}) \in (M_2(\mathbb{R}))^2$ be symmetric matrices. Must their product \mathbf{AB} be symmetric? Prove or disprove!

153 Problem Given square matrices $(\mathbf{A}, \mathbf{B}) \in (M_7(\mathbb{R}))^2$ such that $\text{tr}(\mathbf{A}^2) = \text{tr}(\mathbf{B}^2) = 1$, and

$$(\mathbf{A} - \mathbf{B})^2 = 3\mathbf{I}_7,$$

find $\text{tr}(\mathbf{BA})$.

154 Problem Consider the matrix $\mathbf{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in M_2(\mathbb{R})$.

Find necessary and sufficient conditions on a, b, c, d so that $\text{tr}(\mathbf{A}^2) = (\text{tr}(\mathbf{A}))^2$.

155 Problem Given a square matrix $\mathbf{A} \in M_4(\mathbb{R})$ such that $\text{tr}(\mathbf{A}^2) = -4$, and

$$(\mathbf{A} - \mathbf{I}_4)^2 = 3\mathbf{I}_4,$$

find $\text{tr}(\mathbf{A})$.

156 Problem Prove or disprove! If \mathbf{A}, \mathbf{B} are square matrices of the same size, then it is always true that $\text{tr}(\mathbf{AB}) = \text{tr}(\mathbf{A}) \text{tr}(\mathbf{B})$.

157 Problem Prove or disprove! If $(\mathbf{A}, \mathbf{B}, \mathbf{C}) \in (M_3(\mathbb{F}))^3$ then $\text{tr}(\mathbf{ABC}) = \text{tr}(\mathbf{BAC})$.

158 Problem Let \mathbf{A} be a square matrix. Prove that the matrix \mathbf{AA}^T is symmetric.

159 Problem Let \mathbf{A}, \mathbf{B} be square matrices of the same size, with \mathbf{A} symmetric and \mathbf{B} skew-symmetric. Prove that the matrix $\mathbf{AB} - \mathbf{BA}$ is symmetric.

160 Problem Let $\mathbf{A} \in M_n(\mathbb{F}), \mathbf{A} = [a_{ij}]$. Prove that $\text{tr}(\mathbf{AA}^T) = \sum_{i=1}^n \sum_{j=1}^n a_{ij}^2$.

161 Problem Let $\mathbf{X} \in M_n(\mathbb{R})$. Prove that if $\mathbf{XX}^T = \mathbf{0}_n$ then $\mathbf{X} = \mathbf{0}_n$.

162 Problem Let m, n, p be positive integers and $\mathbf{A} \in M_{m \times n}(\mathbb{R}), \mathbf{B} \in M_{n \times p}(\mathbb{R}), \mathbf{C} \in M_{p \times m}(\mathbb{R})$. Prove that $(\mathbf{BA})^T \mathbf{A} = (\mathbf{CA})^T \mathbf{A} \implies \mathbf{BA} = \mathbf{CA}$.

2.4 Special Matrices

163 Definition The *main diagonal* of a square matrix $\mathbf{A} = [a_{ij}] \in M_n(\mathbb{F})$ is the set $\{a_{ii} : i \leq n\}$. The *counter diagonal* of a square matrix $\mathbf{A} = [a_{ij}] \in M_n(\mathbb{F})$ is the set $\{a_{(n-i+1)i} : i \leq n\}$.

164 Example The main diagonal of the matrix

$$\mathbf{A} = \begin{bmatrix} 0 & 1 & 5 \\ 3 & 2 & 4 \\ 9 & 8 & 7 \end{bmatrix}$$

is the set $\{0, 2, 7\}$. The counter diagonal of \mathbf{A} is the set $\{5, 2, 9\}$.

165 Definition A square matrix is a *diagonal* matrix if every entry off its main diagonal is $0_{\mathbb{F}}$.

166 Example The matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$$

is a diagonal matrix.

167 Definition A square matrix is a *scalar* matrix if it is of the form αI_n for some scalar α .

168 Example The matrix

$$\mathbf{A} = \begin{bmatrix} 4 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 4 \end{bmatrix} = 4I_3$$

is a scalar matrix.

169 Definition $\mathbf{A} \in M_{m \times n}(\mathbb{F})$ is said to be *upper triangular* if

$$(\forall(i, j) \in \{1, 2, \dots, n\}^2), (i > j, a_{ij} = 0_{\mathbb{F}}),$$

that is, every element below the main diagonal is $0_{\mathbb{F}}$. Similarly, \mathbf{A} is *lower triangular* if

$$(\forall(i, j) \in \{1, 2, \dots, n\}^2), (i < j, a_{ij} = 0_{\mathbb{F}}),$$

that is, every element above the main diagonal is $0_{\mathbb{F}}$.

170 Example The matrix $\mathbf{A} \in M_{3 \times 4}(\mathbb{R})$ shewn is upper triangular and $\mathbf{B} \in M_4(\mathbb{R})$ is lower triangular.

$$\mathbf{A} = \begin{bmatrix} 1 & a & b & c \\ 0 & 2 & 3 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad \mathbf{B} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & a & 0 & 0 \\ 0 & 2 & 3 & 0 \\ 1 & 1 & t & 1 \end{bmatrix}$$

171 Definition The Kronecker delta δ_{ij} is defined by

$$\delta_{ij} = \begin{cases} 1_F & \text{if } i = j \\ 0_F & \text{if } i \neq j \end{cases}$$

172 Definition The set of matrices $E_{ij} \in M_{m \times n}(F)$, $E_{ij} = (e_{rs})$ such that $e_{ij} = 1_F$ and $e_{i'j'} = 0_F$, $(i', j') \neq (i, j)$ is called the set of *elementary matrices*. Observe that in fact $e_{rs} = \delta_{ir}\delta_{sj}$.

Elementary matrices have interesting effects when we pre-multiply and post-multiply a matrix by them.

173 Example Let

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \end{bmatrix}, \quad E_{23} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

Then

$$E_{23}A = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 9 & 10 & 11 & 12 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad AE_{23} = \begin{bmatrix} 0 & 0 & 2 \\ 0 & 0 & 6 \\ 0 & 0 & 10 \end{bmatrix}.$$

174 Theorem (Multiplication by Elementary Matrices) Let $E_{ij} \in M_{m \times n}(F)$ be an elementary matrix, and let $A \in M_{n \times m}(F)$. Then $E_{ij}A$ has as its i -th row the j -th row of A and 0_F 's everywhere else. Similarly, AE_{ij} has as its j -th column the i -th column of A and 0_F 's everywhere else.

Proof: Put $(\alpha_{uv}) = E_{ij}A$. To obtain $E_{ij}A$ we multiply the rows of E_{ij} by the columns of A . Now

$$\alpha_{uv} = \sum_{k=1}^n e_{uk} a_{kv} = \sum_{k=1}^n \delta_{ui}\delta_{kj} a_{kv} = \delta_{ui} a_{jv}.$$

Therefore, for $u \neq i$, $\alpha_{uv} = 0_F$, i.e., off of the i -th row the entries of $E_{ij}A$ are 0_F , and $\alpha_{iv} = \alpha_{jv}$, that is, the i -th row of $E_{ij}A$ is the j -th row of A . The case for AE_{ij} is similarly argued. \square

The following corollary is immediate.

175 Corollary Let $(E_{ij}, E_{kl}) \in (M_n(F))^2$, be square elementary matrices. Then

$$E_{ij}E_{kl} = \delta_{jk}E_{il}.$$

176 Example Let $M \in M_n(F)$ be a matrix such that $AM = MA$ for all matrices $A \in M_n(F)$. Demonstrate that $M = aI_n$ for some $a \in F$, i.e. M is a scalar matrix.

Solution: Assume $(s, t) \in \{1, 2, \dots, n\}^2$. Let $\mathbf{M} = (m_{ij})$ and $\mathbf{E}_{st} \in M_n(\mathbb{F})$. Since \mathbf{M} commutes with \mathbf{E}_{st} we have

$$\begin{bmatrix} 0 & 0 & \dots & 0 \\ \vdots & \vdots & \dots & \vdots \\ m_{t1} & m_{t2} & \dots & m_{tn} \\ \vdots & \vdots & \dots & \vdots \\ 0 & 0 & \dots & 0 \end{bmatrix} = \mathbf{E}_{st}\mathbf{M} = \mathbf{M}\mathbf{E}_{st} = \begin{bmatrix} 0 & 0 & \dots & m_{1s} & \dots & 0 \\ 0 & 0 & \vdots & m_{2s} & \vdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \vdots & m_{(n-1)s} & \vdots & 0 \\ 0 & 0 & \vdots & m_{ns} & \vdots & 0 \end{bmatrix}$$

For arbitrary $s \neq t$ we have shown that $m_{st} = m_{ts} = 0$, and that $m_{ss} = m_{tt}$. Thus the entries off the main diagonal are zero and the diagonal entries are all equal to one another, whence \mathbf{M} is a scalar matrix.

177 Definition Let $\lambda \in \mathbb{F}$ and $\mathbf{E}_{ij} \in M_n(\mathbb{F})$. A square matrix in $M_n(\mathbb{F})$ of the form $\mathbf{I}_n + \lambda \mathbf{E}_{ij}$ is called a *transvection*.

178 Example The matrix

$$\mathbf{T} = \mathbf{I}_3 + 4\mathbf{E}_{13} = \begin{bmatrix} 1 & 0 & 4 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

is a transvection. Observe that if

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 5 & 6 & 7 \\ 1 & 2 & 3 \end{bmatrix}$$

then

$$\mathbf{T}\mathbf{A} = \begin{bmatrix} 1 & 0 & 4 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 \\ 5 & 6 & 7 \\ 1 & 2 & 3 \end{bmatrix} = \begin{bmatrix} 5 & 9 & 13 \\ 5 & 6 & 7 \\ 1 & 2 & 3 \end{bmatrix},$$

that is, pre-multiplication by \mathbf{T} adds 4 times the third row of \mathbf{A} to the first row of \mathbf{A} . Similarly,

$$\mathbf{AT} = \begin{bmatrix} 1 & 1 & 1 \\ 5 & 6 & 7 \\ 1 & 2 & 3 \end{bmatrix} \begin{bmatrix} 1 & 0 & 4 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 5 \\ 5 & 6 & 27 \\ 1 & 2 & 7 \end{bmatrix},$$

that is, post-multiplication by \mathbf{T} adds 4 times the first column of \mathbf{A} to the third row of \mathbf{A} .

In general, we have the following theorem.

179 Theorem (Multiplication by a Transvection Matrix) Let $\mathbf{I}_n + \lambda \mathbf{E}_{ij} \in M_n(\mathbb{F})$ be a transvection and let $\mathbf{A} \in M_{n \times m}(\mathbb{F})$. Then $(\mathbf{I}_n + \lambda \mathbf{E}_{ij})\mathbf{A}$ adds the j -th row of \mathbf{A} to its i -th row and leaves the other rows unchanged.

Similarly, if $\mathbf{B} \in M_{p \times n}(\mathbb{F})$, $\mathbf{B}(\mathbf{I}_n + \lambda E_{ij})$ adds the i -th column of \mathbf{B} to the j -th column and leaves the other columns unchanged.

Proof: Simply observe that $(\mathbf{I}_n + \lambda E_{ij})\mathbf{A} = \mathbf{A} + \lambda E_{ij}\mathbf{A}$ and $\mathbf{A}(\mathbf{I}_n + \lambda E_{ij}) = \mathbf{A} + \lambda A E_{ij}$ and apply Theorem 174. \square

Observe that the particular transvection $\mathbf{I}_n + (\lambda - 1_F)E_{ii} \in M_n(\mathbb{F})$ consists of a diagonal matrix with 1_F 's everywhere on the diagonal, except on the ii -th position, where it has a λ .

180 Definition If $\lambda \neq 0_F$, we call the matrix $\mathbf{I}_n + (\lambda - 1_F)E_{ii}$ a *dilatation matrix*.

181 Example The matrix

$$\mathbf{S} = \mathbf{I}_3 + (4 - 1)\mathbf{E}_{11} = \begin{bmatrix} 4 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

is a dilatation matrix. Observe that if

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 5 & 6 & 7 \\ 1 & 2 & 3 \end{bmatrix}$$

then

$$\mathbf{S}\mathbf{A} = \begin{bmatrix} 4 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 \\ 5 & 6 & 7 \\ 1 & 2 & 3 \end{bmatrix} = \begin{bmatrix} 4 & 4 & 4 \\ 5 & 6 & 7 \\ 1 & 2 & 3 \end{bmatrix},$$

that is, pre-multiplication by \mathbf{S} multiplies by 4 the first row of \mathbf{A} . Similarly,

$$\mathbf{AS} = \begin{bmatrix} 1 & 1 & 1 \\ 5 & 6 & 7 \\ 1 & 2 & 3 \end{bmatrix} \begin{bmatrix} 4 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 4 & 1 & 1 \\ 20 & 6 & 7 \\ 4 & 2 & 3 \end{bmatrix},$$

that is, post-multiplication by \mathbf{S} multiplies by 4 the first column of \mathbf{A} .

182 Theorem (Multiplication by a Dilatation Matrix) Pre-multiplication of the matrix $\mathbf{A} \in M_{n \times m}(\mathbb{F})$ by the dilatation matrix $\mathbf{I}_n + (\lambda - 1_F)E_{ii} \in M_n(\mathbb{F})$ multiplies the i -th row of \mathbf{A} by λ and leaves the other rows of \mathbf{A} unchanged. Similarly, if $\mathbf{B} \in M_{p \times n}(\mathbb{F})$ post-multiplication of \mathbf{B} by $\mathbf{I}_n + (\lambda - 1_F)E_{ii}$ multiplies the i -th column of \mathbf{B} by λ and leaves the other columns of \mathbf{B} unchanged.

Proof: This follows by direct application of Theorem 179. \square

183 Definition We write \mathbf{I}_n^{ij} for the matrix which permutes the i -th row with the j -th row of the identity matrix. We call \mathbf{I}_n^{ij} a *transposition matrix*.

184 Example We have

$$\mathbf{I}_4^{(23)} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

If

$$\mathbf{A} = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \\ 13 & 14 & 15 & 16 \end{bmatrix},$$

then

$$\mathbf{I}_4^{(23)} \mathbf{A} = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 9 & 10 & 11 & 12 \\ 5 & 6 & 7 & 8 \\ 13 & 14 & 15 & 16 \end{bmatrix},$$

and

$$\mathbf{A} \mathbf{I}_4^{(23)} = \begin{bmatrix} 1 & 3 & 2 & 4 \\ 5 & 7 & 6 & 8 \\ 9 & 11 & 10 & 12 \\ 13 & 15 & 14 & 16 \end{bmatrix}.$$

185 Theorem (Multiplication by a Transposition Matrix) If $\mathbf{A} \in M_{n \times m}(\mathbb{F})$, then $\mathbf{I}_n^{ij} \mathbf{A}$ is the matrix obtained from \mathbf{A} permuting the i -th row with the j -th row of \mathbf{A} . Similarly, if $\mathbf{B} \in M_{p \times n}(\mathbb{F})$, then $\mathbf{B} \mathbf{I}_n^{ij}$ is the matrix obtained from \mathbf{B} by permuting the i -th column with the j -th column of \mathbf{B} .

Proof: We must prove that $\mathbf{I}_n^{ij} \mathbf{A}$ exchanges the i -th and j -th rows but leaves the other rows unchanged. But this follows upon observing that

$$\mathbf{I}_n^{ij} = \mathbf{I}_n + \mathbf{E}_{ij} + \mathbf{E}_{ji} - \mathbf{E}_{ii} - \mathbf{E}_{jj}$$

and appealing to Theorem 174.

□

186 Definition A square matrix which is either a transvection matrix, a dilatation matrix or a transposition matrix is called an *elimination matrix*.

 In a very loose way, we may associate pre-multiplication of a matrix \mathbf{A} by another matrix with an operation on the rows of \mathbf{A} , and post-multiplication of a matrix \mathbf{A} by another with an operation on the columns of \mathbf{A} .

187 Problem Consider the matrices

$$\mathbf{A} = \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ -1 & 1 & 1 & 1 \\ 1 & -1 & 1 & 1 \end{bmatrix}, \quad \mathbf{B} = \begin{bmatrix} 4 & -2 & 4 & 2 \\ 0 & 1 & 0 & 1 \\ 1 & 1 & -1 & 1 \\ 1 & -1 & 1 & 1 \end{bmatrix}.$$

Find a specific dilatation matrix \mathbf{D} , a specific transposition matrix \mathbf{P} , and a specific transvection matrix \mathbf{T} such that $\mathbf{B} = \mathbf{T}\mathbf{D}\mathbf{A}\mathbf{P}$.

188 Problem The matrix

$$\mathbf{A} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$$

is transformed into the matrix

$$\mathbf{B} = \begin{bmatrix} h-g & g & i \\ e-d & d & f \\ 2b-2a & 2a & 2c \end{bmatrix}$$

by a series of row and column operations. Find explicit permutation matrices \mathbf{P}, \mathbf{P}' , an explicit dilatation matrix \mathbf{D} , and an explicit transvection matrix \mathbf{T} such that

$$\mathbf{B} = \mathbf{D}\mathbf{P}\mathbf{A}\mathbf{P}'\mathbf{T}.$$

189 Problem Let $\mathbf{A} \in M_n(\mathbb{F})$. Prove that if

$$(\forall \mathbf{X} \in M_n(\mathbb{F})), (\text{tr}(\mathbf{AX}) = \text{tr}(\mathbf{BX})),$$

then $\mathbf{A} = \mathbf{B}$.

190 Problem Let $\mathbf{A} \in M_n(\mathbb{R})$ be such that

$$(\forall \mathbf{X} \in M_n(\mathbb{R})), ((\mathbf{XA})^2 = \mathbf{0}_n).$$

Prove that $\mathbf{A} = \mathbf{0}_n$.

2.5 Matrix Inversion

191 Definition Let $\mathbf{A} \in M_{m \times n}(\mathbb{F})$. Then \mathbf{A} is said to be *left-invertible* if $\exists \mathbf{L} \in M_{n \times m}(\mathbb{F})$ such that $\mathbf{LA} = \mathbf{I}_n$. \mathbf{A} is said to be *right-invertible* if $\exists \mathbf{R} \in M_{n \times m}(\mathbb{F})$ such that $\mathbf{RA} = \mathbf{I}_m$. A matrix is said to be *invertible* if it possesses a right and a left inverse. A matrix which is not invertible is said to be *singular*.

192 Example The matrix $\mathbf{A} \in M_{2 \times 3}(\mathbb{R})$

$$\mathbf{A} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$

has infinitely many right-inverses of the form

$$\mathbf{R}_{(x,y)} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ x & y \end{bmatrix}.$$

For

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ x & y \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix},$$

regardless of the values of x and y . Observe, however, that A does not have a left inverse, for

$$\begin{bmatrix} a & b \\ c & d \\ f & g \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} = \begin{bmatrix} a & b & 0 \\ c & d & 0 \\ f & g & 0 \end{bmatrix},$$

which will never give I_3 regardless of the values of a, b, c, d, f, g .

193 Example If $\lambda \neq 0$, then the scalar matrix λI_n is invertible, for

$$(\lambda I_n) (\lambda^{-1} I_n) = I_n = (\lambda^{-1} I_n) (\lambda I_n).$$

194 Example The zero matrix 0_n is singular.

195 Theorem Let $A \in M_n(\mathbb{F})$ a square matrix possessing a left inverse L and a right inverse R . Then $L = R$. Thus an invertible square matrix possesses a unique inverse.

Proof: Observe that we have $LA = I_n = AR$. Then

$$L = LI_n = L(AR) = (LA)R = I_n R = R.$$

□

196 Definition The subset of $M_n(\mathbb{F})$ of all invertible $n \times n$ matrices is denoted by $GL_n(\mathbb{F})$, read “the linear group of rank n over \mathbb{F} .”

197 Corollary Let $(A, B) \in (GL_n(\mathbb{F}))^2$. Then AB is also invertible and

$$(AB)^{-1} = B^{-1}A^{-1}.$$

Proof: Since AB is a square matrix, it suffices to notice that

$$B^{-1}A^{-1}(AB) = (AB)B^{-1}A^{-1} = I_n$$

and that since the inverse of a square matrix is unique, we must have $B^{-1}A^{-1} = (AB)^{-1}$. □

198 Corollary If a square matrix $S \in M_n(\mathbb{F})$ is invertible, then S^{-1} is also invertible and $(S^{-1})^{-1} = S$, in view of the uniqueness of the inverses of square matrices.

199 Corollary If a square matrix $A \in M_n(\mathbb{F})$ is invertible, then A^T is also invertible and $(A^T)^{-1} = (A^{-1})^T$.

Proof: We claim that $(A^T)^{-1} = (A^{-1})^T$. For

$$AA^{-1} = I_n \implies (AA^{-1})^T = I_n^T \implies (A^{-1})^T A^T = I_n,$$

where we have used Theorem 146. □

The next few theorems will prove that elimination matrices are invertible matrices.

200 Theorem (Invertibility of Transvections) Let $I_n + \lambda E_{ij} \in M_n(\mathbb{F})$ be a transvection, and let $i \neq j$. Then

$$(I_n + \lambda E_{ij})^{-1} = I_n - \lambda E_{ij}.$$

Proof: Expanding the product

$$\begin{aligned} (\mathbf{I}_n + \lambda \mathbf{E}_{ij})(\mathbf{I}_n - \lambda \mathbf{E}_{ij}) &= \mathbf{I}_n + \lambda \mathbf{E}_{ij} - \lambda \mathbf{E}_{ij} - \lambda^2 \mathbf{E}_{ij} \mathbf{E}_{ij} \\ &= \mathbf{I}_n - \lambda^2 \delta_{ij} \mathbf{E}_{ij} \\ &= \mathbf{I}_n, \end{aligned}$$

since $i \neq j$. \square

201 Example By Theorem 200, we have

$$\begin{bmatrix} 1 & 0 & 3 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & -3 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

202 Theorem (Invertibility of Dilatations) Let $\lambda \neq 0_F$. Then

$$(\mathbf{I}_n + (\lambda - 1_F) \mathbf{E}_{ii})^{-1} = \mathbf{I}_n + (\lambda^{-1} - 1_F) \mathbf{E}_{ii}.$$

Proof: Expanding the product

$$\begin{aligned} (\mathbf{I}_n + (\lambda - 1_F) \mathbf{E}_{ii})(\mathbf{I}_n + (\lambda^{-1} - 1_F) \mathbf{E}_{ii}) &= \mathbf{I}_n + (\lambda - 1_F) \mathbf{E}_{ii} \\ &\quad + (\lambda^{-1} - 1_F) \mathbf{E}_{ii} \\ &\quad + (\lambda - 1_F)(\lambda^{-1} - 1_F) \mathbf{E}_{ii} \\ &= \mathbf{I}_n + (\lambda - 1_F) \mathbf{E}_{ii} \\ &\quad + (\lambda^{-1} - 1_F) \mathbf{E}_{ii} \\ &\quad + (\lambda - 1_F)(\lambda^{-1} - 1_F) \mathbf{E}_{ii} \\ &= \mathbf{I}_n + (\lambda - 1_F + \lambda^{-1} - 1_F + 1_F \\ &\quad - \lambda - \lambda^{-1} - 1_F) \mathbf{E}_{ii} \\ &= \mathbf{I}_n, \end{aligned}$$

proving the assertion. \square

203 Example By Theorem 202, we have

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{2} & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Repeated applications of Theorem 202 gives the following corollary.

204 Corollary If $\lambda_1 \lambda_2 \lambda_3 \cdots \lambda_n \neq 0_F$, then

$$\begin{bmatrix} \lambda_1 & 0 & 0 & 0 & \cdots & 0 \\ 0 & \lambda_2 & 0 & 0 & \cdots & 0 \\ 0 & 0 & \lambda_3 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \cdots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & \lambda_n \end{bmatrix}$$

is invertible and

$$\begin{bmatrix} \lambda_1 & 0 & 0 & 0 & \cdots & 0 \\ 0 & \lambda_2 & 0 & 0 & \cdots & 0 \\ 0 & 0 & \lambda_3 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \cdots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & \lambda_n \end{bmatrix}^{-1} = \begin{bmatrix} \lambda_1^{-1} & 0 & 0 & 0 & \cdots & 0 \\ 0 & \lambda_2^{-1} & 0 & 0 & \cdots & 0 \\ 0 & 0 & \lambda_3^{-1} & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \cdots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & \lambda_n^{-1} \end{bmatrix}$$

205 Theorem (Invertibility of Permutation Matrices) Let $\tau \in S_n$ be a permutation. Then

$$(I_n^{ij})^{-1} = (I_n^{ij})^T.$$

Proof: By Theorem 185 pre-multiplication of I_n^{ij} by I_n^{ij} exchanges the i -th row with the j -th row, meaning that they return to the original position in I_n . Observe in particular that $I_n^{ij} = (I_n^{ij})^T$, and so $I_n^{ij}(I_n^{ij})^T = I_n$. \square

206 Example By Theorem 205, we have

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

207 Corollary If a square matrix can be represented as the product of elimination matrices of the same size, then it is invertible.

Proof: This follows from Corollary 197, and Theorems 200, 202, and 205. \square

208 Example Observe that

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 3 & 4 \\ 0 & 0 & 1 \end{bmatrix}$$

is the transvection $\mathbf{I}_3 + 4\mathbf{E}_{23}$ followed by the dilatation of the second column of this transvection by 3. Thus

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 3 & 4 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 4 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 1 \end{bmatrix},$$

and so

$$\begin{aligned} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 3 & 4 \\ 0 & 0 & 1 \end{bmatrix}^{-1} &= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 4 \\ 0 & 0 & 1 \end{bmatrix}^{-1} \\ &= \begin{bmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{3} & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -4 \\ 0 & 0 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{3} & -\frac{4}{3} \\ 0 & 0 & 1 \end{bmatrix}. \end{aligned}$$

209 Example We have

$$\begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix},$$

hence

$$\begin{aligned}
 \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}^{-1} &= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{-1} \\
 &= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \\
 &= \begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{bmatrix}.
 \end{aligned}$$

In the next section we will give a general method that will permit us to find the inverse of a square matrix when it exists.

210 Example Let $T = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in M_2(\mathbb{R})$. Then

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} = (ad - bc) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

Thus if $ad - bc \neq 0$ we see that

$$T^{-1} = \begin{bmatrix} \frac{d}{ad-bc} & -\frac{b}{ad-bc} \\ -\frac{c}{ad-bc} & \frac{a}{ad-bc} \end{bmatrix}.$$

211 Example If

$$A = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & -1 & 1 \end{bmatrix},$$

then \mathbf{A} is invertible, for an easy computation shews that

$$\mathbf{A}^2 = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & -1 & 1 \end{bmatrix}^2 = 4\mathbf{I}_4,$$

whence the inverse sought is

$$\mathbf{A}^{-1} = \frac{1}{4}\mathbf{A} = \frac{1}{4} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & -1 & 1 \end{bmatrix} = \begin{bmatrix} 1/4 & 1/4 & 1/4 & 1/4 \\ 1/4 & 1/4 & -1/4 & -1/4 \\ 1/4 & -1/4 & 1/4 & -1/4 \\ 1/4 & -1/4 & -1/4 & 1/4 \end{bmatrix}.$$

212 Example A matrix $\mathbf{A} \in M_n(\mathbb{R})$ is said to be *nilpotent* of index k if satisfies $\mathbf{A} \neq \mathbf{0}_n, \mathbf{A}^2 \neq \mathbf{0}_n, \dots, \mathbf{A}^{k-1} \neq \mathbf{0}_n$ and $\mathbf{A}^k = \mathbf{0}_n$ for integer $k \geq 1$. Prove that if \mathbf{A} is nilpotent, then $\mathbf{I}_n - \mathbf{A}$ is invertible and find its inverse.

Solution: To motivate the solution, think that instead of a matrix, we had a real number x with $|x| < 1$. Then the inverse of $1 - x$ is

$$(1 - x)^{-1} = \frac{1}{1 - x} = 1 + x + x^2 + x^3 + \dots.$$

Notice now that since $\mathbf{A}^k = \mathbf{0}_n$, then $\mathbf{A}^p = \mathbf{0}_n$ for $p \geq k$. We conjecture thus that

$$(\mathbf{I}_n - \mathbf{A})^{-1} = \mathbf{I}_n + \mathbf{A} + \mathbf{A}^2 + \dots + \mathbf{A}^{k-1}.$$

The conjecture is easily verified, as

$$\begin{aligned} (\mathbf{I}_n - \mathbf{A})(\mathbf{I}_n + \mathbf{A} + \mathbf{A}^2 + \dots + \mathbf{A}^{k-1}) &= \mathbf{I}_n + \mathbf{A} + \mathbf{A}^2 + \dots + \mathbf{A}^{k-1} \\ &\quad - (\mathbf{A} + \mathbf{A}^2 + \mathbf{A}^3 + \dots + \mathbf{A}^k) \\ &= \mathbf{I}_n \end{aligned}$$

and

$$\begin{aligned} (\mathbf{I}_n + \mathbf{A} + \mathbf{A}^2 + \dots + \mathbf{A}^{k-1})(\mathbf{I}_n - \mathbf{A}) &= \mathbf{I}_n - \mathbf{A} + \mathbf{A} - \mathbf{A}^2 + \mathbf{A}^3 - \mathbf{A}^4 + \dots \\ &\quad \dots + \mathbf{A}^{k-2} - \mathbf{A}^{k-1} + \mathbf{A}^{k-1} - \mathbf{A}^k \\ &= \mathbf{I}_n. \end{aligned}$$

213 Example The inverse of $\mathbf{A} \in M_3(\mathbb{Z}_5)$,

$$\mathbf{A} = \begin{bmatrix} \bar{2} & \bar{0} & \bar{0} \\ \bar{0} & \bar{3} & \bar{0} \\ \bar{0} & \bar{0} & \bar{4} \end{bmatrix}$$

is

$$\mathbf{A}^{-1} = \begin{bmatrix} \bar{3} & \bar{0} & \bar{0} \\ \bar{0} & \bar{2} & \bar{0} \\ \bar{0} & \bar{0} & \bar{4} \end{bmatrix},$$

as

$$\mathbf{AA}^{-1} = \begin{bmatrix} \bar{2} & \bar{0} & \bar{0} \\ \bar{0} & \bar{3} & \bar{0} \\ \bar{0} & \bar{0} & \bar{4} \end{bmatrix} \begin{bmatrix} \bar{3} & \bar{0} & \bar{0} \\ \bar{0} & \bar{2} & \bar{0} \\ \bar{0} & \bar{0} & \bar{4} \end{bmatrix} = \begin{bmatrix} \bar{1} & \bar{0} & \bar{0} \\ \bar{0} & \bar{1} & \bar{0} \\ \bar{0} & \bar{0} & \bar{1} \end{bmatrix}$$

214 Example (Putnam Exam, 1991) Let \mathbf{A} and \mathbf{B} be different $n \times n$ matrices with real entries. If $\mathbf{A}^3 = \mathbf{B}^3$ and $\mathbf{A}^2\mathbf{B} = \mathbf{B}^2\mathbf{A}$, prove that $\mathbf{A}^2 + \mathbf{B}^2$ is not invertible.

Solution: Observe that

$$(\mathbf{A}^2 + \mathbf{B}^2)(\mathbf{A} - \mathbf{B}) = \mathbf{A}^3 - \mathbf{A}^2\mathbf{B} + \mathbf{B}^2\mathbf{A} - \mathbf{B}^3 = \mathbf{0}_n.$$

If $\mathbf{A}^2 + \mathbf{B}^2$ were invertible, then we would have

$$\mathbf{A} - \mathbf{B} = (\mathbf{A}^2 + \mathbf{B}^2)^{-1}(\mathbf{A}^2 + \mathbf{B}^2)(\mathbf{A} - \mathbf{B}) = \mathbf{0}_n,$$

contradicting the fact that \mathbf{A} and \mathbf{B} are different matrices.

215 Lemma If $\mathbf{A} \in M_n(\mathbb{F})$ has a row or a column consisting all of $0_{\mathbb{F}}$'s, then \mathbf{A} is singular.

Proof: If \mathbf{A} were invertible, the (i, i) -th entry of the product of its inverse with \mathbf{A} would be $1_{\mathbb{F}}$. But if the i -th row of \mathbf{A} is all $0_{\mathbb{F}}$'s, then $\sum_{k=1}^n a_{ik}b_{ki} = 0_{\mathbb{F}}$, so the (i, i) entry of any matrix product with \mathbf{A} is $0_{\mathbb{F}}$, and never $1_{\mathbb{F}}$. \square

216 Problem The inverse of the matrix $\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 2 \\ 1 & 2 & 3 \end{bmatrix}$ is

the matrix $\mathbf{A}^{-1} = \begin{bmatrix} a & 1 & -1 \\ 1 & b & 1 \\ -1 & 1 & 0 \end{bmatrix}$. Determine a and b .

217 Problem A square matrix \mathbf{A} satisfies $\mathbf{A}^3 \neq \mathbf{0}_n$ but $\mathbf{A}^4 = \mathbf{0}_n$. Demonstrate that $\mathbf{I}_n + \mathbf{A}$ is invertible and find, with proof, its inverse.

218 Problem Prove or disprove! If $(\mathbf{A}, \mathbf{B}, \mathbf{A} + \mathbf{B}) \in (GL_n(\mathbb{R}))^3$ then $(\mathbf{A} + \mathbf{B})^{-1} = \mathbf{A}^{-1} + \mathbf{B}^{-1}$.

219 Problem Let $\mathbf{S} \in GL_n(\mathbb{F})$, $(\mathbf{A}, \mathbf{B}) \in (M_n(\mathbb{F}))^2$, and

k a positive integer. Prove that if $\mathbf{B} = \mathbf{S}\mathbf{A}\mathbf{S}^{-1}$ then $\mathbf{B}^k = \mathbf{S}\mathbf{A}^k\mathbf{S}^{-1}$.

220 Problem Let $\mathbf{A} \in M_n(\mathbb{F})$ and let k be a positive integer. Prove that \mathbf{A} is invertible if and only if \mathbf{A}^k is invertible.

221 Problem Let $\mathbf{S} \in GL_n(\mathbb{C})$, $\mathbf{A} \in M_n(\mathbb{C})$ with $\mathbf{A}^k = \mathbf{0}_n$ for some positive integer k . Prove that both $\mathbf{I}_n - \mathbf{S}\mathbf{A}\mathbf{S}^{-1}$ and $\mathbf{I}_n - \mathbf{S}^{-1}\mathbf{A}\mathbf{S}$ are invertible and find their inverses.

222 Problem Let \mathbf{A} and \mathbf{B} be square matrices of the same size such that both $\mathbf{A} - \mathbf{B}$ and $\mathbf{A} + \mathbf{B}$ are invertible. Put $\mathbf{C} = (\mathbf{A} - \mathbf{B})^{-1} + (\mathbf{A} + \mathbf{B})^{-1}$. Prove that

$$\mathbf{ACA} - \mathbf{ACB} + \mathbf{BCA} - \mathbf{BCB} = 2\mathbf{A}.$$

223 Problem Let $\mathbf{A}, \mathbf{B}, \mathbf{C}$ be non-zero square matrices of the same size over the same field and such that $\mathbf{ABC} = \mathbf{0}_n$. Prove that at least two of these three matrices are not invertible.

224 Problem Let $(\mathbf{A}, \mathbf{B}) \in (M_n(\mathbb{F}))^2$ be such that $\mathbf{A}^2 = \mathbf{B}^2 = (\mathbf{AB})^2 = \mathbf{I}_n$. Prove that $\mathbf{AB} = \mathbf{BA}$.

225 **Problem** Let $\mathbf{A} = \begin{bmatrix} a & b & b & \cdots & b \\ b & a & b & \cdots & b \\ b & b & a & \cdots & b \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ b & b & b & \cdots & a \end{bmatrix} \in M_n(\mathbb{F})$, $n > 1$, $(a, b) \in \mathbb{F}^2$. Determine when \mathbf{A} is invertible and find this inverse when it exists.

226 **Problem** Let $(\mathbf{A}, \mathbf{B}) \in (M_n(\mathbb{F}))^2$ be matrices such that

$\mathbf{A} + \mathbf{B} = \mathbf{AB}$. Demonstrate that $\mathbf{A} - \mathbf{I}_n$ is invertible and find this inverse.

227 **Problem** Let $\mathbf{S} \in GL_n(\mathbb{F})$ and $\mathbf{A} \in M_n(\mathbb{F})$. Prove that $\text{tr}(\mathbf{A}) = \text{tr}(\mathbf{SAS}^{-1})$.

228 **Problem** Let $\mathbf{A} \in M_n(\mathbb{R})$ be a skew-symmetric matrix. Prove that $\mathbf{I}_n + \mathbf{A}$ is invertible. Furthermore, if $\mathbf{B} = (\mathbf{I}_n - \mathbf{A})(\mathbf{I}_n + \mathbf{A})^{-1}$, prove that $\mathbf{B}^{-1} = \mathbf{B}^T$.

229 **Problem** A matrix $\mathbf{A} \in M_n(\mathbb{F})$ is said to be a *magic square* if the sum of each individual row equals the sum of each individual column. Assume that \mathbf{A} is a magic square and invertible. Prove that \mathbf{A}^{-1} is also a magic square.

2.6 Block Matrices

230 **Definition** Let $\mathbf{A} \in M_{m \times n}(\mathbb{F})$, $\mathbf{B} \in M_{m \times s}(\mathbb{F})$, $\mathbf{C} \in M_{r \times n}(\mathbb{F})$, $\mathbf{D} \in M_{r \times s}(\mathbb{F})$. We use the notation

$$\mathbf{L} = \left[\begin{array}{c|c} \mathbf{A} & \mathbf{B} \\ \hline \mathbf{C} & \mathbf{D} \end{array} \right]$$

for the *block matrix* $\mathbf{L} \in M_{(m+r) \times (n+s)}(\mathbb{F})$.

 If $(\mathbf{A}, \mathbf{A}') \in (M_m(\mathbb{F}))^2$, $(\mathbf{B}, \mathbf{B}') \in (M_{m \times n}(\mathbb{F}))^2$, $(\mathbf{C}, \mathbf{C}') \in (M_{n \times m}(\mathbb{F}))^2$, $(\mathbf{D}, \mathbf{D}') \in (M_m(\mathbb{F}))^2$, and

$$\mathbf{S} = \left[\begin{array}{c|c} \mathbf{A} & \mathbf{B} \\ \hline \mathbf{C} & \mathbf{D} \end{array} \right], \quad \mathbf{T} = \left[\begin{array}{c|c} \mathbf{A}' & \mathbf{B}' \\ \hline \mathbf{C}' & \mathbf{D}' \end{array} \right],$$

then it is easy to verify that

$$\mathbf{ST} = \left[\begin{array}{c|c} \mathbf{AA}' + \mathbf{BC}' & \mathbf{AB}' + \mathbf{BD}' \\ \hline \mathbf{CA}' + \mathbf{DC}' & \mathbf{CB}' + \mathbf{DD}' \end{array} \right].$$

231 **Lemma** Let $\mathbf{L} \in M_{(m+r) \times (m+r)}(\mathbb{F})$ be the square block matrix

$$\mathbf{L} = \left[\begin{array}{c|c} \mathbf{A} & \mathbf{C} \\ \hline \mathbf{0}_{r \times m} & \mathbf{B} \end{array} \right],$$

with square matrices $\mathbf{A} \in M_m(\mathbb{F})$ and $\mathbf{B} \in M_r(\mathbb{F})$, and a matrix $\mathbf{C} \in M_{m \times r}(\mathbb{F})$. Then \mathbf{L} is invertible if and only if \mathbf{A} and \mathbf{B} are, in which case

$$\mathbf{L}^{-1} = \left[\begin{array}{c|c} \mathbf{A}^{-1} & -\mathbf{A}^{-1}\mathbf{CB}^{-1} \\ \hline \mathbf{0}_{r \times m} & \mathbf{B}^{-1} \end{array} \right]$$

Proof: Assume first that \mathbf{A} , and \mathbf{B} are invertible. Direct calculation yields

$$\begin{aligned} \left[\begin{array}{c|c} \mathbf{A} & \mathbf{C} \\ \hline \mathbf{0}_{r \times m} & \mathbf{B} \end{array} \right] \left[\begin{array}{c|c} \mathbf{A}^{-1} & -\mathbf{A}^{-1}\mathbf{C}\mathbf{B}^{-1} \\ \hline \mathbf{0}_{r \times m} & \mathbf{B}^{-1} \end{array} \right] &= \left[\begin{array}{c|c} \mathbf{AA}^{-1} & -\mathbf{AA}^{-1}\mathbf{C}\mathbf{B}^{-1} + \mathbf{CB}^{-1} \\ \hline \mathbf{0}_{r \times m} & \mathbf{BB}^{-1} \end{array} \right] \\ &= \left[\begin{array}{c|c} \mathbf{I}_m & \mathbf{0}_{m \times r} \\ \hline \mathbf{0}_{r \times m} & \mathbf{I}_r \end{array} \right] \\ &= \mathbf{I}_{m+r}. \end{aligned}$$

Assume now that \mathbf{L} is invertible, $\mathbf{L}^{-1} = \left[\begin{array}{c|c} \mathbf{E} & \mathbf{H} \\ \hline \mathbf{J} & \mathbf{K} \end{array} \right]$, with $\mathbf{E} \in M_m(\mathbb{F})$ and $\mathbf{K} \in M_r(\mathbb{F})$, but that, say, \mathbf{B} is singular. Then

$$\begin{aligned} \left[\begin{array}{c|c} \mathbf{I}_m & \mathbf{0}_{m \times r} \\ \hline \mathbf{0}_{r \times m} & \mathbf{I}_r \end{array} \right] &= \mathbf{LL}^{-1} \\ &= \left[\begin{array}{c|c} \mathbf{A} & \mathbf{C} \\ \hline \mathbf{0}_{r \times m} & \mathbf{B} \end{array} \right] \left[\begin{array}{c|c} \mathbf{E} & \mathbf{H} \\ \hline \mathbf{J} & \mathbf{K} \end{array} \right] \\ &= \left[\begin{array}{c|c} \mathbf{AE} + \mathbf{CJ} & \mathbf{AH} + \mathbf{BK} \\ \hline \mathbf{BJ} & \mathbf{BK} \end{array} \right], \end{aligned}$$

which gives $\mathbf{BK} = \mathbf{I}_r$, i.e., \mathbf{B} is invertible, a contradiction. \square

2.7 Rank of a Matrix

232 Definition Let $(\mathbf{A}, \mathbf{B}) \in (M_{m \times n}(\mathbb{F}))^2$. We say that \mathbf{A} is *row-equivalent* to \mathbf{B} if there exists a matrix $\mathbf{R} \in GL_m(\mathbb{F})$ such that $\mathbf{B} = \mathbf{RA}$. Similarly, we say that \mathbf{A} is *column-equivalent* to \mathbf{B} if there exists a matrix $\mathbf{C} \in GL_n(\mathbb{F})$ such that $\mathbf{B} = \mathbf{AC}$. We say that \mathbf{A} and \mathbf{B} are *equivalent* if $\exists (\mathbf{P}, \mathbf{Q}) \in GL_m(\mathbb{F}) \times GL_n(\mathbb{F})$ such that $\mathbf{B} = \mathbf{PAQ}$.

233 Theorem Row equivalence, column equivalence, and equivalence are equivalence relations.

Proof: We prove the result for row equivalence. The result for column equivalence, and equivalence are analogously proved.

Since $\mathbf{I}_m \in GL_m(\mathbb{F})$ and $\mathbf{A} = \mathbf{I}_m \mathbf{A}$, row equivalence is a reflexive relation. Assume $(\mathbf{A}, \mathbf{B}) \in (M_{m \times n}(\mathbb{F}))^2$ and that $\exists \mathbf{P} \in GL_m(\mathbb{F})$ such that $\mathbf{B} = \mathbf{PA}$. Then $\mathbf{A} = \mathbf{P}^{-1}\mathbf{B}$ and since $\mathbf{P}^{-1} \in GL_m(\mathbb{F})$, we see that row equivalence is a symmetric relation. Finally assume $(\mathbf{A}, \mathbf{B}, \mathbf{C}) \in (M_{m \times n}(\mathbb{F}))^3$ and that $\exists \mathbf{P} \in GL_m(\mathbb{F})$, $\exists \mathbf{P}' \in GL_n(\mathbb{F})$ such that $\mathbf{A} = \mathbf{PB}$, $\mathbf{B} = \mathbf{P}'\mathbf{C}$. Then $\mathbf{A} = \mathbf{PP}'\mathbf{C}$. But $\mathbf{PP}' \in GL_m(\mathbb{F})$ in view of Corollary 197. This completes the proof. \square

234 Theorem Let $\mathbf{A} \in M_{m \times n}(\mathbb{F})$. Then \mathbf{A} can be reduced, by means of pre-multiplication and post-multiplication

by elimination matrices, to a unique matrix of the form

$$\mathbf{D}_{m,n,r} = \left[\begin{array}{c|c} \mathbf{I}_r & \mathbf{0}_{r \times (n-r)} \\ \hline \mathbf{0}_{(m-r) \times r} & \mathbf{0}_{(m-r) \times (n-r)} \end{array} \right], \quad (2.16)$$

called the *Hermite normal form* of \mathbf{A} . Thus there exist $\mathbf{P} \in \text{GL}_m(\mathbb{F})$, $\mathbf{Q} \in \text{GL}_n(\mathbb{F})$ such that $\mathbf{D}_{m,n,r} = \mathbf{PAQ}$. The integer $r \geq 0$ is called the *rank* of the matrix \mathbf{A} which we denote by $\text{rank}(\mathbf{A})$.

Proof: If \mathbf{A} is the $m \times n$ zero matrix, then the theorem is obvious, taking $r = 0$. Assume hence that \mathbf{A} is not the zero matrix. We proceed as follows using the Gauß-Jordan Algorithm.

GJ-1 Since \mathbf{A} is a non-zero matrix, it has a non-zero column. By means of permutation matrices we move this column to the first column.

GJ-2 Since this column is a non-zero column, it must have an entry $a \neq 0_{\mathbb{F}}$. Again, by means of permutation matrices, we move the row on which this entry is to the first row.

GJ-3 By means of a dilatation matrix with scale factor a^{-1} , we make this new $(1,1)$ entry into a $1_{\mathbb{F}}$.

GJ-4 By means of transvections (adding various multiples of row 1 to the other rows) we now annihilate every entry below the entry $(1,1)$.

This process ends up in a matrix of the form

$$\mathbf{P}_1 \mathbf{A} \mathbf{Q}_1 = \left[\begin{array}{c|ccccc} 1_{\mathbb{F}} & * & * & \cdots & * \\ \hline 0_{\mathbb{F}} & b_{22} & b_{23} & \cdots & b_{2n} \\ 0_{\mathbb{F}} & b_{32} & b_{33} & \cdots & b_{3n} \\ 0_{\mathbb{F}} & \vdots & \vdots & \cdots & \vdots \\ 0_{\mathbb{F}} & b_{m2} & b_{m3} & \cdots & b_{mn} \end{array} \right]. \quad (2.17)$$

Here the asterisks represent unknown entries. Observe that the \mathbf{b} 's form a $(m-1) \times (n-1)$ matrix.

GJ-5 Apply GJ-1 through GJ-4 to the matrix of the \mathbf{b} 's.

Observe that this results in a matrix of the form

$$\mathbf{P}_2 \mathbf{A} \mathbf{Q}_2 = \left[\begin{array}{c|ccccc} 1_{\mathbb{F}} & * & * & \cdots & * \\ \hline 0_{\mathbb{F}} & 1_{\mathbb{F}} & * & \cdots & * \\ \hline 0_{\mathbb{F}} & 0_{\mathbb{F}} & c_{33} & \cdots & c_{3n} \\ 0_{\mathbb{F}} & \vdots & \vdots & \cdots & \vdots \\ 0_{\mathbb{F}} & 0_{\mathbb{F}} & c_{m3} & \cdots & c_{mn} \end{array} \right]. \quad (2.18)$$

GJ-6 Add the appropriate multiple of column 1 to column 2, that is, apply a transvection, in order to make the entry in the $(1,2)$ position $0_{\mathbb{F}}$.

This now gives a matrix of the form

$$\mathbf{P}_3 \mathbf{A} \mathbf{Q}_3 = \left[\begin{array}{cc|ccc} 1_{\mathbb{F}} & 0_{\mathbb{F}} & * & \cdots & * \\ 0_{\mathbb{F}} & 1_{\mathbb{F}} & * & \cdots & * \\ \hline 0_{\mathbb{F}} & 0_{\mathbb{F}} & c_{33} & \cdots & c_{3n} \\ 0_{\mathbb{F}} & \vdots & \vdots & \cdots & \vdots \\ 0_{\mathbb{F}} & 0_{\mathbb{F}} & c_{m3} & \cdots & c_{mn} \end{array} \right]. \quad (2.19)$$

The matrix of the c 's has size $(m - 2) \times (n - 2)$.

GJ-7 Apply GJ-1 through GJ-6 to the matrix of the c 's, etc.

Observe that this process eventually stops, and in fact, it is clear that $\text{rank}(\mathbf{A}) \leq \min(m, n)$.

Suppose now that \mathbf{A} were equivalent to a matrix $\mathbf{D}_{m,n,s}$ with $s > r$. Since matrix equivalence is an equivalence relation, $\mathbf{D}_{m,n,s}$ and $\mathbf{D}_{m,n,r}$ would be equivalent, and so there would be $\mathbf{R} \in \text{GL}_m(\mathbb{F})$, $\mathbf{S} \in \text{GL}_n(\mathbb{F})$, such that $\mathbf{R}\mathbf{D}_{m,n,r}\mathbf{S} = \mathbf{D}_{m,n,s}$, that is, $\mathbf{R}\mathbf{D}_{m,n,r} = \mathbf{D}_{m,n,s}\mathbf{S}^{-1}$. Partition \mathbf{R} and \mathbf{S}^{-1} as follows

$$\mathbf{R} = \left[\begin{array}{c|c} \mathbf{R}_{11} & \mathbf{R}_{12} \\ \hline \mathbf{R}_{21} & \mathbf{R}_{22} \end{array} \right], \quad \mathbf{S}^{-1} = \left[\begin{array}{c|cc} \mathbf{S}_{11} & \mathbf{S}_{12} & \mathbf{S}_{13} \\ \hline \mathbf{S}_{21} & \mathbf{S}_{22} & \mathbf{S}_{23} \\ \mathbf{S}_{31} & \mathbf{S}_{32} & \mathbf{S}_{33} \end{array} \right],$$

with $(\mathbf{R}_{11}, \mathbf{S}_{11})^2 \in (\mathbf{M}_r(\mathbb{F}))^2$, $\mathbf{S}_{22} \in \mathbf{M}_{(s-r) \times (s-r)}(\mathbb{F})$. We have

$$\mathbf{R}\mathbf{D}_{m,n,r} = \left[\begin{array}{c|c} \mathbf{R}_{11} & \mathbf{R}_{12} \\ \hline \mathbf{R}_{21} & \mathbf{R}_{22} \end{array} \right] \left[\begin{array}{c|c} \mathbf{I}_r & \mathbf{0}_{(m-r) \times r} \\ \hline \mathbf{0}_{(m-r) \times r} & \mathbf{0}_{r \times (m-r)} \end{array} \right] = \left[\begin{array}{c|c} \mathbf{R}_{11} & \mathbf{0}_{(m-r) \times r} \\ \hline \mathbf{R}_{21} & \mathbf{0}_{r \times (m-r)} \end{array} \right],$$

and

$$\begin{aligned} \mathbf{D}_{m,n,s}\mathbf{S}^{-1} &= \left[\begin{array}{c|cc} \mathbf{I}_r & \mathbf{0}_{r \times (s-r)} & \mathbf{0}_{r \times (n-s)} \\ \hline \mathbf{0}_{(s-r) \times r} & \mathbf{I}_{s-r} & \mathbf{0}_{(s-r) \times (n-s)} \\ \mathbf{0}_{(m-s) \times r} & \mathbf{0}_{(m-s) \times (s-r)} & \mathbf{0}_{(m-s) \times (n-s)} \end{array} \right] \left[\begin{array}{c|cc} \mathbf{S}_{11} & \mathbf{S}_{12} & \mathbf{S}_{13} \\ \hline \mathbf{S}_{21} & \mathbf{S}_{22} & \mathbf{S}_{23} \\ \mathbf{S}_{31} & \mathbf{S}_{32} & \mathbf{S}_{33} \end{array} \right] \\ &= \left[\begin{array}{c|ccc} \mathbf{S}_{11} & \mathbf{S}_{12} & \mathbf{S}_{13} & & \\ \hline \mathbf{S}_{21} & \mathbf{S}_{22} & \mathbf{S}_{23} & & \\ \mathbf{0}_{(m-s) \times r} & \mathbf{0}_{(m-s) \times (s-r)} & \mathbf{0}_{(m-s) \times (n-s)} & & \end{array} \right]. \end{aligned}$$

Since we are assuming

$$\left[\begin{array}{c|c} \mathbf{R}_{11} & \mathbf{0}_{(m-r) \times r} \\ \hline \mathbf{R}_{21} & \mathbf{0}_{r \times (m-r)} \end{array} \right] = \left[\begin{array}{c|ccc} \mathbf{S}_{11} & \mathbf{S}_{12} & \mathbf{S}_{13} & & \\ \hline \mathbf{S}_{21} & \mathbf{S}_{22} & \mathbf{S}_{23} & & \\ \mathbf{0}_{(m-s) \times r} & \mathbf{0}_{(m-s) \times (s-r)} & \mathbf{0}_{(m-s) \times (n-s)} & & \end{array} \right],$$

we must have $S_{12} = \mathbf{0}_{r \times (s-r)}$, $S_{13} = \mathbf{0}_{r \times (n-s)}$, $S_{22} = \mathbf{0}_{(s-r) \times (s-r)}$, $S_{23} = \mathbf{0}_{(s-r) \times (n-s)}$. Hence

$$S^{-1} = \left[\begin{array}{c|cc} S_{11} & \mathbf{0}_{r \times (s-r)} & \mathbf{0}_{r \times (n-s)} \\ \hline S_{21} & \mathbf{0}_{(s-r) \times (s-r)} & \mathbf{0}_{(s-r) \times (n-s)} \\ S_{31} & S_{32} & S_{33} \end{array} \right].$$

The matrix

$$\left[\begin{array}{c|c} \mathbf{0}_{(s-r) \times (s-r)} & \mathbf{0}_{(s-r) \times (n-s)} \\ \hline S_{32} & S_{33} \end{array} \right]$$

is non-invertible, by virtue of Lemma 215. This entails that S^{-1} is non-invertible by virtue of Lemma 231. This is a contradiction, since S is assumed invertible, and hence S^{-1} must also be invertible. \square

 Albeit the rank of a matrix is unique, the matrices P and Q appearing in Theorem 234 are not necessarily unique. For example, the matrix

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}$$

has rank 2, the matrix

$$\begin{bmatrix} 1 & 0 & x \\ 0 & 1 & y \\ 0 & 0 & 1 \end{bmatrix}$$

is invertible, and an easy computation shews that

$$\begin{bmatrix} 1 & 0 & x \\ 0 & 1 & y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix},$$

regardless of the values of x and y .

235 Corollary Let $A \in M_{m \times n}(\mathbb{F})$. Then $\text{rank}(A) = \text{rank}(A^T)$.

Proof: Let $P, Q, D_{m,n,r}$ as in Theorem 234. Observe that P^T, Q^T are invertible. Then

$$PAQ = D_{m,n,r} \implies Q^T A^T P^T = D_{m,n,r}^T = D_{n,m,r},$$

and since this last matrix has the same number of $1_{\mathbb{F}}$'s as $D_{m,n,r}$, the corollary is proven. \square

236 Example Show that

$$\mathbf{A} = \begin{bmatrix} 0 & 2 & 3 \\ 0 & 1 & 0 \end{bmatrix}$$

has $\text{rank } (\mathbf{A}) = 2$ and find invertible matrices $\mathbf{P} \in \text{GL}_2(\mathbb{R})$ and $\mathbf{Q} \in \text{GL}_3(\mathbb{R})$ such that

$$\mathbf{PAQ} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}.$$

Solution: We first transpose the first and third columns by effecting

$$\begin{bmatrix} 0 & 2 & 3 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 3 & 2 & 0 \\ 0 & 1 & 0 \end{bmatrix}.$$

We now subtract twice the second row from the first, by effecting

$$\begin{bmatrix} 1 & -2 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 3 & 2 & 0 \\ 0 & 1 & 0 \end{bmatrix} = \begin{bmatrix} 3 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}.$$

Finally, we divide the first row by 3,

$$\begin{bmatrix} 1/3 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 3 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}.$$

We conclude that

$$\begin{bmatrix} 1/3 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & -2 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 2 & 3 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix},$$

from where we may take

$$\mathbf{P} = \begin{bmatrix} 1/3 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & -2 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1/3 & -2/3 \\ 0 & 1 \end{bmatrix}$$

and

$$\mathbf{Q} = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}.$$

In practice it is easier to do away with the multiplication by elimination matrices and perform row and column operations on the *augmented* $(m + n) \times (m + n)$ matrix

$$\left[\begin{array}{c|c} I_n & 0_{n \times m} \\ \hline A & I_m \end{array} \right].$$

237 Definition Denote the rows of a matrix $A \in M_{m \times n}(\mathbb{F})$ by R_1, R_2, \dots, R_m , and its columns by C_1, C_2, \dots, C_n . The elimination operations will be denoted as follows.

- Exchanging the i -th row with the j -th row, which we denote by $R_i \leftrightarrow R_j$, and the s -th column by the t -th column by $C_s \leftrightarrow C_t$.
- A dilatation of the i -th row by a non-zero scalar $\alpha \in \mathbb{F} \setminus \{0_{\mathbb{F}}\}$, we will denote by $\alpha R_i \rightarrow R_i$. Similarly, $\beta C_j \rightarrow C_j$ denotes the dilatation of the j -th column by the non-zero scalar β .
- A transvection on the rows will be denoted by $R_i + \alpha R_j \rightarrow R_i$, and one on the columns by $C_s + \beta C_t \rightarrow C_s$.

238 Example Find the Hermite normal form of

$$A = \begin{bmatrix} -1 & 0 \\ 0 & 0 \\ 1 & 1 \\ 1 & 2 \end{bmatrix}.$$

Solution: First observe that $\text{rank}(A) \leq \min(4, 2) = 2$, so the rank can be either 1 or 2 (why not 0?). Form the augmented matrix

$$\left[\begin{array}{cc|cccc} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ \hline -1 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 & 1 & 0 \\ 1 & 2 & 0 & 0 & 0 & 1 \end{array} \right].$$

Perform $R_5 + R_3 \rightarrow R_5$ and $R_6 + R_3 \rightarrow R_6$ successively, obtaining

$$\left[\begin{array}{cc|cccc} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ \hline -1 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 & 1 & 0 \\ 0 & 2 & 1 & 0 & 0 & 1 \end{array} \right].$$

Perform $R_6 - 2R_5 \rightarrow R_6$

$$\left[\begin{array}{cc|cccc} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ \hline -1 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & -1 & 0 & -2 & 1 \end{array} \right].$$

Perform $R_4 \leftrightarrow R_5$

$$\left[\begin{array}{cc|cccc} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ \hline -1 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 & -2 & 1 \end{array} \right].$$

Finally, perform $-R_3 \rightarrow R_3$

$$\left[\begin{array}{cc|cccc} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ \hline 1 & 0 & -1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 & -2 & 1 \end{array} \right].$$

We conclude that

$$\begin{bmatrix} -1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & -2 & 1 \end{bmatrix} \begin{bmatrix} -1 & 0 \\ 0 & 0 \\ 1 & 1 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}.$$

239 Theorem Let $A \in M_{m \times n}(\mathbb{F})$, $B \in M_{n \times p}(\mathbb{F})$. Then

$$\text{rank}(AB) \leq \min(\text{rank}(A), \text{rank}(B)).$$

Proof: We prove that $\text{rank}(\mathbf{A}) \geq \text{rank}(\mathbf{AB})$. The proof that $\text{rank}(\mathbf{B}) \geq \text{rank}(\mathbf{AB})$ is similar and left to the reader. Put $r = \text{rank}(\mathbf{A})$, $s = \text{rank}(\mathbf{AB})$. There exist matrices $P \in \text{GL}_m(\mathbb{F})$, $Q \in \text{GL}_n(\mathbb{F})$, $S \in \text{GL}_m(\mathbb{F})$, $T \in \text{GL}_p(\mathbb{F})$ such that

$$P\mathbf{A}Q = D_{m,n,r}, \quad S\mathbf{AB}T = D_{m,p,s}.$$

Now

$$D_{m,p,s} = S\mathbf{AB}T = SP^{-1}D_{m,n,r}Q^{-1}BT,$$

from where it follows that

$$PS^{-1}D_{m,p,s} = D_{m,n,r}Q^{-1}BT.$$

Now the proof is analogous to the uniqueness proof of Theorem 234. Put $U = PS^{-1} \in \text{GL}_m(\mathbb{R})$ and $V = Q^{-1}BT \in M_{n \times p}(\mathbb{F})$, and partition U and V as follows:

$$U = \begin{bmatrix} U_{11} & U_{12} \\ \hline U_{21} & U_{22} \end{bmatrix}, \quad V = \begin{bmatrix} V_{11} & V_{12} \\ \hline V_{21} & V_{22} \end{bmatrix},$$

with $U_{11} \in M_s(\mathbb{F})$, $V_{11} \in M_r(\mathbb{F})$. Then

$$UD_{m,p,s} = \begin{bmatrix} U_{11} & U_{12} \\ \hline U_{21} & U_{22} \end{bmatrix} \begin{bmatrix} I_s & & \\ \hline & 0_{s \times (p-s)} & \\ 0_{(m-s) \times s} & & 0_{(m-s) \times (p-s)} \end{bmatrix} \in M_{m \times p}(\mathbb{F}),$$

and

$$D_{m,p,s}V = \begin{bmatrix} I_r & & \\ \hline & 0_{r \times (n-r)} & \\ 0_{(m-r) \times r} & & 0_{(m-r) \times (n-r)} \end{bmatrix} \begin{bmatrix} V_{11} & V_{12} \\ \hline V_{21} & V_{22} \end{bmatrix} \in M_{m \times p}(\mathbb{F}).$$

From the equality of these two $m \times p$ matrices, it follows that

$$\begin{bmatrix} U_{11} & & 0_{s \times (p-s)} \\ \hline U_{21} & & 0_{(m-s) \times (p-s)} \end{bmatrix} = \begin{bmatrix} V_{11} & & V_{12} \\ \hline 0_{(m-r) \times r} & & 0_{(m-r) \times (n-r)} \end{bmatrix}.$$

If $s > r$ then (i) U_{11} would have at least one row of $0_{\mathbb{F}}$'s meaning that U_{11} is non-invertible by Lemma 215. (ii) $U_{21} = 0_{(m-s) \times s}$. Thus from (i) and (ii) and from Lemma 231, U is not invertible, which is a contradiction. \square

240 Corollary Let $A \in M_{m \times n}(\mathbb{F})$, $B \in M_{n \times p}(\mathbb{F})$. If A is invertible then $\text{rank}(\mathbf{AB}) = \text{rank}(\mathbf{B})$. If B is invertible then $\text{rank}(\mathbf{AB}) = \text{rank}(\mathbf{A})$.

Proof: Using Theorem 239, if A is invertible

$$\text{rank}(\mathbf{AB}) \leq \text{rank}(\mathbf{B}) = \text{rank}(\mathbf{A}^{-1}\mathbf{AB}) \leq \text{rank}(\mathbf{AB}),$$

and so $\text{rank}(\mathbf{B}) = \text{rank}(\mathbf{AB})$. A similar argument works when B is invertible.

\square

241 Example Study the various possibilities for the rank of the matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ b+c & c+a & a+b \\ bc & ca & ab \end{bmatrix}.$$

Solution: Performing $R_2 - (b+c)R_1 \rightarrow R_2$ and $R_3 - bcR_1 \rightarrow R_3$, we find

$$\begin{bmatrix} 1 & 1 & 1 \\ 0 & a-b & a-c \\ 0 & 0 & (b-c)(a-c) \end{bmatrix}.$$

Performing $C_2 - C_1 \rightarrow C_2$ and $C_3 - C_1 \rightarrow C_3$, we find

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & a-b & a-c \\ 0 & 0 & (b-c)(a-c) \end{bmatrix}.$$

We now examine the various ways of getting rows consisting only of 0's. If $a = b = c$, the last two rows are 0-rows and so $\text{rank } (\mathbf{A}) = 1$. If exactly two of a, b, c are equal, the last row is a 0-row, but the middle one is not, and so $\text{rank } (\mathbf{A}) = 2$ in this case. If none of a, b, c are equal, then the rank is clearly 3.

242 Problem On a symmetric matrix $\mathbf{A} \in M_n(\mathbb{R})$ with $n \geq 3$,

$$R_3 - 3R_1 \rightarrow R_3$$

successively followed by

$$C_3 - 3C_1 \rightarrow C_3$$

are performed. Is the resulting matrix still symmetric?

$$\begin{bmatrix} 1 & a & 1 & b \\ a & 1 & b & 1 \\ 1 & b & 1 & a \\ b & 1 & a & 1 \end{bmatrix}$$

243 Problem Find the rank of

$$\begin{bmatrix} a+1 & a+2 & a+3 & a+4 & a+5 \\ a+2 & a+3 & a+4 & a+5 & a+6 \\ a+3 & a+4 & a+5 & a+6 & a+7 \\ a+4 & a+5 & a+6 & a+7 & a+8 \end{bmatrix} \in M_5(\mathbb{R}).$$

244 Problem Let A, B be arbitrary $n \times n$ matrices over \mathbb{R} . Prove or disprove! $\text{rank } (AB) = \text{rank } (BA)$.

when $(a, b) \in \mathbb{R}^2$.

246 Problem Find the rank of $\begin{bmatrix} 1 & -1 & 0 & 1 \\ m & 1 & -1 & -1 \\ 1 & -m & 1 & 0 \\ 1 & -1 & m & 2 \end{bmatrix}$ as a function of $m \in \mathbb{C}$.

245 Problem Study the various possibilities for the rank of

247 Problem Determine the rank of the matrix

$$\begin{bmatrix} a^2 & ab & ab & b^2 \\ ab & a^2 & b^2 & ab \\ ab & b^2 & a^2 & ab \\ b^2 & ab & ab & a^2 \end{bmatrix}.$$

248 Problem Determine the rank of the matrix

$$\begin{bmatrix} 1 & 1 & 1 & 1 \\ a & b & a & b \\ c & c & d & d \\ ac & bc & ad & bd \end{bmatrix}.$$

249 Problem Let $A \in M_{3 \times 2}(\mathbb{R})$, $B \in M_2(\mathbb{R})$, and $C \in$

$$M_{2 \times 3}(\mathbb{R}) \text{ be such that } ABC = \begin{bmatrix} 1 & 1 & 2 \\ -2 & x & 1 \\ 1 & -2 & 1 \end{bmatrix}. \text{ Find } x.$$

250 Problem Let A, B be matrices of the same size. Prove that $\text{rank}(A + B) \leq \text{rank}(A) + \text{rank}(B)$.

251 Problem Let B be the matrix obtained by adjoining a row (or column) to a matrix A . Prove that either $\text{rank}(B) = \text{rank}(A)$ or $\text{rank}(B) = \text{rank}(A) + 1$.

252 Problem Let $A \in M_n(\mathbb{R})$. Prove that $\text{rank}(A) = \text{rank}(AA^T)$. Find a counterexample in the case $A \in M_n(\mathbb{C})$.

253 Problem Prove that the rank of a skew-symmetric matrix is an even number.

2.8 Rank and Invertibility

254 Theorem A matrix $A \in M_{m \times n}(\mathbb{F})$ is left-invertible if and only if $\text{rank}(A) = n$. A matrix $A \in M_{m \times n}(\mathbb{F})$ is right-invertible if and only if $\text{rank}(A) = m$.

Proof: Observe that we always have $\text{rank}(A) \leq n$. If A is left invertible, then $\exists L \in M_{n \times m}(\mathbb{F})$ such that $LA = I_n$. By Theorem 239,

$$n = \text{rank}(I_n) = \text{rank}(LA) \leq \text{rank}(A),$$

whence the two inequalities give $\text{rank}(A) = n$.

Conversely, assume that $\text{rank}(A) = n$. Then $\text{rank}(A^T) = n$ by Corollary 235, and so by Theorem 234 there exist $P \in GL_m(\mathbb{F})$, $Q \in GL_n(\mathbb{F})$, such that

$$PAQ = \begin{bmatrix} I_n \\ 0_{(m-n) \times n} \end{bmatrix}, \quad Q^T A^T P^T = \begin{bmatrix} I_n & 0_{n \times (m-n)} \end{bmatrix}.$$

This gives

$$\begin{aligned} Q^T A^T P^T PAQ &= I_n \implies A^T P^T PA = (Q^T)^{-1} Q^{-1} \\ &\implies ((Q^T)^{-1} Q^{-1})^{-1} A^T P^T PA = I_n, \end{aligned}$$

and so $((Q^T)^{-1} Q^{-1})^{-1} A^T P^T P$ is a left inverse for A .

The right-invertibility case is argued similarly. \square

By combining Theorem 254 and Theorem 195, the following corollary is thus immediate.

255 Corollary If $A \in M_{m \times n}(\mathbb{F})$ possesses a left inverse L and a right inverse R then $m = n$ and $L = R$.

We use Gauß-Jordan Reduction to find the inverse of $\mathbf{A} \in \text{GL}_n(\mathbb{F})$. We form the *augmented matrix* $\mathbf{T} = [\mathbf{A} | \mathbf{I}_n]$ which is obtained by putting \mathbf{A} side by side with the identity matrix \mathbf{I}_n . We perform permissible row operations on \mathbf{T} until instead of \mathbf{A} we obtain \mathbf{I}_n , which will appear if the matrix is invertible. The matrix on the right will be \mathbf{A}^{-1} . We finish with $[\mathbf{I}_n | \mathbf{A}^{-1}]$.

 If $\mathbf{A} \in M_n(\mathbb{R})$ is non-invertible, then the left hand side in the procedure above will not reduce to \mathbf{I}_n .

256 Example Find the inverse of the matrix $\mathbf{B} \in M_3(\mathbb{Z}_7)$,

$$\mathbf{B} = \begin{bmatrix} \bar{6} & \bar{0} & \bar{1} \\ \bar{3} & \bar{2} & \bar{0} \\ \bar{1} & \bar{0} & \bar{1} \end{bmatrix}.$$

Solution: We have

$$\begin{array}{c} \left[\begin{array}{ccc|ccc} \bar{6} & \bar{0} & \bar{1} & \bar{1} & \bar{0} & \bar{0} \\ \bar{3} & \bar{2} & \bar{0} & \bar{0} & \bar{1} & \bar{0} \\ \bar{1} & \bar{0} & \bar{1} & \bar{0} & \bar{0} & \bar{1} \end{array} \right] \xrightarrow[\sim]{\substack{\mathbf{R}_1 \leftrightarrow \mathbf{R}_3 \\ \mathbf{R}_2 - \bar{3}\mathbf{R}_1 \rightarrow \mathbf{R}_2}} \left[\begin{array}{ccc|ccc} \bar{1} & \bar{0} & \bar{1} & \bar{0} & \bar{0} & \bar{1} \\ \bar{3} & \bar{2} & \bar{0} & \bar{0} & \bar{1} & \bar{0} \\ \bar{6} & \bar{0} & \bar{1} & \bar{1} & \bar{0} & \bar{0} \end{array} \right] \\ \xrightarrow[\sim]{\substack{\mathbf{R}_3 - \bar{6}\mathbf{R}_1 \rightarrow \mathbf{R}_3 \\ \mathbf{R}_2 - \bar{3}\mathbf{R}_1 \rightarrow \mathbf{R}_2}} \left[\begin{array}{ccc|ccc} \bar{1} & \bar{0} & \bar{1} & \bar{0} & \bar{0} & \bar{1} \\ \bar{0} & \bar{2} & \bar{4} & \bar{0} & \bar{1} & \bar{4} \\ \bar{0} & \bar{0} & \bar{2} & \bar{1} & \bar{0} & \bar{1} \end{array} \right] \\ \xrightarrow[\sim]{\substack{\mathbf{R}_2 - \bar{2}\mathbf{R}_3 \rightarrow \mathbf{R}_2 \\ 5\mathbf{R}_1 + \mathbf{R}_3 \rightarrow \mathbf{R}_1}} \left[\begin{array}{ccc|ccc} \bar{5} & \bar{0} & \bar{0} & \bar{1} & \bar{0} & \bar{6} \\ \bar{0} & \bar{2} & \bar{0} & \bar{5} & \bar{1} & \bar{2} \\ \bar{0} & \bar{0} & \bar{2} & \bar{1} & \bar{0} & \bar{1} \end{array} \right] \\ \xrightarrow[\sim]{\substack{\bar{3}\mathbf{R}_1 \rightarrow \mathbf{R}_1; \bar{4}\mathbf{R}_3 \rightarrow \mathbf{R}_3 \\ \bar{4}\mathbf{R}_2 \rightarrow \mathbf{R}_2}} \left[\begin{array}{ccc|ccc} \bar{1} & \bar{0} & \bar{0} & \bar{3} & \bar{0} & \bar{4} \\ \bar{0} & \bar{1} & \bar{0} & \bar{6} & \bar{4} & \bar{1} \\ \bar{0} & \bar{0} & \bar{1} & \bar{4} & \bar{0} & \bar{4} \end{array} \right]. \end{array}$$

We conclude that

$$\begin{bmatrix} \bar{6} & \bar{0} & \bar{1} \\ \bar{3} & \bar{2} & \bar{0} \\ \bar{1} & \bar{0} & \bar{1} \end{bmatrix}^{-1} = \begin{bmatrix} \bar{3} & \bar{0} & \bar{4} \\ \bar{6} & \bar{4} & \bar{1} \\ \bar{4} & \bar{0} & \bar{4} \end{bmatrix}.$$

$$\begin{bmatrix} 0 & 1 & -1 \\ 4 & -3 & 4 \\ 3 & -3 & 4 \end{bmatrix}$$

257 Example Use Gauß-Jordan reduction to find the inverse of the matrix $\mathbf{A} = \begin{bmatrix} 0 & 1 & -1 \\ 4 & -3 & 4 \\ 3 & -3 & 4 \end{bmatrix}$. Also, find \mathbf{A}^{2001} .

Solution: Operating on the augmented matrix

$$\begin{array}{c}
 \left[\begin{array}{ccc|ccc} 0 & 1 & -1 & 1 & 0 & 0 \\ 4 & -3 & 4 & 0 & 1 & 0 \\ 3 & -3 & 4 & 0 & 0 & 1 \end{array} \right] \xrightarrow{\text{R}_2 - \text{R}_3 \rightarrow \text{R}_2} \left[\begin{array}{ccc|ccc} 0 & 1 & -1 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & -1 \\ 3 & -3 & 4 & 0 & 0 & 1 \end{array} \right] \\
 \qquad\qquad\qquad \xrightarrow{\text{R}_3 - 3\text{R}_2 \rightarrow \text{R}_3} \left[\begin{array}{ccc|ccc} 0 & 1 & -1 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & -1 \\ 0 & -3 & 4 & 0 & -3 & 4 \end{array} \right] \\
 \qquad\qquad\qquad \xrightarrow{\text{R}_3 + 3\text{R}_1 \rightarrow \text{R}_3} \left[\begin{array}{ccc|ccc} 0 & 1 & -1 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 1 & 3 & -3 & 4 \end{array} \right] \\
 \qquad\qquad\qquad \xrightarrow{\text{R}_1 + \text{R}_3 \rightarrow \text{R}_1} \left[\begin{array}{ccc|ccc} 0 & 1 & 0 & 4 & -3 & 4 \\ 1 & 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 1 & 3 & -3 & 4 \end{array} \right] \\
 \qquad\qquad\qquad \xrightarrow{\text{R}_1 \leftrightarrow \text{R}_2} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 0 & 1 & -1 \\ 0 & 1 & 0 & 4 & -3 & 4 \\ 0 & 0 & 1 & 3 & -3 & 4 \end{array} \right].
 \end{array}$$

Thus we deduce that

$$\mathbf{A}^{-1} = \begin{bmatrix} 0 & 1 & -1 \\ 4 & -3 & 4 \\ 3 & -3 & 4 \end{bmatrix} = \mathbf{A}.$$

From $\mathbf{A}^{-1} = \mathbf{A}$ we deduce $\mathbf{A}^2 = \mathbf{I}_n$. Hence $\mathbf{A}^{2000} = (\mathbf{A}^2)^{1000} = \mathbf{I}_n^{1000} = \mathbf{I}_n$ and $\mathbf{A}^{2001} = \mathbf{A}(\mathbf{A}^{2000}) = \mathbf{A}\mathbf{I}_n = \mathbf{A}$.

258 Example Find the inverse of the triangular matrix $\mathbf{A} \in M_n(\mathbb{R})$,

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 & \cdots & 1 \\ 0 & 1 & 1 & \cdots & 1 \\ 0 & 0 & 1 & \cdots & 1 \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \end{bmatrix}.$$

Solution: Form the augmented matrix

$$\left[\begin{array}{ccccc|ccccc} 1 & 1 & 1 & \cdots & 1 & 1 & 0 & 0 & \cdots & 0 \\ 0 & 1 & 1 & \cdots & 1 & 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 1 & 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \cdots & \vdots & \vdots & \vdots & \vdots & \cdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & 0 & 0 & 0 & \cdots & 1 \end{array} \right],$$

and perform $R_k - R_{k+1} \rightarrow R_k$ successively for $k = 1, 2, \dots, n-1$, obtaining

$$\left[\begin{array}{ccccc|ccccc} 1 & 0 & 0 & \cdots & 0 & 1 & -1 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & 0 & 0 & 1 & -1 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 & 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \cdots & \vdots & \vdots & \vdots & \vdots & \cdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & 0 & 0 & 0 & \cdots & 1 \end{array} \right],$$

whence

$$\mathbf{A}^{-1} = \begin{bmatrix} 1 & -1 & 0 & \cdots & 0 \\ 0 & 1 & -1 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \end{bmatrix},$$

that is, the inverse of \mathbf{A} has 1's on the diagonal and -1 's on the superdiagonal.

259 Theorem Let $\mathbf{A} \in M_n(\mathbb{F})$ be a triangular matrix such that $a_{11}a_{22}\cdots a_{nn} \neq 0_{\mathbb{F}}$. Then \mathbf{A} is invertible.

Proof: Since the entry $a_{kk} \neq 0_{\mathbb{F}}$ we multiply the k -th row by a_{kk}^{-1} and then proceed to subtract the appropriate multiples of the preceding $k-1$ rows at each stage. \square

260 Example (Putnam Exam, 1969) Let \mathbf{A} and \mathbf{B} be matrices of size 3×2 and 2×3 respectively. Suppose that their product \mathbf{AB} is given by

$$\mathbf{AB} = \begin{bmatrix} 8 & 2 & -2 \\ 2 & 5 & 4 \\ -2 & 4 & 5 \end{bmatrix}.$$

Demonstrate that the product \mathbf{BA} is given by

$$\mathbf{BA} = \begin{bmatrix} 9 & 0 \\ 0 & 9 \end{bmatrix}.$$

Solution: Observe that

$$(\mathbf{AB})^2 = \begin{bmatrix} 8 & 2 & -2 \\ 2 & 5 & 4 \\ -2 & 4 & 5 \end{bmatrix} \begin{bmatrix} 8 & 2 & -2 \\ 2 & 5 & 4 \\ -2 & 4 & 5 \end{bmatrix} = \begin{bmatrix} 72 & 18 & -18 \\ 18 & 45 & 36 \\ -18 & 36 & 45 \end{bmatrix} = 9\mathbf{AB}.$$

Performing $R_3 + R_2 \rightarrow R_3$, $R_1 - 4R_2 \rightarrow R_2$, and $2R_3 + R_1 \rightarrow R_3$ in succession we see that

$$\begin{bmatrix} 8 & 2 & -2 \\ 2 & 5 & 4 \\ -2 & 4 & 5 \end{bmatrix} \rightsquigarrow \begin{bmatrix} 0 & -18 & -18 \\ 2 & 5 & 4 \\ 0 & 0 & 0 \end{bmatrix} \rightsquigarrow \begin{bmatrix} 0 & -18 & 0 \\ 2 & 5 & -1 \\ 0 & 0 & 0 \end{bmatrix} \rightsquigarrow \begin{bmatrix} 0 & -18 & 0 \\ 0 & 5 & -1 \\ 0 & 0 & 0 \end{bmatrix},$$

and so $\text{rank } (\mathbf{AB}) = 2$. This entails that $\text{rank } ((\mathbf{AB})^2) = 2$. Now, since \mathbf{BA} is a 2×2 matrix, $\text{rank } (\mathbf{BA}) \leq 2$. Also

$$2 = \text{rank } ((\mathbf{AB})^2) = \text{rank } (\mathbf{ABA}\mathbf{B}) \leq \text{rank } (\mathbf{ABA}) \leq \text{rank } (\mathbf{BA}),$$

and we must conclude that $\text{rank } (\mathbf{BA}) = 2$. This means that \mathbf{BA} is invertible and so

$$\begin{aligned} (\mathbf{AB})^2 = 9\mathbf{AB} &\implies \mathbf{A}(\mathbf{BA} - 9\mathbf{I}_2)\mathbf{B} = \mathbf{0}_3 \\ &\implies \mathbf{BA}(\mathbf{BA} - 9\mathbf{I}_2)\mathbf{BA} = \mathbf{B}\mathbf{0}_3\mathbf{A} \\ &\implies \mathbf{BA}(\mathbf{BA} - 9\mathbf{I}_2)\mathbf{BA} = \mathbf{0}_2 \\ &\implies (\mathbf{BA})^{-1}\mathbf{BA}(\mathbf{BA} - 9\mathbf{I}_2)\mathbf{BA}(\mathbf{BA})^{-1} = (\mathbf{BA})^{-1}\mathbf{0}_2(\mathbf{BA})^{-1} \\ &\implies \mathbf{BA} - 9\mathbf{I}_2 = \mathbf{0}_2 \end{aligned}$$

261 Problem Find the inverse of the matrix

$$\begin{bmatrix} \bar{1} & \bar{2} & \bar{3} \\ \bar{2} & \bar{3} & \bar{1} \\ \bar{3} & \bar{1} & \bar{2} \end{bmatrix} \in M_3(\mathbb{Z}_7).$$

262 Problem Let $(A, B) \in M_3(\mathbb{R})$ be given by

$$A = \begin{bmatrix} a & b & c \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}, \quad B = \begin{bmatrix} 0 & 0 & -1 \\ 0 & -1 & a \\ -1 & a & b \end{bmatrix}.$$

Find B^{-1} and prove that $A^T = BAB^{-1}$.

263 Problem Find all the values of the parameter a for which the matrix B given below is not invertible.

$$B = \begin{bmatrix} -1 & a+2 & 2 \\ 0 & a & 1 \\ 2 & 1 & a \end{bmatrix}$$

264 Problem Find the inverse of the triangular matrix

$$\begin{bmatrix} a & 2a & 3a \\ 0 & b & 2b \\ 0 & 0 & c \end{bmatrix} \in M_3(\mathbb{R})$$

assuming that $abc \neq 0$.

265 Problem Under what conditions is the matrix

$$\begin{bmatrix} b & a & 0 \\ c & 0 & a \\ 0 & c & b \end{bmatrix}$$

invertible? Find the inverse under these conditions.

266 Problem Find the inverse of the matrix

$$\begin{bmatrix} 1+a & 1 & 1 \\ 1 & 1+b & 1 \\ 1 & 1 & 1+c \end{bmatrix}$$

267 Problem Prove that for the $n \times n$ ($n > 1$) matrix

$$\begin{bmatrix} 0 & 1 & 1 & \dots & 1 \\ 1 & 0 & 1 & \dots & 1 \\ 1 & 1 & 0 & \dots & 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & 1 & \dots & 0 \end{bmatrix}^{-1} = \frac{1}{n-1} \begin{bmatrix} 2-n & 1 & 1 & \dots & 1 \\ 1 & 2-n & 1 & \dots & 1 \\ 1 & 1 & 2-n & \dots & 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & 1 & \dots & 2-n \end{bmatrix}$$

268 Problem Prove that the $n \times n$ ($n > 1$) matrix

$$\begin{bmatrix} 1+a & 1 & 1 & \dots & 1 \\ 1 & 1+a & 1 & \dots & 1 \\ 1 & 1 & 1+a & \dots & 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & 1 & \dots & 1+a \end{bmatrix}$$

has inverse

$$-\frac{1}{a(n+a)} \begin{bmatrix} 1-n-a & 1 & 1 & \dots & 1 \\ 1 & 1-n-a & 1 & \dots & 1 \\ 1 & 1 & 1-n-a & \dots & 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & 1 & \dots & 1-n-a \end{bmatrix}$$

269 Problem Prove that

$$\begin{bmatrix} 1 & 3 & 5 & 7 & \dots & (2n-1) \\ (2n-1) & 1 & 3 & 5 & \dots & (2n-3) \\ (2n-3) & (2n-1) & 1 & 3 & \dots & (2n-5) \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 3 & 5 & 7 & 9 & \dots & 1 \end{bmatrix}$$

has inverse

$$\frac{1}{2n^3} \begin{bmatrix} 2-n^2 & 2+n^2 & 2 & 2 & \dots & 2 \\ 2 & 2-n^2 & 2+n^2 & 2 & \dots & 2 \\ 2 & 2 & 2-n^2 & 2+n^2 & \dots & 2 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 2+n^2 & 2 & 2 & 2 & \dots & 2-n^2 \end{bmatrix}.$$

270 Problem Prove that the $n \times n$ ($n > 1$) matrix

$$\begin{bmatrix} 1 + a_1 & 1 & 1 & \dots & 1 \\ 1 & 1 + a_2 & 1 & \dots & 1 \\ 1 & 1 & 1 + a_3 & \dots & 1 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 1 & 1 & 1 & \dots & 1 + a_n \end{bmatrix}$$

has inverse

$$-\frac{1}{s} \begin{bmatrix} \frac{1 - a_1 s}{a_1^2} & \frac{1}{a_1 a_2} & \frac{1}{a_1 a_3} & \dots & \frac{1}{a_1 a_n} \\ \frac{1}{a_2 a_1} & \frac{1 - a_2 s}{a_2^2} & \frac{1}{a_2 a_3} & \dots & \frac{1}{a_2 a_n} \\ \frac{1}{a_3 a_1} & \frac{1}{a_3 a_2} & \frac{1 - a_3 s}{a_3^2} & \dots & \frac{1}{a_3 a_1 n} \\ \vdots & \vdots & \vdots & \dots & \vdots \\ \frac{1}{a_n a_1} & \frac{1}{a_n a_2} & \frac{1}{a_n a_3} & \dots & \frac{1 - a_n s}{a_n^2} \end{bmatrix},$$

where $s = 1 + \frac{1}{a_1} + \frac{1}{a_2} + \dots + \frac{1}{a_n}$.

271 Problem Let $A \in M_5(\mathbb{R})$. Show that if $\text{rank}(A^2) < 5$, then $\text{rank}(A) < 5$.

272 Problem Let $A \in M_{3,2}(\mathbb{R})$ and $B \in M_{2,3}(\mathbb{R})$ be mat-

ces such that $AB = \begin{bmatrix} 0 & -1 & -1 \\ -1 & 0 & -1 \\ 1 & 1 & 2 \end{bmatrix}$. Prove that $BA = I_2$.

Chapter 3

Linear Equations

3.1 Definitions

We can write a system of m linear equations in n variables over a field \mathbb{F}

$$a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \cdots + a_{1n}x_n = y_1,$$

$$a_{21}x_1 + a_{22}x_2 + a_{23}x_3 + \cdots + a_{2n}x_n = y_2,$$

⋮

$$a_{m1}x_1 + a_{m2}x_2 + a_{m3}x_3 + \cdots + a_{mn}x_n = y_m,$$

in matrix form as

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \vdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_m \end{bmatrix}. \quad (3.1)$$

We write the above matrix relation in the abbreviated form

$$\mathbf{AX} = \mathbf{Y}, \quad (3.2)$$

where \mathbf{A} is the matrix of coefficients, \mathbf{X} is the matrix of variables and \mathbf{Y} is the matrix of constants. Most often we will dispense with the matrix of variables \mathbf{X} and will simply write the *augmented matrix* of the system as

$$[\mathbf{A}|\mathbf{Y}] = \left[\begin{array}{cccc|c} a_{11} & a_{12} & \cdots & a_{1n} & y_1 \\ a_{21} & a_{22} & \cdots & a_{2n} & y_2 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} & y_m \end{array} \right]. \quad (3.3)$$

273 Definition Let $\mathbf{AX} = \mathbf{Y}$ be as in 3.1. If $\mathbf{Y} = \mathbf{0}_{m \times 1}$, then the system is called *homogeneous*, otherwise it is called *inhomogeneous*. The set

$$\{\mathbf{X} \in M_{n \times 1}(\mathbb{F}) : \mathbf{AX} = \mathbf{0}_{m \times 1}\}$$

is called the *kernel* or *nullspace* of \mathbf{A} and it is denoted by $\ker(\mathbf{A})$.

 Observe that we always have $\mathbf{0}_{n \times 1} \in \ker(\mathbf{A}) \in M_{m \times n}(\mathbb{F})$.

274 Definition A system of linear equations is *consistent* if it has a solution. If the system does not have a solution then we say that it is *inconsistent*.

275 Definition If a row of a matrix is non-zero, we call the first non-zero entry of this row a *pivot* for this row.

276 Definition A matrix $\mathbf{M} \in M_{m \times n}(\mathbb{F})$ is a *row-echelon* matrix if

- All the zero rows of \mathbf{M} , if any, are at the bottom of \mathbf{M} .
- For any two consecutive rows R_i and R_{i+1} , either R_{i+1} is all $0_{\mathbb{F}}$'s or the pivot of R_{i+1} is immediately to the right of the pivot of R_i .

The variables accompanying these pivots are called the *leading variables*. Those variables which are not leading variables are the *free parameters*.

277 Example The matrices

$$\left[\begin{array}{cccc} 1 & 0 & 1 & 1 \\ 0 & 0 & 2 & 2 \\ 0 & 0 & 0 & 3 \\ 0 & 0 & 0 & 0 \end{array} \right], \quad \left[\begin{array}{cccc} 1 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right],$$

are in row-echelon form, with the pivots circled, but the matrices

$$\left[\begin{array}{cccc} 1 & 0 & 1 & 1 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{array} \right], \quad \left[\begin{array}{cccc} 1 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{array} \right],$$

are not in row-echelon form.

 Observe that given a matrix $\mathbf{A} \in M_{m \times n}(\mathbb{F})$, by following Gauß-Jordan reduction à la Theorem 284, we can find a matrix $\mathbf{P} \in GL_m(\mathbb{F})$ such that $\mathbf{PA} = \mathbf{B}$ is in row-echelon form.

278 Example Solve the system of linear equations

$$\left[\begin{array}{cccc} 1 & 1 & 1 & 1 \\ 0 & 2 & 1 & 0 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 2 \end{array} \right] \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} = \begin{bmatrix} -3 \\ -1 \\ 4 \\ -6 \end{bmatrix}.$$

Solution: Observe that the matrix of coefficients is already in row-echelon form. Clearly every variable is a leading variable, and by back substitution

$$2w = -6 \implies w = -\frac{6}{2} = -3,$$

$$z - w = 4 \implies z = 4 + w = 4 - 3 = 1,$$

$$2y + z = -1 \implies y = -\frac{1}{2} - \frac{1}{2}z = -1,$$

$$x + y + z + w = -3 \implies x = -3 - y - z - w = 0.$$

The (unique) solution is thus

$$\begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} = \begin{bmatrix} 0 \\ -1 \\ 1 \\ -3 \end{bmatrix}.$$

279 Example Solve the system of linear equations

$$\begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 2 & 1 & 0 \\ 0 & 0 & 1 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} = \begin{bmatrix} -3 \\ -1 \\ 4 \end{bmatrix}.$$

Solution: The system is already in row-echelon form, and we see that x, y, z are leading variables while w is a free parameter. We put $w = t$. Using back substitution, and operating from the bottom up, we find

$$z - w = 4 \implies z = 4 + w = 4 + t,$$

$$2y + z = -1 \implies y = -\frac{1}{2} - \frac{1}{2}z = -\frac{1}{2} - 2 - \frac{1}{2}t = -\frac{5}{2} - \frac{1}{2}t,$$

$$x + y + z + w = -3 \implies x = -3 - y - z - w = -3 + \frac{5}{2} + \frac{1}{2}t - 4 - t - t = -\frac{9}{2} - \frac{3}{2}t.$$

The solution is thus

$$\begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} = \begin{bmatrix} -\frac{9}{2} - \frac{3}{2}t \\ -\frac{5}{2} - \frac{1}{2}t \\ 4 + t \\ t \end{bmatrix}, \quad t \in \mathbb{R}.$$

280 Example Solve the system of linear equations

$$\begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 2 & 1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} = \begin{bmatrix} -3 \\ -1 \end{bmatrix}.$$

Solution: We see that x, y are leading variables, while z, w are free parameters. We put $z = s, w = t$. Operating from the bottom up, we find

$$2y + z = -1 \implies y = -\frac{1}{2} - \frac{1}{2}z = -\frac{1}{2} - \frac{1}{2}s,$$

$$x + y + z + w = -3 \implies x = -3 - y - z - w = -\frac{5}{2} - \frac{3}{2}s - t.$$

The solution is thus

$$\begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} = \begin{bmatrix} -\frac{5}{2} - \frac{3}{2}s - t \\ -\frac{1}{2} - \frac{1}{2}s \\ s \\ t \end{bmatrix}, \quad (s, t) \in \mathbb{R}^2.$$

281 Example Find all the solutions of the system

$$x + \bar{2}y + \bar{2}z = \bar{0},$$

$$y + \bar{2}z = \bar{1},$$

working in \mathbb{Z}_3 .

Solution: The augmented matrix of the system is

$$\left[\begin{array}{ccc|c} \bar{1} & \bar{2} & \bar{2} & \bar{0} \\ \bar{0} & \bar{1} & \bar{2} & \bar{1} \end{array} \right].$$

The system is already in row-echelon form and x, y are leading variables while z is a free parameter. We find

$$y = \bar{1} - \bar{2}z = \bar{1} + \bar{1}z,$$

and

$$x = -\bar{2}y - \bar{2}z = \bar{1} + \bar{2}z.$$

Thus

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} \bar{1} + \bar{2}z \\ \bar{1} + \bar{1}z \\ z \end{bmatrix}, \quad z \in \mathbb{Z}_3.$$

Letting $z = \bar{0}, \bar{1}, \bar{2}$ successively, we find the three solutions

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} \bar{1} \\ \bar{1} \\ \bar{0} \end{bmatrix},$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} \bar{0} \\ \bar{2} \\ \bar{1} \end{bmatrix},$$

and

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} \bar{2} \\ \bar{0} \\ \bar{2} \end{bmatrix}.$$

282 Problem Find all the solutions in \mathbb{Z}_3 of the system

$$x + y + z + w = \bar{0},$$

$$\bar{2}y + w = \bar{2}.$$

find all solutions of the system

$$\bar{1}x + \bar{2}y + \bar{3}z = \bar{5};$$

$$\bar{2}x + \bar{3}y + \bar{1}z = \bar{6};$$

$$\bar{3}x + \bar{1}y + \bar{2}z = \bar{0}.$$

283 Problem In \mathbb{Z}_7 , given that

$$\begin{bmatrix} \bar{1} & \bar{2} & \bar{3} \\ \bar{2} & \bar{3} & \bar{1} \\ \bar{3} & \bar{1} & \bar{2} \end{bmatrix}^{-1} = \begin{bmatrix} \bar{4} & \bar{2} & \bar{0} \\ \bar{2} & \bar{0} & \bar{4} \\ \bar{0} & \bar{4} & \bar{2} \end{bmatrix},$$

3.2 Existence of Solutions

We now answer the question of deciding when a system of linear equations is solvable.

284 Lemma Let $A \in M_{m \times n}(\mathbb{F})$ be in row-echelon form, and let $X \in M_{n \times 1}(\mathbb{F})$ be a matrix of variables. The homogeneous system $AX = 0_{m \times 1}$ of m linear equations in n variables has (i) a unique solution if $m = n$, (ii) multiple solutions if $m < n$.

Proof: If $m = n$ then A is a square triangular matrix whose diagonal elements are different from $0_{\mathbb{F}}$. As such, it is invertible by virtue of Theorem 259. Thus

$$AX = 0_{n \times 1} \implies X = A^{-1}0_{n \times 1} = 0_{n \times 1}$$

so there is only the unique solution $X = 0_{n \times 1}$, called the trivial solution.

If $m < n$ then there are $n - m$ free variables. Letting these variables run through the elements of the field, we obtain multiple solutions. Thus if the field has infinitely many elements, we obtain infinitely many solutions, and if the field has k elements, we obtain k^{n-m} solutions. Observe that in this case there is always a non-trivial solution.

□

285 Theorem Let $\mathbf{A} \in M_{m \times n}(\mathbb{F})$, and let $\mathbf{X} \in M_{n \times 1}(\mathbb{F})$ be a matrix of variables. The homogeneous system $\mathbf{AX} = \mathbf{0}_{m \times 1}$ of m linear equations in n variables always has a non-trivial solution if $m < n$.

Proof: We can find a matrix $\mathbf{P} \in GL_m(\mathbb{F})$ such that $\mathbf{B} = \mathbf{PA}$ is in row-echelon form. Now

$$\mathbf{AX} = \mathbf{0}_{m \times 1} \iff \mathbf{PAX} = \mathbf{0}_{m \times 1} \iff \mathbf{BX} = \mathbf{0}_{m \times 1}.$$

That is, the systems $\mathbf{AX} = \mathbf{0}_{m \times 1}$ and $\mathbf{BX} = \mathbf{0}_{m \times 1}$ have the same set of solutions. But by Lemma 284 there is a non-trivial solution. \square

286 Theorem (Kronecker-Capelli) Let $\mathbf{A} \in M_{m \times n}(\mathbb{F})$, $\mathbf{Y} \in M_{m \times 1}(\mathbb{F})$ be constant matrices and $\mathbf{X} \in M_{n \times 1}(\mathbb{F})$ be a matrix of variables. The matrix equation $\mathbf{AX} = \mathbf{Y}$ is solvable if and only if

$$\text{rank}(\mathbf{A}) = \text{rank}([\mathbf{A}|\mathbf{Y}]).$$

Proof: Assume first that $\mathbf{AX} = \mathbf{Y}$,

$$\mathbf{X} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}.$$

Let the columns of $[\mathbf{A}|\mathbf{X}]$ be denoted by \mathbf{C}_i , $1 \leq i \leq n$. Observe that that $[\mathbf{A}|\mathbf{X}] \in M_{m \times (n+1)}(\mathbb{F})$ and that the $(n+1)$ -th column of $[\mathbf{A}|\mathbf{X}]$ is

$$\mathbf{C}_{n+1} = \mathbf{AX} = \begin{bmatrix} x_1 a_{11} + x_2 a_{12} + \cdots + x_n a_{1n} \\ x_1 a_{21} + x_2 a_{22} + \cdots + x_n a_{2n} \\ \vdots \\ x_1 a_{n1} + x_2 a_{n2} + \cdots + x_n a_{nn} \end{bmatrix} = \sum_{i=1}^n x_i \mathbf{C}_i.$$

By performing $\mathbf{C}_{n+1} - \sum_{j=1}^n x_j \mathbf{C}_j \rightarrow \mathbf{C}_{n+1}$ on $[\mathbf{A}|\mathbf{Y}] = [\mathbf{A}|\mathbf{AX}]$ we obtain $[\mathbf{A}|\mathbf{0}_{n \times 1}]$. Thus $\text{rank}([\mathbf{A}|\mathbf{Y}]) = \text{rank}([\mathbf{A}|\mathbf{0}_{n \times 1}]) = \text{rank}(\mathbf{A})$.

Now assume that $r = \text{rank}(\mathbf{A}) = \text{rank}([\mathbf{A}|\mathbf{Y}])$. This means that adding an extra column to \mathbf{A} does not change the rank, and hence, by a sequence column operations $[\mathbf{A}|\mathbf{Y}]$ is equivalent to $[\mathbf{A}|\mathbf{0}_{n \times 1}]$. Observe that none of these operations is a permutation of the columns, since the first n columns of $[\mathbf{A}|\mathbf{Y}]$ and $[\mathbf{A}|\mathbf{0}_{n \times 1}]$ are the same. This means that \mathbf{Y} can be obtained from the columns \mathbf{C}_i , $1 \leq i \leq n$ of \mathbf{A} by means of transvections and dilatations. But then

$$\mathbf{Y} = \sum_{i=1}^n x_i \mathbf{C}_i.$$

The solutions sought is thus

$$\mathbf{X} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}.$$

\square

287 Problem Let $A \in M_{n \times p}(\mathbb{F})$, $B \in M_{n \times q}(\mathbb{F})$ and put $C = [A \quad B] \in M_{n \times (p+q)}(\mathbb{F})$. Prove that $\text{rank}(A) = \left| \begin{array}{l} \text{rank}(C) \iff \exists P \in M_p(q) \text{ such that } B = AP. \\ \text{rank}(B) \end{array} \right|$

3.3 Examples of Linear Systems

288 Example Use row reduction to solve the system

$$\begin{aligned} x + 2y + 3z + 4w &= 8 \\ x + 2y + 4z + 7w &= 12 \\ 2x + 4y + 6z + 8w &= 16 \end{aligned}$$

Solutions: Form the expanded matrix of coefficients and apply row operations to obtain

$$\left[\begin{array}{cccc|c} 1 & 2 & 3 & 4 & 8 \\ 1 & 2 & 4 & 7 & 12 \\ 2 & 4 & 6 & 8 & 16 \end{array} \right] \xrightarrow{\substack{R_3 - 2R_1 \rightarrow R_3 \\ R_2 - R_1 \rightarrow R_2}} \left[\begin{array}{cccc|c} 1 & 2 & 3 & 4 & 8 \\ 0 & 0 & 1 & 3 & 4 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right].$$

The matrix is now in row-echelon form. The variables x and z are the pivots, so w and y are free. Setting $w = s, y = t$ we have

$$z = 4 - 3s,$$

$$x = 8 - 4w - 3z - 2y = 8 - 4s - 3(4 - 3s) - 2t = -4 + 5s - 2t.$$

Hence the solution is given by

$$\begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} = \begin{bmatrix} -4 + 5s - 2t \\ t \\ 4 - 3s \\ s \end{bmatrix}.$$

289 Example Find $\alpha \in \mathbb{R}$ such that the system

$$x + y - z = 1,$$

$$2x + 3y + \alpha z = 3,$$

$$x + \alpha y + 3z = 2,$$

posses (i) no solution, (ii) infinitely many solutions, (iii) a unique solution.

Solution: The augmented matrix of the system is

$$\left[\begin{array}{ccc|c} 1 & 1 & -1 & 1 \\ 2 & 3 & \alpha & 3 \\ 1 & \alpha & 3 & 2 \end{array} \right].$$

By performing $R_2 - 2R_1 \rightarrow R_2$ and $R_3 - R_1 \rightarrow R_3$ we obtain

$$\rightsquigarrow \left[\begin{array}{ccc|c} 1 & 1 & -1 & 1 \\ 0 & 1 & \alpha + 2 & 1 \\ 0 & \alpha - 1 & 4 & 1 \end{array} \right].$$

By performing $R_3 - (\alpha - 1)R_2 \rightarrow R_3$ on this last matrix we obtain

$$\rightsquigarrow \left[\begin{array}{ccc|c} 1 & 1 & -1 & 1 \\ 0 & 1 & \alpha + 2 & 1 \\ 0 & 0 & (\alpha - 2)(\alpha + 3) & \alpha - 2 \end{array} \right].$$

If $\alpha = -3$, we obtain no solution. If $\alpha = 2$, there is an infinity of solutions

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 5t \\ 1 - 4t \\ t \end{bmatrix}, \quad t \in \mathbb{R}.$$

If $\alpha \neq 2$ and $\alpha \neq -3$, there is a unique solution

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ \frac{1}{\alpha + 3} \\ \frac{1}{\alpha + 3} \end{bmatrix}.$$

290 Example Solve the system

$$\begin{bmatrix} \bar{6} & \bar{0} & \bar{1} \\ \bar{3} & \bar{2} & \bar{0} \\ \bar{1} & \bar{0} & \bar{1} \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} \bar{1} \\ \bar{0} \\ \bar{2} \end{bmatrix},$$

for $(x, y, z) \in (\mathbb{Z}_7)^3$.

Solution: Performing operations on the augmented matrix we have

$$\begin{array}{c} \begin{bmatrix} \bar{6} & \bar{0} & \bar{1} & \bar{1} \\ \bar{3} & \bar{2} & \bar{0} & \bar{0} \\ \bar{1} & \bar{0} & \bar{1} & \bar{2} \end{bmatrix} \xrightarrow{\substack{R_1 \leftrightarrow R_3 \\ \sim}} \begin{bmatrix} \bar{1} & \bar{0} & \bar{1} & \bar{2} \\ \bar{3} & \bar{2} & \bar{0} & \bar{0} \\ \bar{6} & \bar{0} & \bar{1} & \bar{1} \end{bmatrix} \\ \xrightarrow{\substack{R_3 - \bar{6}R_1 \rightarrow R_3 \\ R_2 - \bar{3}R_1 \rightarrow R_2}} \begin{bmatrix} \bar{1} & \bar{0} & \bar{1} & \bar{2} \\ \bar{0} & \bar{2} & \bar{4} & \bar{1} \\ \bar{0} & \bar{0} & \bar{2} & \bar{3} \end{bmatrix} \end{array}$$

This gives

$$\begin{aligned}\bar{2}z &= \bar{3} \implies z = \bar{5}, \\ \bar{2}y - \bar{4}z &= \bar{2} \implies y = \bar{1}, \\ x &= \bar{2} - z = \bar{4}.\end{aligned}$$

The solution is thus

$$(x, y, z) = (\bar{4}, \bar{1}, \bar{5}).$$

291 Problem Find the general solution to the system

$$\left[\begin{array}{ccccc} 1 & 1 & 1 & 1 & 1 \\ 1 & 0 & 1 & 0 & 1 \\ 2 & 1 & 2 & 1 & 2 \\ 4 & 2 & 4 & 2 & 4 \\ 1 & 0 & 0 & 0 & 1 \end{array} \right] \left[\begin{array}{c} a \\ b \\ c \\ d \\ f \end{array} \right] = \left[\begin{array}{c} 1 \\ -1 \\ 0 \\ 0 \\ 0 \end{array} \right]$$

or shew that there is no solution.

292 Problem Find all solutions of the system

$$\left[\begin{array}{ccccc} 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 2 \\ 1 & 1 & 1 & 3 & 3 \\ 1 & 1 & 4 & 4 & 4 \\ 1 & 2 & 3 & 4 & 5 \end{array} \right] \left[\begin{array}{c} a \\ b \\ c \\ d \\ f \end{array} \right] = \left[\begin{array}{c} 3 \\ 4 \\ 7 \\ 6 \\ 9 \end{array} \right],$$

if any.

293 Problem Study the system

$$x + 2my + z = 4m;$$

$$2mx + y + z = 2;$$

$$x + y + 2mz = 2m^2,$$

with real parameter m . You must determine, with proof, for which m this system has (i) no solution, (ii) exactly one solution, and (iii) infinitely many solutions.

294 Problem Study the following system of linear equations with parameter a .

$$(2a - 1)x + ay - (a + 1)z = 1,$$

$$ax + y - 2z = 1,$$

$$2x + (3 - a)y + (2a - 6)z = 1.$$

You must determine for which a there is: (i) no solution, (ii) a unique solution, (iii) infinitely many solutions.

295 Problem Determine the values of the parameter m for which the system

$$\begin{aligned}x + y + (1 - m)z &= m + 2 \\ (1 + m)x - y + 2z &= 0 \\ 2x - my + 3z &= m + 2\end{aligned}$$

is solvable.

296 Problem Determine the values of the parameter m for which the system

$$\begin{aligned}x + y + z + t &= 4a \\ x - y - z + t &= 4b \\ -x - y + z + t &= 4c \\ x - y + z - t &= 4d\end{aligned}$$

is solvable.

297 Problem It is known that the system

$$ay + bx = c;$$

$$cx + az = b;$$

$$bz + cy = a$$

possesses a unique solution. What conditions must $(a, b, c) \in \mathbb{R}^3$ fulfill in this case? Find this unique solution.

298 Problem Find strictly positive real numbers x, y, z such that

$$\begin{aligned}x^3y^2z^6 &= 1 \\ x^4y^5z^{12} &= 2 \\ x^2y^2z^5 &= 3.\end{aligned}$$

299 Problem (Leningrad Mathematical Olympiad, 1987, Grade 5)

The numbers 1, 2, ..., 16 are arranged in a 4×4 matrix A as shewn below. We may add 1 to all the numbers of any row or subtract 1 from all numbers of any column. Using only the allowed operations, how can we obtain A^T ?

$$\mathbf{A} = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \\ 13 & 14 & 15 & 16 \end{bmatrix}$$

300 Problem (International Mathematics Olympiad, 1963)

Find all solutions x_1, x_2, x_3, x_4, x_5 of the system

$$x_5 + x_2 = yx_1;$$

$$x_1 + x_3 = yx_2;$$

$$x_2 + x_4 = yx_3;$$

$$x_3 + x_5 = yx_4;$$

$$x_4 + x_1 = yx_5,$$

where y is a parameter.

Chapter 4

\mathbb{R}^2 , \mathbb{R}^3 and \mathbb{R}^n

4.1 Points and Bi-points in \mathbb{R}^2

\mathbb{R}^2 is the set of all points $\mathbf{A} = \begin{pmatrix} a_1 \\ a_2 \end{pmatrix}$ with real number coordinates on the plane, as in figure 4.1. We use the notation $\mathbf{O} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$ to denote the *origin*.




Figure 4.1: Rectangular coordinates in \mathbb{R}^2 .

Given $\mathbf{A} = \begin{pmatrix} a_1 \\ a_2 \end{pmatrix} \in \mathbb{R}^2$ and $\mathbf{B} = \begin{pmatrix} b_1 \\ b_2 \end{pmatrix} \in \mathbb{R}^2$ we define their addition as

$$\mathbf{A} + \mathbf{B} = \begin{pmatrix} a_1 \\ a_2 \end{pmatrix} + \begin{pmatrix} b_1 \\ b_2 \end{pmatrix} = \begin{pmatrix} a_1 + b_1 \\ a_2 + b_2 \end{pmatrix}. \quad (4.1)$$

Similarly, we define the scalar multiplication of a point of \mathbb{R}^2 by the scalar $\alpha \in \mathbb{R}$ as

$$\alpha \mathbf{A} = \alpha \begin{pmatrix} a_1 \\ a_2 \end{pmatrix} = \begin{pmatrix} \alpha a_1 \\ \alpha a_2 \end{pmatrix}. \quad (4.2)$$

 Throughout this chapter, unless otherwise noted, we will use the convention that a point $\mathbf{A} \in \mathbb{R}^2$ will have its coordinates named after its letter, thus

$$\mathbf{A} = \begin{pmatrix} a_1 \\ a_2 \end{pmatrix}.$$

301 Definition Consider the points $\mathbf{A} \in \mathbb{R}^2, \mathbf{B} \in \mathbb{R}^2$. By the *bi-point* starting at \mathbf{A} and ending at \mathbf{B} , denoted by $[\mathbf{A}, \mathbf{B}]$, we mean the directed line segment from \mathbf{A} to \mathbf{B} . We define

$$[\mathbf{A}, \mathbf{A}] = \mathbf{O} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

 The bi-point $[\mathbf{A}, \mathbf{B}]$ can be thus interpreted as an arrow starting at \mathbf{A} and finishing, with the arrow tip, at \mathbf{B} . We say that \mathbf{A} is the tail of the bi-point $[\mathbf{A}, \mathbf{B}]$ and that \mathbf{B} is its head. Some authors use the terminology “fixed vector” instead of “bi-point.”

302 Definition Let $\mathbf{A} \neq \mathbf{B}$ be points on the plane and let \mathbf{L} be the line passing through \mathbf{A} and \mathbf{B} . The *direction* of the bi-point $[\mathbf{A}, \mathbf{B}]$ is the direction of the line \mathbf{L} , that is, the angle $\theta \in]-\frac{\pi}{2}, \frac{\pi}{2}]$ that the line \mathbf{L} makes with the horizontal. See figure 4.2.

303 Definition Let \mathbf{A}, \mathbf{B} lie on line \mathbf{L} , and let \mathbf{C}, \mathbf{D} lie on line \mathbf{L}' . If $\mathbf{L} \parallel \mathbf{L}'$ then we say that $[\mathbf{A}, \mathbf{B}]$ has the same direction as $[\mathbf{C}, \mathbf{D}]$. We say that the bi-points $[\mathbf{A}, \mathbf{B}]$ and $[\mathbf{C}, \mathbf{D}]$ have the *same sense* if they have the same direction and if both their heads lie on the same half-plane made by the line joining their tails. They have *opposite sense* if they have the same direction and if both their heads lie on alternative half-planes made by the line joining their tails. See figures 4.3 and 4.4.




Figure 4.2: Direction of a bi-point




Figure 4.3: Bi-points with the same sense.




Figure 4.4: Bi-points with opposite sense.

 Bi-point $[\mathbf{B}, \mathbf{A}]$ has the opposite sense of $[\mathbf{A}, \mathbf{B}]$ and so we write

$$[\mathbf{B}, \mathbf{A}] = -[\mathbf{A}, \mathbf{B}].$$

304 Definition Let $\mathbf{A} \neq \mathbf{B}$. The *Euclidean length or norm* of bi-point $[\mathbf{A}, \mathbf{B}]$ is simply the distance between \mathbf{A} and \mathbf{B} and it is denoted by

$$\|[\mathbf{A}, \mathbf{B}]\| = \sqrt{(\mathbf{a}_1 - \mathbf{b}_1)^2 + (\mathbf{a}_2 - \mathbf{b}_2)^2}.$$

We define

$$\|[\mathbf{A}, \mathbf{A}]\| = \|\mathbf{O}\| = 0.$$

A bi-point is said to have *unit length* if it has norm 1.

 A bi-point is completely determined by three things: (i) its norm, (ii) its direction, and (iii) its sense.

305 Definition (Chasles' Rule) Two bi-points are said to be *contiguous* if one has as tail the head of the other. In such case we define the sum of contiguous bi-points $[\mathbf{A}, \mathbf{B}]$ and $[\mathbf{B}, \mathbf{C}]$ by *Chasles' Rule*

$$[\mathbf{A}, \mathbf{B}] + [\mathbf{B}, \mathbf{C}] = [\mathbf{A}, \mathbf{C}].$$

See figure 4.5.

306 Definition (Scalar Multiplication of Bi-points) Let $\lambda \in \mathbb{R} \setminus \{0\}$ and $\mathbf{A} \neq \mathbf{B}$. We define

$$0[\mathbf{A}, \mathbf{B}] = \mathbf{O}$$

and

$$\lambda[\mathbf{A}, \mathbf{A}] = \mathbf{O}.$$

We define $\lambda[\mathbf{A}, \mathbf{B}]$ as follows.

1. $\lambda[\mathbf{A}, \mathbf{B}]$ has the direction of $[\mathbf{A}, \mathbf{B}]$.
2. $\lambda[\mathbf{A}, \mathbf{B}]$ has the sense of $[\mathbf{A}, \mathbf{B}]$ if $\lambda > 0$ and sense opposite $[\mathbf{A}, \mathbf{B}]$ if $\lambda < 0$.
3. $\lambda[\mathbf{A}, \mathbf{B}]$ has norm $|\lambda| \|[\mathbf{A}, \mathbf{B}]\|$ which is a contraction of $[\mathbf{A}, \mathbf{B}]$ if $0 < |\lambda| < 1$ or a dilatation of $[\mathbf{A}, \mathbf{B}]$ if $|\lambda| > 1$.

See figure 4.6 for some examples.




Figure 4.5: Chasles' Rule.




Figure 4.6: Scalar multiplication of bi-points.

4.2 Vectors in \mathbb{R}^2

307 Definition (Midpoint) Let \mathbf{A}, \mathbf{B} be points in \mathbb{R}^2 . We define the *midpoint* of the bi-point $[\mathbf{A}, \mathbf{B}]$ as

$$\frac{\mathbf{A} + \mathbf{B}}{2} = \begin{pmatrix} \frac{\mathbf{a}_1 + \mathbf{b}_1}{2} \\ \frac{\mathbf{a}_2 + \mathbf{b}_2}{2} \end{pmatrix}.$$

308 Definition (Equipollence) Two bi-points $[\mathbf{X}, \mathbf{Y}]$ and $[\mathbf{A}, \mathbf{B}]$ are said to be *equipollent* written $[\mathbf{X}, \mathbf{Y}] \sim [\mathbf{A}, \mathbf{B}]$ if the midpoints of the bi-points $[\mathbf{X}, \mathbf{B}]$ and $[\mathbf{Y}, \mathbf{A}]$ coincide, that is,

$$[\mathbf{X}, \mathbf{Y}] \sim [\mathbf{A}, \mathbf{B}] \Leftrightarrow \frac{\mathbf{X} + \mathbf{B}}{2} = \frac{\mathbf{Y} + \mathbf{A}}{2}.$$

See figure 4.7.

Geometrically, equipollence means that the quadrilateral \mathbf{XYBA} is a parallelogram. Thus the bi-points $[\mathbf{X}, \mathbf{Y}]$ and $[\mathbf{A}, \mathbf{B}]$ have the same norm, sense, and direction.




Figure 4.7: Equipollent bi-points.

309 Lemma Two bi-points $[\mathbf{X}, \mathbf{Y}]$ and $[\mathbf{A}, \mathbf{B}]$ are equipollent if and only if

$$\begin{pmatrix} \mathbf{y}_1 - \mathbf{x}_1 \\ \mathbf{y}_2 - \mathbf{x}_2 \end{pmatrix} = \begin{pmatrix} \mathbf{b}_1 - \mathbf{a}_1 \\ \mathbf{b}_2 - \mathbf{a}_2 \end{pmatrix}.$$

Proof: This is immediate, since

$$\begin{aligned} [\mathbf{X}, \mathbf{Y}] \sim [\mathbf{A}, \mathbf{B}] &\iff \begin{pmatrix} \frac{\mathbf{a}_1 + \mathbf{y}_1}{2} \\ \frac{\mathbf{a}_2 + \mathbf{y}_2}{2} \end{pmatrix} = \begin{pmatrix} \frac{\mathbf{b}_1 + \mathbf{x}_1}{2} \\ \frac{\mathbf{b}_2 + \mathbf{x}_2}{2} \end{pmatrix} \\ &\iff \begin{pmatrix} \mathbf{y}_1 - \mathbf{x}_1 \\ \mathbf{y}_2 - \mathbf{x}_2 \end{pmatrix} = \begin{pmatrix} \mathbf{b}_1 - \mathbf{a}_1 \\ \mathbf{b}_2 - \mathbf{a}_2 \end{pmatrix}, \end{aligned}$$

as desired. \square

From Lemma 309, equipollent bi-points have the same norm, the same direction, and the same sense.

310 Theorem Equipollence is an equivalence relation.

Proof: Write $[X, Y] \sim [A, B]$ if $[X, Y]$ is equipollent to $[A, B]$. Now $[X, Y] \sim [X, Y]$ since $\begin{pmatrix} y_1 - x_1 \\ y_2 - x_2 \end{pmatrix} =$

$\begin{pmatrix} y_1 - x_1 \\ y_2 - x_2 \end{pmatrix}$ and so the relation is reflexive. Also

$$\begin{aligned} [X, Y] \sim [A, B] &\iff \begin{pmatrix} y_1 - x_1 \\ y_2 - x_2 \end{pmatrix} = \begin{pmatrix} b_1 - a_1 \\ b_2 - a_2 \end{pmatrix} \\ &\iff \begin{pmatrix} b_1 - a_1 \\ b_2 - a_2 \end{pmatrix} = \begin{pmatrix} y_1 - x_1 \\ y_2 - x_2 \end{pmatrix} \\ &\iff [A, B] \sim [X, Y], \end{aligned}$$

and the relation is symmetric. Finally

$$\begin{aligned} [X, Y] \sim [A, B] \wedge [A, B] \sim [U, V] &\iff \begin{pmatrix} y_1 - x_1 \\ y_2 - x_2 \end{pmatrix} = \begin{pmatrix} b_1 - a_1 \\ b_2 - a_2 \end{pmatrix} \\ &\quad \wedge \begin{pmatrix} b_1 - a_1 \\ b_2 - a_2 \end{pmatrix} = \begin{pmatrix} v_1 - u_1 \\ v_2 - u_2 \end{pmatrix} \\ &\iff \begin{pmatrix} y_1 - x_1 \\ y_2 - x_2 \end{pmatrix} = \begin{pmatrix} v_1 - u_1 \\ v_2 - u_2 \end{pmatrix} \\ &\iff [X, Y] \sim [U, V], \end{aligned}$$

and the relation is transitive. \square

311 Definition (Vectors on the Plane) The equivalence class in which the bi-point $[X, Y]$ falls is called the *vector* (or *free vector*) from X to Y , and is denoted by \overrightarrow{XY} . Thus we write

$$[X, Y] \in \overrightarrow{XY} = \begin{bmatrix} y_1 - x_1 \\ y_2 - x_2 \end{bmatrix}.$$

If we desire to talk about a vector without mentioning a bi-point representative, we write, say, v , thus denoting vectors with boldface lowercase letters. If it is necessary to mention the coordinates of v we will write

$$v = \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}.$$

 For any point X on the plane, we have $\overrightarrow{XX} = \mathbf{0}$, the zero vector. If $[X, Y] \in v$ then $[Y, X] \in -v$.

312 Definition (Position Vector) For any particular point $P = \begin{pmatrix} p_1 \\ p_2 \end{pmatrix} \in \mathbb{R}^2$ we may form the vector $\overrightarrow{OP} = \begin{pmatrix} p_1 \\ p_2 \end{pmatrix}$.

We call \overrightarrow{OP} the *position vector* of P and we use boldface lowercase letters to denote the equality $\overrightarrow{OP} = p$.

313 Example The vector into which the bi-point with tail at $A = \begin{pmatrix} -1 \\ 2 \end{pmatrix}$ and head at $B = \begin{pmatrix} 3 \\ 4 \end{pmatrix}$ falls is

$$\overrightarrow{AB} = \begin{bmatrix} 3 - (-1) \\ 4 - 2 \end{bmatrix} = \begin{bmatrix} 4 \\ 2 \end{bmatrix}.$$

314 Example The bi-points $[A, B]$ and $[X, Y]$ with

$$A = \begin{pmatrix} -1 \\ 2 \end{pmatrix}, B = \begin{pmatrix} 3 \\ 4 \end{pmatrix},$$

$$X = \begin{pmatrix} 3 \\ 7 \end{pmatrix}, Y = \begin{pmatrix} 7 \\ 9 \end{pmatrix}$$

represent the same vector

$$\overrightarrow{AB} = \begin{bmatrix} 3 - (-1) \\ 4 - 2 \end{bmatrix} = \begin{bmatrix} 4 \\ 2 \end{bmatrix} = \begin{bmatrix} 7 - 3 \\ 9 - 7 \end{bmatrix} = \overrightarrow{XY}.$$

In fact, if $S = \begin{pmatrix} -1 + n \\ 2 + m \end{pmatrix}, T = \begin{pmatrix} 3 + n \\ 4 + m \end{pmatrix}$ then the infinite number of bi-points $[S, T]$ are representatives of the vectors $\overrightarrow{AB} = \overrightarrow{XY} = \overrightarrow{ST}$.

Given two vectors u, v we define their sum $u + v$ as follows. Find a bi-point representative $\overrightarrow{AB} \in u$ and a contiguous bi-point representative $\overrightarrow{BC} \in v$. Then by Chasles' Rule

$$u + v = \overrightarrow{AB} + \overrightarrow{BC} = \overrightarrow{AC}.$$

Again, by virtue of Chasles' Rule we then have

$$\overrightarrow{AB} = \overrightarrow{AO} + \overrightarrow{OB} = -\overrightarrow{OA} + \overrightarrow{OB} = b - a \quad (4.3)$$

Similarly we define scalar multiplication of a vector by scaling one of its bi-point representatives. We define the norm of a vector $v \in \mathbb{R}^2$ to be the norm of any of its bi-point representatives.

Componentwise we may see that given vectors $\mathbf{u} = \begin{bmatrix} \mathbf{u}_1 \\ \mathbf{u}_2 \end{bmatrix}$, $\mathbf{v} = \begin{bmatrix} \mathbf{v}_1 \\ \mathbf{v}_2 \end{bmatrix}$, and a scalar $\lambda \in \mathbb{R}$ then their sum and scalar multiplication take the form

$$\mathbf{u} + \mathbf{v} = \begin{bmatrix} \mathbf{u}_1 \\ \mathbf{u}_2 \end{bmatrix} + \begin{bmatrix} \mathbf{v}_1 \\ \mathbf{v}_2 \end{bmatrix}, \quad \lambda\mathbf{u} = \begin{bmatrix} \lambda\mathbf{u}_1 \\ \lambda\mathbf{u}_2 \end{bmatrix}.$$




Figure 4.8: Addition of Vectors.




Figure 4.9: Scalar multiplication of vectors.

315 Example Diagonals are drawn in a rectangle ABCD. If $\overrightarrow{AB} = \mathbf{x}$ and $\overrightarrow{AC} = \mathbf{y}$, then $\overrightarrow{BC} = \mathbf{y} - \mathbf{x}$, $\overrightarrow{CD} = -\mathbf{x}$, $\overrightarrow{DA} = \mathbf{x} - \mathbf{y}$, and $\overrightarrow{BD} = \mathbf{y} - 2\mathbf{x}$.

316 Definition (Parallel Vectors) Two vectors \mathbf{u} and \mathbf{v} are said to be *parallel* if there is a scalar λ such that $\mathbf{u} = \lambda\mathbf{v}$. If \mathbf{u} is parallel to \mathbf{v} we write $\mathbf{u} \parallel \mathbf{v}$. We denote by $\mathbb{R}\mathbf{v} = \{\alpha\mathbf{v} : \alpha \in \mathbb{R}\}$, the set of all vectors parallel to \mathbf{v} .

$\mathbf{0}$ is parallel to every vector.

317 Definition If $\mathbf{u} = \begin{bmatrix} \mathbf{u}_1 \\ \mathbf{u}_2 \end{bmatrix}$, then we define its *norm* as $\|\mathbf{u}\| = \sqrt{\mathbf{u}_1^2 + \mathbf{u}_2^2}$. The distance between two vectors \mathbf{u} and \mathbf{v} is $d(\mathbf{u}, \mathbf{v}) = \|\mathbf{u} - \mathbf{v}\|$.

318 Example Let $a \in \mathbb{R}$, $a > 0$ and let $\mathbf{v} \neq \mathbf{0}$. Find a vector with norm a and parallel to \mathbf{v} .

Solution: Observe that $\frac{\mathbf{v}}{\|\mathbf{v}\|}$ has norm 1 as

$$\left\| \frac{\mathbf{v}}{\|\mathbf{v}\|} \right\| = \frac{\|\mathbf{v}\|}{\|\mathbf{v}\|} = 1.$$

Hence the vector $a \frac{\mathbf{v}}{\|\mathbf{v}\|}$ has norm a and it is in the direction of \mathbf{v} . One may also take $-a \frac{\mathbf{v}}{\|\mathbf{v}\|}$.

319 Example If M is the midpoint of the bi-point [X, Y] then $\overrightarrow{XM} = \overrightarrow{MY}$ from where $\overrightarrow{XM} = \frac{1}{2}\overrightarrow{XY}$. Moreover, if T

is any point, by Chasles' Rule

$$\begin{aligned}\overrightarrow{TX} + \overrightarrow{TY} &= \overrightarrow{TM} + \overrightarrow{MX} + \overrightarrow{TM} + \overrightarrow{MY} \\ &= 2\overrightarrow{TM} - \overrightarrow{XM} + \overrightarrow{MY} \\ &= 2\overrightarrow{TM}.\end{aligned}$$

320 Example Let $\triangle ABC$ be a triangle on the plane. Prove that the line joining the midpoints of two sides of the triangle is parallel to the third side and measures half its length.

Solution: Let the midpoints of $[A, B]$ and $[A, C]$ be M_C and M_B , respectively. We shew that $\overrightarrow{BC} = 2\overrightarrow{M_C M_B}$. We have $2\overrightarrow{AM_C} = \overrightarrow{AB}$ and $2\overrightarrow{AM_B} = \overrightarrow{AC}$. Thus

$$\begin{aligned}\overrightarrow{BC} &= \overrightarrow{BA} + \overrightarrow{AC} \\ &= -\overrightarrow{AB} + \overrightarrow{AC} \\ &= -2\overrightarrow{AM_C} + 2\overrightarrow{AM_B} \\ &= 2\overrightarrow{M_C A} + 2\overrightarrow{AM_B} \\ &= 2(\overrightarrow{M_C A} + \overrightarrow{AM_B}) \\ &= 2\overrightarrow{M_C M_B},\end{aligned}$$

as we wanted to shew.

321 Example In $\triangle ABC$, let M_C be the midpoint of side AB . Shew that

$$\overrightarrow{CM_C} = \frac{1}{2} (\overrightarrow{CA} + \overrightarrow{CB}).$$

Solution: Since $\overrightarrow{AM_C} = \overrightarrow{M_C B}$, we have

$$\begin{aligned}\overrightarrow{CA} + \overrightarrow{CB} &= \overrightarrow{CM_C} + \overrightarrow{M_C A} + \overrightarrow{CM_C} + \overrightarrow{M_C B} \\ &= 2\overrightarrow{CM_C} - \overrightarrow{AM_C} + \overrightarrow{M_C B} \\ &= 2\overrightarrow{CM_C},\end{aligned}$$

which yields the desired result.

322 Theorem (Section Formula) Let APB be a straight line and λ and μ be real numbers such that

$$\frac{\| [A, P] \|}{\| [P, B] \|} = \frac{\lambda}{\mu}.$$

With $a = \overrightarrow{OA}$, $b = \overrightarrow{OB}$, and $p = \overrightarrow{OP}$, then

$$p = \frac{\lambda b + \mu a}{\lambda + \mu}. \quad (4.4)$$

Proof: Using Chasles' Rule for vectors,

$$\overrightarrow{AB} = \overrightarrow{AO} + \overrightarrow{OB} = -\mathbf{a} + \mathbf{b},$$

$$\overrightarrow{AP} = \overrightarrow{AO} + \overrightarrow{OP} = -\mathbf{a} + \mathbf{p}.$$

Also, using Chasles' Rule for bi-points,

$$[\mathbf{A}, \mathbf{P}] \mu = \lambda([\mathbf{P}, \mathbf{B}]) = \lambda([\mathbf{P}, \mathbf{A}] + [\mathbf{A}, \mathbf{B}]) = \lambda(-[\mathbf{A}, \mathbf{P}] + [\mathbf{A}, \mathbf{B}]),$$

whence

$$[\mathbf{A}, \mathbf{P}] = \frac{\lambda}{\lambda + \mu} [\mathbf{A}, \mathbf{B}] \implies \overrightarrow{AP} = \frac{\lambda}{\lambda + \mu} \overrightarrow{AB} \implies \mathbf{p} - \mathbf{a} = \frac{\lambda}{\lambda + \mu} (\mathbf{b} - \mathbf{a}).$$

On combining these formulæ

$$(\lambda + \mu)(\mathbf{p} - \mathbf{a}) = \lambda(\mathbf{b} - \mathbf{a}) \implies (\lambda + \mu)\mathbf{p} = \lambda\mathbf{b} + \mu\mathbf{a},$$

from where the result follows. \square




Figure 4.10: [A]. Problem 328.




Figure 4.11: [B]. Problem 328.




Figure 4.12: [C]. Problem 328.




Figure 4.13: [D]. Problem 328.




Figure 4.14: [E]. Problem 328.




Figure 4.15: [F]. Problem 328.

323 Problem Let a be a real number. Find the distance between $\begin{bmatrix} 1 \\ a \end{bmatrix}$ and $\begin{bmatrix} 1-a \\ 1 \end{bmatrix}$.

324 Problem Find all scalars λ for which $\|\lambda\mathbf{v}\| = \frac{1}{2}$, where

$$\mathbf{v} = \begin{bmatrix} 1 \\ -1 \end{bmatrix}.$$

325 Problem Given a pentagon ABCDE, find $\overrightarrow{AB} + \overrightarrow{BC} + \overrightarrow{CD} + \overrightarrow{DE} + \overrightarrow{EA}$.

326 Problem For which values of a will the vectors

$$\mathbf{a} = \begin{bmatrix} a+1 \\ a^2-1 \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} 2a+5 \\ a^2-4a+3 \end{bmatrix}$$

will be parallel?

327 Problem In $\triangle ABC$ let the midpoints of $[\mathbf{A}, \mathbf{B}]$ and $[\mathbf{A}, \mathbf{C}]$ be \mathbf{M}_B and \mathbf{M}_C , respectively. Put $\overrightarrow{M_C B} = \mathbf{x}$, $\overrightarrow{M_B C} = \mathbf{y}$, and $\overrightarrow{CA} = \mathbf{z}$. Express [A] $\overrightarrow{AB} + \overrightarrow{BC} + \overrightarrow{MC MB}$, [B] $\overrightarrow{AM} + \overrightarrow{MC MB} + \overrightarrow{MB C}$, [C] $\overrightarrow{AC} + \overrightarrow{MC A} - \overrightarrow{BM B}$ in terms of \mathbf{x} , \mathbf{y} , and \mathbf{z} .

328 Problem A circle is divided into three, four equal, or six equal parts (figures 4.10 through 4.15). Find the sum of the vectors. Assume that the divisions start or stop at the centre of the circle, as suggested in the figures.

329 Problem Diagonals are drawn in a square (figures ?? through ??). Find the vectorial sum $\mathbf{a} + \mathbf{b} + \mathbf{c}$. Assume that the diagonals either start, stop, or pass through the centre of the square, as suggested by the figures.

330 Problem Prove that the mid-points of the sides of a skew quadrilateral form the vertices of a parallelogram.

331 Problem ABCD is a parallelogram. E is the midpoint of [B, C] and F is the midpoint of [D, C]. Prove that

$$\overrightarrow{AC} + \overrightarrow{BD} = 2\overrightarrow{BC}.$$

332 Problem Let A, B be two points on the plane. Construct two points I and J such that

$$\overrightarrow{IA} = -3\overrightarrow{IB}, \quad \overrightarrow{JA} = -\frac{1}{3}\overrightarrow{JB},$$

and then demonstrate that for any arbitrary point M on the plane

$$\overrightarrow{MA} + 3\overrightarrow{MB} = 4\overrightarrow{MI}$$

and

$$3\overrightarrow{MA} + \overrightarrow{MB} = 4\overrightarrow{MJ}.$$

333 Problem You find an ancient treasure map in your great-grandfather's sea-chest. The sketch indicates that from the gallows you should walk to the oak tree, turn right 90° and walk a like distance, putting x at the point where you stop; then go back to the gallows, walk to the pine tree, turn left 90° , walk the same distance, mark point Y. Then you will find the treasure at the midpoint of the segment XY. So you charter a sailing vessel and go to the remote south-seas island. On arrival, you readily locate the oak and pine trees, but unfortunately, the gallows was struck by lightning, burned to dust and dispersed to the winds. No trace of it remains. What do you do?

4.3 Dot Product in \mathbb{R}^2

334 Definition Let $(\mathbf{a}, \mathbf{b}) \in (\mathbb{R}^2)^2$. The *dot product* $\mathbf{a} \bullet \mathbf{b}$ of \mathbf{a} and \mathbf{b} is defined by

$$\mathbf{a} \bullet \mathbf{b} = \begin{bmatrix} \mathbf{a}_1 \\ \mathbf{a}_2 \end{bmatrix} \cdot \begin{bmatrix} \mathbf{b}_1 \\ \mathbf{b}_2 \end{bmatrix} = \mathbf{a}_1 \mathbf{b}_1 + \mathbf{a}_2 \mathbf{b}_2.$$

The following properties of the dot product are easy to deduce from the definition.

DP1 Bilinearity

$$(\mathbf{x} + \mathbf{y}) \bullet \mathbf{z} = \mathbf{x} \bullet \mathbf{z} + \mathbf{y} \bullet \mathbf{z}, \quad \mathbf{x} \bullet (\mathbf{y} + \mathbf{z}) = \mathbf{x} \bullet \mathbf{y} + \mathbf{x} \bullet \mathbf{z} \quad (4.5)$$

DP2 Scalar Homogeneity

$$(\alpha \mathbf{x}) \bullet \mathbf{y} = \mathbf{x} \bullet (\alpha \mathbf{y}) = \alpha (\mathbf{x} \bullet \mathbf{y}), \quad \alpha \in \mathbb{R}. \quad (4.6)$$

DP3 Commutativity

$$\mathbf{x} \bullet \mathbf{y} = \mathbf{y} \bullet \mathbf{x} \quad (4.7)$$

DP4

$$\mathbf{x} \bullet \mathbf{x} \geq 0 \quad (4.8)$$

DP5

$$\mathbf{x} \bullet \mathbf{x} = 0 \Leftrightarrow \mathbf{x} = \mathbf{0} \quad (4.9)$$

DP6

$$\|\mathbf{x}\| = \sqrt{\mathbf{x} \bullet \mathbf{x}} \quad (4.10)$$

335 Example If we put

$$\mathbf{i} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad \mathbf{j} = \begin{bmatrix} 0 \\ 1 \end{bmatrix},$$

then we can write any vector $\mathbf{a} = \begin{bmatrix} \mathbf{a}_1 \\ \mathbf{a}_2 \end{bmatrix}$ as a sum

$$\mathbf{a} = \mathbf{a}_1 \mathbf{i} + \mathbf{a}_2 \mathbf{j}.$$

The vectors

$$\mathbf{i} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad \mathbf{j} = \begin{bmatrix} 0 \\ 1 \end{bmatrix},$$

satisfy $\mathbf{i} \cdot \mathbf{j} = 0$, and $\|\mathbf{i}\| = \|\mathbf{j}\| = 1$.

336 Definition Given vectors \mathbf{a} and \mathbf{b} , we define the angle between them, denoted by $\widehat{(\mathbf{a}, \mathbf{b})}$, as the angle between any two contiguous bi-point representatives of \mathbf{a} and \mathbf{b} .

337 Theorem

$$\mathbf{a} \cdot \mathbf{b} = \|\mathbf{a}\| \|\mathbf{b}\| \cos \widehat{(\mathbf{a}, \mathbf{b})}.$$

Proof: Using Al-Kashi's Law of Cosines on the length of the vectors, we have

$$\begin{aligned} \|\mathbf{b} - \mathbf{a}\|^2 &= \|\mathbf{a}\|^2 + \|\mathbf{b}\|^2 - 2\|\mathbf{a}\| \|\mathbf{b}\| \cos \widehat{(\mathbf{a}, \mathbf{b})} \\ \Leftrightarrow (\mathbf{b} - \mathbf{a}) \cdot (\mathbf{b} - \mathbf{a}) &= \|\mathbf{a}\|^2 + \|\mathbf{b}\|^2 - 2\|\mathbf{a}\| \|\mathbf{b}\| \cos \widehat{(\mathbf{a}, \mathbf{b})} \\ \Leftrightarrow \mathbf{b} \cdot \mathbf{b} - 2\mathbf{a} \cdot \mathbf{b} + \mathbf{a} \cdot \mathbf{a} &= \|\mathbf{a}\|^2 + \|\mathbf{b}\|^2 - 2\|\mathbf{a}\| \|\mathbf{b}\| \cos \widehat{(\mathbf{a}, \mathbf{b})} \\ \Leftrightarrow \|\mathbf{b}\|^2 - 2\mathbf{a} \cdot \mathbf{b} + \|\mathbf{b}\|^2 &= \|\mathbf{a}\|^2 + \|\mathbf{b}\|^2 - 2\|\mathbf{a}\| \|\mathbf{b}\| \cos \widehat{(\mathbf{a}, \mathbf{b})} \\ \Leftrightarrow \mathbf{a} \cdot \mathbf{b} &= \|\mathbf{a}\| \|\mathbf{b}\| \cos \widehat{(\mathbf{a}, \mathbf{b})}, \end{aligned}$$

as we wanted to show. \square

Putting $\widehat{(\mathbf{a}, \mathbf{b})} = \frac{\pi}{2}$ in Theorem 337 we obtain the following corollary.

338 Corollary Two vectors in \mathbb{R}^2 are perpendicular if and only if their dot product is 0.




Figure 4.19: Theorem 337.

339 Definition Two vectors are said to be *orthogonal* if they are perpendicular. If \mathbf{a} is orthogonal to \mathbf{b} , we write $\mathbf{a} \perp \mathbf{b}$.

340 Definition If $\mathbf{a} \perp \mathbf{b}$ and $\|\mathbf{a}\| = \|\mathbf{b}\| = 1$ we say that \mathbf{a} and \mathbf{b} are *orthonormal*.

 It follows that the vector $\mathbf{0}$ is simultaneously parallel and perpendicular to any vector!

341 Definition Let $\mathbf{a} \in \mathbb{R}^2$ be fixed. Then the *orthogonal space* to \mathbf{a} is defined and denoted by

$$\mathbf{a}^\perp = \{\mathbf{x} \in \mathbb{R}^2 : \mathbf{x} \perp \mathbf{a}\}.$$

Since $|\cos \theta| \leq 1$ we also have

342 Corollary (Cauchy-Bunyakovsky-Schwarz Inequality)

$$|\mathbf{a} \cdot \mathbf{b}| \leq \|\mathbf{a}\| \|\mathbf{b}\|.$$

343 Corollary (Triangle Inequality)

$$\|\mathbf{a} + \mathbf{b}\| \leq \|\mathbf{a}\| + \|\mathbf{b}\|.$$

Proof:

$$\begin{aligned} \|\mathbf{a} + \mathbf{b}\|^2 &= (\mathbf{a} + \mathbf{b}) \cdot (\mathbf{a} + \mathbf{b}) \\ &= \mathbf{a} \cdot \mathbf{a} + 2\mathbf{a} \cdot \mathbf{b} + \mathbf{b} \cdot \mathbf{b} \\ &\leq \|\mathbf{a}\|^2 + 2\|\mathbf{a}\| \|\mathbf{b}\| + \|\mathbf{b}\|^2 \\ &= (\|\mathbf{a}\| + \|\mathbf{b}\|)^2, \end{aligned}$$

from where the desired result follows. \square

344 Corollary (Pythagorean Theorem) If $\mathbf{a} \perp \mathbf{b}$ then

$$\|\mathbf{a} + \mathbf{b}\|^2 = \|\mathbf{a}\|^2 + \|\mathbf{b}\|^2.$$

Proof: Since $\mathbf{a} \cdot \mathbf{b} = 0$, we have

$$\begin{aligned} \|\mathbf{a} + \mathbf{b}\|^2 &= (\mathbf{a} + \mathbf{b}) \cdot (\mathbf{a} + \mathbf{b}) \\ &= \mathbf{a} \cdot \mathbf{a} + 2\mathbf{a} \cdot \mathbf{b} + \mathbf{b} \cdot \mathbf{b} \\ &= \mathbf{a} \cdot \mathbf{a} + 0 + \mathbf{b} \cdot \mathbf{b} \\ &= \|\mathbf{a}\|^2 + \|\mathbf{b}\|^2, \end{aligned}$$

from where the desired result follows. \square

345 Definition The *projection* of \mathbf{t} onto \mathbf{v} (or the \mathbf{v} -component of \mathbf{t}) is the vector

$$\text{proj}_{\mathbf{v}}^{\mathbf{t}} = (\cos(\widehat{\mathbf{v}, \mathbf{t}})) \|\mathbf{t}\| \frac{1}{\|\mathbf{v}\|} \mathbf{v},$$

where $\widehat{(\mathbf{v}, \mathbf{t})} \in [0; \pi]$ is the convex angle between \mathbf{v} and \mathbf{t} read in the positive sense.

 Given two vectors t and vector $v \neq 0$, find bi-point representatives of them having a common tail and join them together at their tails. The projection of t onto v is the “shadow” of t in the direction of v . To obtain proj_v^t we prolong v if necessary and drop a perpendicular line to it from the head of t . The projection is the portion between the common tails of the vectors and the point where this perpendicular meets t . See figure 4.20.



Figure 4.20: Vector Projections.

346 Corollary Let $a \neq 0$. Then

$$\text{proj}_a^x = \cos(\widehat{x, a}) \|x\| \frac{1}{\|a\|} a = \frac{x \cdot a}{\|a\|^2} a.$$

347 Theorem Let $a \in \mathbb{R}^2 \setminus \{0\}$. Then any $x \in \mathbb{R}^2$ can be decomposed as

$$x = u + v,$$

where $u \in \mathbb{R}a$ and $v \in a^\perp$.

Proof: We know that proj_a^x is parallel to a , so we take $u = \text{proj}_a^x$. This means that we must then take $v = x - \text{proj}_a^x$. We must demonstrate that v is indeed perpendicular to a . But this is clear, as

$$\begin{aligned} a \cdot v &= a \cdot x - a \cdot \text{proj}_a^x \\ &= a \cdot x - a \cdot \frac{x \cdot a}{\|a\|^2} a \\ &= a \cdot x - x \cdot a \\ &= 0, \end{aligned}$$

completing the proof. \square

348 Corollary Let $v \perp w$ be non-zero vectors in \mathbb{R}^2 . Then any vector $a \in \mathbb{R}^2$ has a unique representation as a linear combination of v, w ,

$$a = sv + tw, \quad (s, t) \in \mathbb{R}^2.$$

Proof: By Theorem 347, there exists a decomposition

$$a = sv + s'v',$$

where v' is orthogonal to v . But then $v' \parallel w$ and hence there exists $\alpha \in \mathbb{R}$ with $v' = \alpha w$. Taking $t = s'\alpha$ we achieve the decomposition

$$a = sv + tw.$$

To prove uniqueness, assume

$$sv + tw = a = pw + qw.$$

Then $(s - p)v = (q - t)w$. We must have $s = p$ and $q = t$ since otherwise v would be parallel to w . This completes the proof. \square

349 Corollary Let \mathbf{p}, \mathbf{q} be non-zero, non-parallel vectors in \mathbb{R}^2 . Then any vector $\mathbf{a} \in \mathbb{R}^2$ has a unique representation as a linear combination of \mathbf{p}, \mathbf{q} ,

$$\mathbf{a} = \mathbf{l}\mathbf{p} + \mathbf{m}\mathbf{q}, \quad (\mathbf{l}, \mathbf{m}) \in \mathbb{R}^2.$$

Proof: Consider $\mathbf{z} = \mathbf{q} - \text{proj}_{\mathbf{p}}^{\mathbf{q}}$. Clearly $\mathbf{p} \perp \mathbf{z}$ and so by Corollary 348, there exists unique $(s, t) \in \mathbb{R}^2$ such that

$$\begin{aligned} \mathbf{a} &= s\mathbf{p} + t\mathbf{z} \\ &= s\mathbf{p} - t\text{proj}_{\mathbf{p}}^{\mathbf{q}} + t\mathbf{q} \\ &= \left(s - t\frac{\mathbf{q} \cdot \mathbf{p}}{\|\mathbf{p}\|^2}\right)\mathbf{p} + t\mathbf{q}, \end{aligned}$$

establishing the result upon choosing $\mathbf{l} = s - t\frac{\mathbf{q} \cdot \mathbf{p}}{\|\mathbf{p}\|^2}$ and $\mathbf{m} = t$. \square

350 Example Let $\mathbf{p} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$, $\mathbf{q} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$. Write \mathbf{p} as the sum of two vectors, one parallel to \mathbf{q} and the other perpendicular to \mathbf{q} .

Solution: We use Theorem 347. We know that $\text{proj}_{\mathbf{q}}^{\mathbf{p}}$ is parallel to \mathbf{q} , and we find

$$\text{proj}_{\mathbf{q}}^{\mathbf{p}} = \frac{\mathbf{p} \cdot \mathbf{q}}{\|\mathbf{q}\|^2} \mathbf{q} = \frac{3}{5} \mathbf{q} = \begin{bmatrix} \frac{3}{5} \\ \frac{6}{5} \end{bmatrix}.$$

We also compute

$$\mathbf{p} - \text{proj}_{\mathbf{q}}^{\mathbf{p}} = \begin{bmatrix} 1 - \frac{3}{5} \\ 1 - \frac{6}{5} \end{bmatrix} = \begin{bmatrix} \frac{2}{5} \\ -\frac{1}{5} \end{bmatrix}.$$

Observe that

$$\begin{bmatrix} \frac{3}{5} \\ \frac{6}{5} \end{bmatrix} \cdot \begin{bmatrix} \frac{2}{5} \\ -\frac{1}{5} \end{bmatrix} = \frac{6}{25} - \frac{6}{25} = 0,$$

and the desired decomposition is

$$\begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} \frac{3}{5} \\ \frac{6}{5} \end{bmatrix} + \begin{bmatrix} \frac{2}{5} \\ -\frac{1}{5} \end{bmatrix}.$$




Figure 4.21: Orthocentre.

351 Example Prove that the altitudes of a triangle $\triangle ABC$ on the plane are concurrent. This point is called the *orthocentre* of the triangle.

Solution: Put $\mathbf{a} = \overrightarrow{OA}$, $\mathbf{b} = \overrightarrow{OB}$, $\mathbf{c} = \overrightarrow{OC}$. First observe that for any \mathbf{x} , we have, upon expanding,

$$(\mathbf{x} - \mathbf{a}) \cdot (\mathbf{b} - \mathbf{c}) + (\mathbf{x} - \mathbf{b}) \cdot (\mathbf{c} - \mathbf{a}) + (\mathbf{x} - \mathbf{c}) \cdot (\mathbf{a} - \mathbf{b}) = 0. \quad (4.11)$$

Let \mathbf{H} be the point of intersection of the altitude from A and the altitude from B . Then

$$\mathbf{0} = \overrightarrow{AH} \cdot \overrightarrow{CB} = (\overrightarrow{OH} - \overrightarrow{OA}) \cdot (\overrightarrow{OB} - \overrightarrow{OC}) = (\overrightarrow{OH} - \mathbf{a}) \cdot (\mathbf{b} - \mathbf{c}), \quad (4.12)$$

and

$$\mathbf{0} = \overrightarrow{BH} \cdot \overrightarrow{AC} = (\overrightarrow{OH} - \overrightarrow{OB}) \cdot (\overrightarrow{OC} - \overrightarrow{OA}) = (\overrightarrow{OH} - \mathbf{b}) \cdot (\mathbf{c} - \mathbf{a}). \quad (4.13)$$

Putting $\mathbf{x} = \overrightarrow{OH}$ in (4.11) and subtracting from it (4.12) and (4.13), we gather that

$$\mathbf{0} = (\overrightarrow{OH} - \mathbf{c}) \cdot (\mathbf{a} - \mathbf{b}) = \overrightarrow{CH} \cdot \overrightarrow{AB},$$

which gives the result.

352 Problem Determine the value of a so that $\begin{bmatrix} a \\ 1-a \end{bmatrix}$ be perpendicular to $\begin{bmatrix} 1 \\ -1 \end{bmatrix}$.

353 Problem Demonstrate that

$$(\mathbf{b} + \mathbf{c} = \mathbf{0}) \wedge (\|\mathbf{a}\| = \|\mathbf{b}\|) \iff (\mathbf{a} - \mathbf{b}) \cdot (\mathbf{a} - \mathbf{c}) = 0.$$

354 Problem Let $\mathbf{p} = \begin{bmatrix} 4 \\ 5 \end{bmatrix}$, $\mathbf{r} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$, $\mathbf{s} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$. Write \mathbf{p} as the sum of two vectors, one parallel to \mathbf{r} and the other parallel to \mathbf{s} .

355 Problem Prove that

$$\|\mathbf{a}\|^2 = (\mathbf{a} \cdot \mathbf{i})^2 + (\mathbf{a} \cdot \mathbf{j})^2.$$

356 Problem Let $\mathbf{a} \neq \mathbf{0} \neq \mathbf{b}$ be vectors in \mathbb{R}^2 such that $\mathbf{a} \cdot \mathbf{b} = 0$. Prove that

$$\alpha\mathbf{a} + \beta\mathbf{b} = \mathbf{0} \implies \alpha = \beta = 0.$$

357 Problem Let $(\mathbf{x}, \mathbf{y}) \in (\mathbb{R}^2)^2$ with $\|\mathbf{x}\| = \frac{3}{2}\|\mathbf{y}\|$. Show that $2\mathbf{x} + 3\mathbf{y}$ and $2\mathbf{x} - 3\mathbf{y}$ are perpendicular.

358 Problem Let \mathbf{a}, \mathbf{b} be fixed vectors in \mathbb{R}^2 . Prove that if

$$\forall \mathbf{v} \in \mathbb{R}^2, \mathbf{v} \cdot \mathbf{a} = \mathbf{v} \cdot \mathbf{b},$$

then $\mathbf{a} = \mathbf{b}$.

359 Problem Let $(\mathbf{a}, \mathbf{b}) \in (\mathbb{R}^2)^2$. Prove that

$$\|\mathbf{a} + \mathbf{b}\|^2 + \|\mathbf{a} - \mathbf{b}\|^2 = 2\|\mathbf{a}\|^2 + 2\|\mathbf{b}\|^2.$$

360 Problem Let \mathbf{u}, \mathbf{v} be vectors in \mathbb{R}^2 . Prove the *polarisation identity*:

$$\mathbf{u} \bullet \mathbf{v} = \frac{1}{4} (\|\mathbf{u} + \mathbf{v}\|^2 - \|\mathbf{u} - \mathbf{v}\|^2).$$

361 Problem Let \mathbf{x}, \mathbf{a} be non-zero vectors in \mathbb{R}^2 . Prove that

$$\text{proj}_{\mathbf{a}}^{\mathbf{x}} = \alpha \mathbf{a},$$

with $0 \leq \alpha \leq 1$.

362 Problem Let $(\lambda, \mathbf{a}) \in \mathbb{R} \times \mathbb{R}^2$ be fixed. Solve the equation

$$\mathbf{a} \bullet \mathbf{x} = \lambda$$

for $\mathbf{x} \in \mathbb{R}^2$.

4.4 Lines on the Plane

363 Definition Three points \mathbf{A} , \mathbf{B} , and \mathbf{C} are *collinear* if they lie on the same line.

It is clear that the points \mathbf{A} , \mathbf{B} , and \mathbf{C} are collinear if and only if \overrightarrow{AB} is parallel to \overrightarrow{AC} . Thus we have the following definition.

364 Definition The parametric equation with parameter $t \in \mathbb{R}$ of the straight line passing through the point

$P = \begin{pmatrix} p_1 \\ p_2 \end{pmatrix}$ in the direction of the vector $v \neq 0$ is

$$\begin{bmatrix} x - p_1 \\ y - p_2 \end{bmatrix} = tv.$$

If $r = \begin{bmatrix} x \\ y \end{bmatrix}$, then the equation of the line can be written in the form

$$r - p = tv. \quad (4.14)$$

The *Cartesian equation of a line* is an equation of the form $ax + by = c$, where $a^2 + b^2 \neq 0$. We write (AB) for the line passing through the points A and B .

365 Theorem Let $v \neq 0$ and let $n \perp v$. An alternative form for the equation of the line $r - p = tv$ is

$$(r - p) \cdot n = 0.$$

Moreover, the vector $\begin{bmatrix} a \\ b \end{bmatrix}$ is perpendicular to the line with Cartesian equation $ax + by = c$.

Proof: The first part follows at once by observing that $v \cdot n = 0$ and taking dot products to both sides of 4.14. For the second part observe that at least one of a and b is $\neq 0$. First assume that $a \neq 0$. Then we can put $y = t$ and $x = -\frac{b}{a}t + \frac{c}{a}$ and the parametric equation of this line is

$$\begin{bmatrix} x - \frac{c}{a} \\ y \end{bmatrix} = t \begin{bmatrix} -\frac{b}{a} \\ 1 \end{bmatrix},$$

and we have

$$\begin{bmatrix} -\frac{b}{a} \\ 1 \end{bmatrix} \cdot \begin{bmatrix} a \\ b \end{bmatrix} = -\frac{b}{a} \cdot a + b = 0.$$

Similarly if $b \neq 0$ we can put $x = t$ and $y = -\frac{a}{b}t + \frac{c}{b}$ and the parametric equation of this line is

$$\begin{bmatrix} x \\ y - \frac{c}{b} \end{bmatrix} = t \begin{bmatrix} 1 \\ -\frac{a}{b} \end{bmatrix},$$

and we have

$$\begin{bmatrix} 1 \\ -\frac{a}{b} \end{bmatrix} \cdot \begin{bmatrix} a \\ b \end{bmatrix} = a - \frac{a}{b} \cdot b = 0,$$

proving the theorem in this case. \square

 The vector $\begin{bmatrix} \frac{a}{\sqrt{a^2+b^2}} \\ \frac{b}{\sqrt{a^2+b^2}} \end{bmatrix}$ has norm 1 and is orthogonal to the line $ax + by = c$.

366 Example The equation of the line passing through $A = \begin{pmatrix} 2 \\ 3 \end{pmatrix}$ and in the direction of $v = \begin{bmatrix} -4 \\ 5 \end{bmatrix}$ is

$$\begin{bmatrix} x - 2 \\ y - 3 \end{bmatrix} = \lambda \begin{bmatrix} -4 \\ 5 \end{bmatrix}.$$

367 Example Find the equation of the line passing through $A = \begin{pmatrix} -1 \\ 1 \end{pmatrix}$ and $B = \begin{pmatrix} -2 \\ 3 \end{pmatrix}$.

Solution: The direction of this line is that of

$$\overrightarrow{AB} = \begin{bmatrix} -2 - (-1) \\ 3 - 1 \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \end{bmatrix}.$$

The equation is thus

$$\begin{bmatrix} x + 1 \\ y - 1 \end{bmatrix} = \lambda \begin{bmatrix} -1 \\ 2 \end{bmatrix}, \quad \lambda \in \mathbb{R}.$$

368 Example Suppose that $(m, b) \in \mathbb{R}^2$. Write the Cartesian equation of the line $y = mx + b$ in parametric form.

Solution: Here is a way. Put $x = t$. Then $y = mt + b$ and so the desired parametric form is

$$\begin{bmatrix} x \\ y - b \end{bmatrix} = t \begin{bmatrix} 1 \\ m \end{bmatrix}.$$

369 Example Let $(m_1, m_2, b_1, b_2) \in \mathbb{R}^4$, $m_1 m_2 \neq 0$. Consider the lines $L_1 : y = m_1 x + b_1$ and $L_2 : y = m_2 x + b_2$. By translating this problem in the language of vectors in \mathbb{R}^2 , shew that $L_1 \perp L_2$ if and only if $m_1 m_2 = -1$.

Solution: The parametric equations of the lines are

$$L_1 : \begin{bmatrix} x \\ y - b_1 \end{bmatrix} = s \begin{bmatrix} 1 \\ m_1 \end{bmatrix}, \quad L_2 : \begin{bmatrix} x \\ y - b_2 \end{bmatrix} = t \begin{bmatrix} 1 \\ m_2 \end{bmatrix}.$$

Put $\mathbf{v} = \begin{bmatrix} 1 \\ \mathbf{m}_1 \end{bmatrix}$ and $\mathbf{w} = \begin{bmatrix} 1 \\ \mathbf{m}_2 \end{bmatrix}$. Since the lines are perpendicular we must have $\mathbf{v} \cdot \mathbf{w} = 0$, which yields

$$0 = \mathbf{v} \cdot \mathbf{w} = 1(1) + \mathbf{m}_1(\mathbf{m}_2) \implies \mathbf{m}_1 \mathbf{m}_2 = -1.$$

370 Theorem (Distance Between a Point and a Line) Let $(\mathbf{r} - \mathbf{a}) \cdot \mathbf{n} = 0$ be a line passing through the point \mathbf{A} and perpendicular to vector \mathbf{n} . If \mathbf{B} is not a point on the line, then the distance from \mathbf{B} to the line is

$$\frac{|(\mathbf{a} - \mathbf{b}) \cdot \mathbf{n}|}{\|\mathbf{n}\|}.$$

If the line has Cartesian equation $a\mathbf{x} + b\mathbf{y} = c$, then this distance is

$$\frac{|ab_1 + bb_2 - c|}{\sqrt{a^2 + b^2}}.$$

Proof: Let \mathbf{R}_0 be the point on the line that is nearest to \mathbf{B} . Then $\overrightarrow{\mathbf{BR}_0} = \mathbf{r}_0 - \mathbf{b}$ is orthogonal to the line, and the distance we seek is

$$\|\text{proj}_{\mathbf{n}}^{\mathbf{r}_0 - \mathbf{b}}\| = \left\| \frac{(\mathbf{r}_0 - \mathbf{b}) \cdot \mathbf{n}}{\|\mathbf{n}\|^2} \mathbf{n} \right\| = \frac{|(\mathbf{r}_0 - \mathbf{b}) \cdot \mathbf{n}|}{\|\mathbf{n}\|}.$$

Since \mathbf{R}_0 is on the line, $\mathbf{r}_0 \cdot \mathbf{n} = \mathbf{a} \cdot \mathbf{n}$, and so

$$\|\text{proj}_{\mathbf{n}}^{\mathbf{r}_0 - \mathbf{b}}\| = \frac{|\mathbf{r}_0 \cdot \mathbf{n} - \mathbf{b} \cdot \mathbf{n}|}{\|\mathbf{n}\|} = \frac{|\mathbf{a} \cdot \mathbf{n} - \mathbf{b} \cdot \mathbf{n}|}{\|\mathbf{n}\|} = \frac{|(\mathbf{a} - \mathbf{b}) \cdot \mathbf{n}|}{\|\mathbf{n}\|},$$

as we wanted to shew.

If the line has Cartesian equation $a\mathbf{x} + b\mathbf{y} = c$, then at least one of a and b is $\neq 0$. Let us suppose $a \neq 0$, as the argument when $a = 0$ and $b \neq 0$ is similar. Then $a\mathbf{x} + b\mathbf{y} = c$ is equivalent to

$$\left(\begin{bmatrix} \mathbf{x} \\ \mathbf{y} \end{bmatrix} - \begin{bmatrix} \frac{c}{a} \\ 0 \end{bmatrix} \right) \cdot \begin{bmatrix} \mathbf{a} \\ \mathbf{b} \end{bmatrix} = 0.$$

We use the result obtained above with $\mathbf{a} = \begin{bmatrix} \frac{c}{a} \\ 0 \end{bmatrix}$, $\mathbf{n} = \begin{bmatrix} \mathbf{a} \\ \mathbf{b} \end{bmatrix}$, and $\mathbf{B} = \begin{bmatrix} \mathbf{b}_1 \\ \mathbf{b}_2 \end{bmatrix}$. Then $\|\mathbf{n}\| = \sqrt{a^2 + b^2}$

and

$$|(\mathbf{a} - \mathbf{b}) \cdot \mathbf{n}| = \left| \begin{bmatrix} \frac{c}{a} - \mathbf{b}_1 \\ -\mathbf{b}_2 \end{bmatrix} \cdot \begin{bmatrix} \mathbf{a} \\ \mathbf{b} \end{bmatrix} \right| = |c - ab_1 - bb_2| = |ab_1 + bb_2 - c|,$$

giving the result. \square

371 Example Recall that the medians of $\triangle ABC$ are lines joining the vertices of $\triangle ABC$ with the midpoints of the side opposite the vertex. Prove that the medians of a triangle are concurrent, that is, that they pass through a common point.

 This point of concurrency is called, alternatively, the isobarycentre, centroid, or centre of gravity of the triangle.

Solution: Let \mathbf{M}_A , \mathbf{M}_B , and \mathbf{M}_C denote the midpoints of the lines opposite A , B , and C , respectively. The equation

of the line passing through A and in the direction of $\overrightarrow{AM_A}$ is (with $\mathbf{r} = \begin{bmatrix} x \\ y \end{bmatrix}$)

$$\mathbf{r} = \overrightarrow{OA} + r\overrightarrow{AM_A}.$$

Similarly, the equation of the line passing through B and in the direction of $\overrightarrow{BM_B}$ is

$$\mathbf{r} = \overrightarrow{OB} + s\overrightarrow{BM_B}.$$

These two lines must intersect at a point G inside the triangle. We will show that \overrightarrow{GC} is parallel to $\overrightarrow{CM_C}$, which means that the three points G, C, M_C are collinear.

Now, $\exists (r_0, s_0) \in \mathbb{R}^2$ such that

$$\overrightarrow{OA} + r_0\overrightarrow{AM_A} = \overrightarrow{OG} = \overrightarrow{OB} + s_0\overrightarrow{BM_B},$$

that is

$$r_0\overrightarrow{AM_A} - s_0\overrightarrow{BM_B} = \overrightarrow{OB} - \overrightarrow{OA},$$

or

$$r_0(\overrightarrow{AB} + \overrightarrow{BM_A}) - s_0(\overrightarrow{BA} + \overrightarrow{AM_B}) = \overrightarrow{AB}.$$

Since M_A and M_B are the midpoints of $[B, C]$ and $[C, A]$ respectively, we have $2\overrightarrow{BM_A} = \overrightarrow{BC}$ and $2\overrightarrow{AM_B} = \overrightarrow{AC} = \overrightarrow{AB} + \overrightarrow{BC}$. The relationship becomes

$$\begin{aligned} r_0(\overrightarrow{AB} + \frac{1}{2}\overrightarrow{BC}) - s_0(-\overrightarrow{AB} + \frac{1}{2}\overrightarrow{AB} + \frac{1}{2}\overrightarrow{BC}) &= \overrightarrow{AB}, \\ (r_0 + \frac{s_0}{2} - 1)\overrightarrow{AB} &= (-\frac{r_0}{2} + \frac{s_0}{2})\overrightarrow{BC}. \end{aligned}$$

We must have

$$r_0 + \frac{s_0}{2} - 1 = 0,$$

$$-\frac{r_0}{2} + \frac{s_0}{2} = 0,$$

since otherwise the vectors \overrightarrow{AB} and \overrightarrow{BC} would be parallel, and the triangle would be degenerate. Solving, we find $s_0 = r_0 = \frac{2}{3}$. Thus we have $\overrightarrow{OA} + \frac{2}{3}\overrightarrow{AM_A} = \overrightarrow{OG}$, or $\overrightarrow{AG} = \frac{2}{3}\overrightarrow{AM_A}$, and similarly, $\overrightarrow{BG} = \frac{2}{3}\overrightarrow{BM_B}$.

From $\overrightarrow{AG} = \frac{2}{3}\overrightarrow{AM_A}$, we deduce $\overrightarrow{AG} = 2\overrightarrow{GM_A}$. Since M_A is the midpoint of $[B, C]$, we have $\overrightarrow{GB} + \overrightarrow{GC} = 2\overrightarrow{GM_A} = \overrightarrow{AG}$, which is equivalent to

$$\overrightarrow{GA} + \overrightarrow{GB} + \overrightarrow{GC} = 0.$$

As M_C is the midpoint of $[A, B]$ we have $\overrightarrow{GA} + \overrightarrow{GB} = 2\overrightarrow{GM_C}$. Thus

$$\mathbf{0} = \overrightarrow{GA} + \overrightarrow{GB} + \overrightarrow{GC} = 2\overrightarrow{GM_C} + \overrightarrow{GC}.$$

This means that $\overrightarrow{GC} = -2\overrightarrow{GM_C}$, that is, that they are parallel, and so the points G, C and M_C all lie on the same line. This achieves the desired result.

 The centroid of $\triangle ABC$ satisfies thus

$$\overrightarrow{GA} + \overrightarrow{GB} + \overrightarrow{GC} = \mathbf{0},$$

and divides the medians on the ratio $2 : 1$, reckoning from a vertex.

372 Problem Find the angle between the lines $2x - y = 1$ and $x - 3y = 1$.

373 Problem Find the equation of the line passing through $\begin{pmatrix} 1 \\ -1 \end{pmatrix}$ and in a direction perpendicular to $\begin{bmatrix} 2 \\ 1 \end{bmatrix}$.

374 Problem $\triangle ABC$ has centroid G , and $\triangle A'B'C'$ satisfies

$$\overrightarrow{AA'} + \overrightarrow{BB'} + \overrightarrow{CC'} = 0.$$

Prove that G is also the centroid of $\triangle A'B'C'$.

375 Problem Let $ABCD$ be a trapezoid, with bases $[A, B]$ and $[C, D]$. The lines (AC) and (BD) meet at E and the lines (AD) and (BC) meet at F . Prove that the line (EF) passes through the midpoints of $[A, B]$ and $[C, D]$ by proving the following steps.

- ① Let I be the midpoint of $[A, B]$ and let J be the point of intersection of the lines (FI) and (DC) . Prove that J is the midpoint of $[C, D]$. Deduce that F, I, J are collinear.
- ② Prove that E, I, J are collinear.

376 Problem Let $ABCD$ be a parallelogram.

- ① Let E and F be points such that

$$\overrightarrow{AE} = \frac{1}{4}\overrightarrow{AC} \quad \text{and} \quad \overrightarrow{AF} = \frac{3}{4}\overrightarrow{AC}.$$

Demonstrate that the lines (BE) and (DF) are parallel.

- ② Let I be the midpoint of $[A, D]$ and J be the midpoint of $[B, C]$. Demonstrate that the lines (AB) and (IJ) are parallel. What type of quadrilateral is $IEJF$?

377 Problem $ABCD$ is a parallelogram; point I is the midpoint of $[A, B]$. Point E is defined by the relation $\overrightarrow{IE} = \frac{1}{3}\overrightarrow{ID}$. Prove that

$$\overrightarrow{AE} = \frac{1}{3}(\overrightarrow{AB} + \overrightarrow{AD})$$

and prove that the points A, C, E are collinear.

378 Problem Put $\overrightarrow{OA} = \mathbf{a}$, $\overrightarrow{OB} = \mathbf{b}$, $\overrightarrow{OC} = \mathbf{c}$. Prove that A, B, C are collinear if and only if there exist real numbers α, β, γ , not all zero, such that

$$\alpha\mathbf{a} + \beta\mathbf{b} + \gamma\mathbf{c} = 0, \quad \alpha + \beta + \gamma = 0.$$

379 Problem Prove Desargues' Theorem: If $\triangle ABC$ and $\triangle A'B'C'$ (not necessarily in the same plane) are so positioned that (AA') , (BB') , (CC') all pass through the same point V and if (BC) and $(B'C')$ meet at L , (CA) and $(C'A')$ meet at M , and (AB) and $(A'B')$ meet at N , then L, M, N are collinear.

4.5 Vectors in \mathbb{R}^3

We now extend the notions studied for \mathbb{R}^2 to \mathbb{R}^3 . The rectangular coordinate form of a vector in \mathbb{R}^3 is

$$\mathbf{a} = \begin{bmatrix} \mathbf{a}_1 \\ \mathbf{a}_2 \\ \mathbf{a}_3 \end{bmatrix}.$$

In particular, if

$$\mathbf{i} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \quad \mathbf{j} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \quad \mathbf{k} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

then we can write any vector $\mathbf{a} = \begin{bmatrix} \mathbf{a}_1 \\ \mathbf{a}_2 \\ \mathbf{a}_3 \end{bmatrix}$ as a sum

$$\mathbf{a} = \mathbf{a}_1\mathbf{i} + \mathbf{a}_2\mathbf{j} + \mathbf{a}_3\mathbf{k}.$$

Given $\mathbf{a} = \begin{bmatrix} \mathbf{a}_1 \\ \mathbf{a}_2 \\ \mathbf{a}_3 \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} \mathbf{b}_1 \\ \mathbf{b}_2 \\ \mathbf{b}_3 \end{bmatrix}$, their dot product is

$$\mathbf{a} \cdot \mathbf{b} = \mathbf{a}_1 \mathbf{b}_1 + \mathbf{a}_2 \mathbf{b}_2 + \mathbf{a}_3 \mathbf{b}_3,$$

and

$$\|\mathbf{a}\| = \sqrt{\mathbf{a}_1^2 + \mathbf{a}_2^2 + \mathbf{a}_3^2}.$$

We also have

$$\mathbf{i} \cdot \mathbf{j} = \mathbf{j} \cdot \mathbf{k} = \mathbf{k} \cdot \mathbf{i} = 0,$$

and

$$\|\mathbf{i}\| = \|\mathbf{j}\| = \|\mathbf{k}\| = 1.$$

380 Definition A system of unit vectors $\mathbf{i}, \mathbf{j}, \mathbf{k}$ is *right-handed* if the shortest-route rotation which brings \mathbf{i} to coincide with \mathbf{j} is performed in a counter-clockwise manner. It is *left-handed* if the rotation is done in a clockwise manner.

To study points in space we must first agree on the orientation that we will give our coordinate system. We will use, unless otherwise noted, a right-handed orientation, as in figure 4.22.




Figure 4.22: Right-handed system.




Figure 4.23: Left-handed system.

The analogues of the Cauchy-Bunyakovsky-Schwarz and the Triangle Inequality also hold in \mathbb{R}^3 .

We now define the (standard) cross (wedge) product in \mathbb{R}^3 as a product satisfying the following properties.

381 Definition Let $(\mathbf{x}, \mathbf{y}, \mathbf{z}, \alpha) \in \mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{R}$. The wedge product $\times : \mathbb{R}^3 \times \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is a closed binary operation satisfying

CP1 Anti-commutativity:

$$\mathbf{x} \times \mathbf{y} = -(\mathbf{y} \times \mathbf{x}) \quad (4.15)$$

CP2 Bilinearity:

$$(\mathbf{x} + \mathbf{z}) \times \mathbf{y} = \mathbf{x} \times \mathbf{y} + \mathbf{z} \times \mathbf{y}, \quad \mathbf{x} \times (\mathbf{z} + \mathbf{y}) = \mathbf{x} \times \mathbf{z} + \mathbf{x} \times \mathbf{y} \quad (4.16)$$

CP3 Scalar homogeneity:

$$(\alpha \mathbf{x}) \times \mathbf{y} = \mathbf{x} \times (\alpha \mathbf{y}) = \alpha(\mathbf{x} \times \mathbf{y}) \quad (4.17)$$

CP4

$$\mathbf{x} \times \mathbf{x} = \mathbf{0} \quad (4.18)$$

CP5 Right-hand Rule:

$$\mathbf{i} \times \mathbf{j} = \mathbf{k}, \quad \mathbf{j} \times \mathbf{k} = \mathbf{i}, \quad \mathbf{k} \times \mathbf{i} = \mathbf{j} \quad (4.19)$$

382 Theorem Let $\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$ and $\mathbf{y} = \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix}$ be vectors in \mathbb{R}^3 . Then

$$\mathbf{x} \times \mathbf{y} = (x_2 y_3 - x_3 y_2) \mathbf{i} + (x_3 y_1 - x_1 y_3) \mathbf{j} + (x_1 y_2 - x_2 y_1) \mathbf{k}.$$

Proof: Since $\mathbf{i} \times \mathbf{i} = \mathbf{j} \times \mathbf{j} = \mathbf{k} \times \mathbf{k} = \mathbf{0}$ we have

$$\begin{aligned} (\mathbf{x}_1 \mathbf{i} + \mathbf{x}_2 \mathbf{j} + \mathbf{x}_3 \mathbf{k}) \times (\mathbf{y}_1 \mathbf{i} + \mathbf{y}_2 \mathbf{j} + \mathbf{y}_3 \mathbf{k}) &= \mathbf{x}_1 \mathbf{y}_2 \mathbf{i} \times \mathbf{j} + \mathbf{x}_1 \mathbf{y}_3 \mathbf{i} \times \mathbf{k} \\ &\quad + \mathbf{x}_2 \mathbf{y}_1 \mathbf{j} \times \mathbf{i} + \mathbf{x}_2 \mathbf{y}_3 \mathbf{j} \times \mathbf{k} \\ &\quad + \mathbf{x}_3 \mathbf{y}_1 \mathbf{k} \times \mathbf{i} + \mathbf{x}_3 \mathbf{y}_2 \mathbf{k} \times \mathbf{j} \\ &= \mathbf{x}_1 \mathbf{y}_2 \mathbf{k} - \mathbf{x}_1 \mathbf{y}_3 \mathbf{j} - \mathbf{x}_2 \mathbf{y}_1 \mathbf{k} \\ &\quad + \mathbf{x}_2 \mathbf{y}_3 \mathbf{i} + \mathbf{x}_3 \mathbf{y}_1 \mathbf{j} - \mathbf{x}_3 \mathbf{y}_2 \mathbf{i}, \end{aligned}$$

from where the theorem follows. \square

383 Example Find

$$\begin{bmatrix} 1 \\ 0 \\ -3 \end{bmatrix} \times \begin{bmatrix} 0 \\ 1 \\ 2 \end{bmatrix}.$$

Solution: We have

$$\begin{aligned} (\mathbf{i} - 3\mathbf{k}) \times (\mathbf{j} + 2\mathbf{k}) &= \mathbf{i} \times \mathbf{j} + 2\mathbf{i} \times \mathbf{k} - 3\mathbf{k} \times \mathbf{j} - 6\mathbf{k} \times \mathbf{k} \\ &= \mathbf{k} - 2\mathbf{j} - 3\mathbf{i} + 6\mathbf{0} \\ &= -3\mathbf{i} - 2\mathbf{j} + \mathbf{k}. \end{aligned}$$

Hence

$$\begin{bmatrix} 1 \\ 0 \\ -3 \end{bmatrix} \times \begin{bmatrix} 0 \\ 1 \\ 2 \end{bmatrix} = \begin{bmatrix} -3 \\ -2 \\ 1 \end{bmatrix}.$$

384 Theorem The cross product vector $\mathbf{x} \times \mathbf{y}$ is simultaneously perpendicular to \mathbf{x} and \mathbf{y} .

Proof: We will only check the first assertion, the second verification is analogous.

$$\begin{aligned}
 \mathbf{x} \bullet (\mathbf{x} \times \mathbf{y}) &= (x_1\mathbf{i} + x_2\mathbf{j} + x_3\mathbf{k}) \bullet ((x_2y_3 - x_3y_2)\mathbf{i} \\
 &\quad + (x_3y_1 - x_1y_3)\mathbf{j} + (x_1y_2 - x_2y_1)\mathbf{k}) \\
 &= x_1x_2y_3 - x_1x_3y_2 + x_2x_3y_1 - x_2x_1y_3 + x_3x_1y_2 - x_3x_2y_1 \\
 &= 0,
 \end{aligned}$$

completing the proof. \square

385 Theorem $\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) = (\mathbf{a} \bullet \mathbf{c})\mathbf{b} - (\mathbf{a} \bullet \mathbf{b})\mathbf{c}$.

Proof:

$$\begin{aligned}
 \mathbf{a} \times (\mathbf{b} \times \mathbf{c}) &= (a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k}) \times ((b_2c_3 - b_3c_2)\mathbf{i} + \\
 &\quad + (b_3c_1 - b_1c_3)\mathbf{j} + (b_1c_2 - b_2c_1)\mathbf{k}) \\
 &= a_1(b_3c_1 - b_1c_3)\mathbf{k} - a_1(b_1c_2 - b_2c_1)\mathbf{j} \\
 &\quad - a_2(b_2c_3 - b_3c_2)\mathbf{k} + a_2(b_1c_2 - b_2c_1)\mathbf{i} \\
 &\quad + a_3(b_2c_3 - b_3c_2)\mathbf{j} - a_3(b_3c_1 - b_1c_3)\mathbf{i} \\
 &= (a_1c_1 + a_2c_2 + a_3c_3)(b_1\mathbf{i} + b_2\mathbf{j} + b_3\mathbf{i}) \\
 &\quad + (-a_1b_1 - a_2b_2 - a_3b_3)(c_1\mathbf{i} + c_2\mathbf{j} + c_3\mathbf{i}) \\
 &= (\mathbf{a} \bullet \mathbf{c})\mathbf{b} - (\mathbf{a} \bullet \mathbf{b})\mathbf{c},
 \end{aligned}$$

completing the proof. \square

386 Theorem (Jacobi's Identity)

$$\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) + \mathbf{b} \times (\mathbf{c} \times \mathbf{a}) + \mathbf{c} \times (\mathbf{a} \times \mathbf{b}) = 0.$$

Proof: From Theorem 385 we have

$$\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) = (\mathbf{a} \bullet \mathbf{c})\mathbf{b} - (\mathbf{a} \bullet \mathbf{b})\mathbf{c},$$

$$\mathbf{b} \times (\mathbf{c} \times \mathbf{a}) = (\mathbf{b} \bullet \mathbf{a})\mathbf{c} - (\mathbf{b} \bullet \mathbf{c})\mathbf{a},$$

$$\mathbf{c} \times (\mathbf{a} \times \mathbf{b}) = (\mathbf{c} \bullet \mathbf{b})\mathbf{a} - (\mathbf{c} \bullet \mathbf{a})\mathbf{b},$$

and adding yields the result. \square

387 Theorem Let $\widehat{(\mathbf{x}, \mathbf{y})} \in [0; \pi]$ be the convex angle between two vectors \mathbf{x} and \mathbf{y} . Then

$$\|\mathbf{x} \times \mathbf{y}\| = \|\mathbf{x}\| \|\mathbf{y}\| \sin \widehat{(\mathbf{x}, \mathbf{y})}.$$

Proof: We have

$$\begin{aligned}
 \|\mathbf{x} \times \mathbf{y}\|^2 &= (x_2 y_3 - x_3 y_2)^2 + (x_3 y_1 - x_1 y_3)^2 + (x_1 y_2 - x_2 y_1)^2 \\
 &= x_2^2 y_3^2 - 2x_2 y_3 x_3 y_2 + x_3^2 y_2^2 + x_3^2 y_1^2 - 2x_3 y_1 x_1 y_3 + \\
 &\quad + x_1^2 y_3^2 + x_1^2 y_2^2 - 2x_1 y_2 x_2 y_1 + x_2^2 y_1^2 \\
 &= (x_1^2 + x_2^2 + x_3^2)(y_1^2 + y_2^2 + y_3^2) - (x_1 y_1 + x_2 y_2 + x_3 y_3)^2 \\
 &= \|\mathbf{x}\|^2 \|\mathbf{y}\|^2 - (\mathbf{x} \cdot \mathbf{y})^2 \\
 &= \|\mathbf{x}\|^2 \|\mathbf{y}\|^2 - \|\mathbf{x}\|^2 \|\mathbf{y}\|^2 \cos^2(\widehat{\mathbf{x}, \mathbf{y}}) \\
 &= \|\mathbf{x}\|^2 \|\mathbf{y}\|^2 \sin^2(\widehat{\mathbf{x}, \mathbf{y}}),
 \end{aligned}$$

whence the theorem follows. The Theorem is illustrated in Figure 4.24. Geometrically it means that the area of the parallelogram generated by joining \mathbf{x} and \mathbf{y} at their heads is $\|\mathbf{x} \times \mathbf{y}\|$. \square




Figure 4.24: Theorem 387.

The following corollaries are now obvious.

388 Corollary Two non-zero vectors \mathbf{x}, \mathbf{y} satisfy $\mathbf{x} \times \mathbf{y} = \mathbf{0}$ if and only if they are parallel.

389 Corollary (Lagrange's Identity)

$$\|\mathbf{x} \times \mathbf{y}\|^2 = \|\mathbf{x}\|^2 \|\mathbf{y}\|^2 - (\mathbf{x} \cdot \mathbf{y})^2.$$

390 Example Let $\mathbf{x} \in \mathbb{R}^3, \|\mathbf{x}\| = 1$. Find

$$\|\mathbf{x} \times \mathbf{i}\|^2 + \|\mathbf{x} \times \mathbf{j}\|^2 + \|\mathbf{x} \times \mathbf{k}\|^2.$$

Solution: By Lagrange's Identity,

$$\begin{aligned}
 \|\mathbf{x} \times \mathbf{i}\|^2 &= \|\mathbf{x}\|^2 \|\mathbf{i}\|^2 - (\mathbf{x} \cdot \mathbf{i})^2 = 1 - (\mathbf{x} \cdot \mathbf{i})^2, \\
 \|\mathbf{x} \times \mathbf{k}\|^2 &= \|\mathbf{x}\|^2 \|\mathbf{k}\|^2 - (\mathbf{x} \cdot \mathbf{k})^2 = 1 - (\mathbf{x} \cdot \mathbf{k})^2, \\
 \|\mathbf{x} \times \mathbf{j}\|^2 &= \|\mathbf{x}\|^2 \|\mathbf{j}\|^2 - (\mathbf{x} \cdot \mathbf{j})^2 = 1 - (\mathbf{x} \cdot \mathbf{j})^2,
 \end{aligned}$$

and since $(\mathbf{x} \cdot \mathbf{i})^2 + (\mathbf{x} \cdot \mathbf{j})^2 + (\mathbf{x} \cdot \mathbf{k})^2 = \|\mathbf{x}\|^2 = 1$, the desired sum equals $3 - 1 = 2$.

391 Problem Consider a tetrahedron ABCS. [A] Find $\overrightarrow{AB} + \overrightarrow{BC} + \overrightarrow{CS}$. [B] Find $\overrightarrow{AC} + \overrightarrow{CS} + \overrightarrow{SA} + \overrightarrow{AB}$.

392 Problem Find a vector simultaneously perpendicular to

$\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ and $\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$ and having norm 3.

393 Problem Find the area of the triangle whose vertices are

$$\text{at } P = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, Q = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \text{ and } R = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}.$$

394 Problem Prove or disprove! The cross product is associative.

395 Problem Prove that $x \times x = 0$ follows from the anti-commutativity of the cross product.

396 Problem Expand the product $(\mathbf{a} - \mathbf{b}) \times (\mathbf{a} + \mathbf{b})$.

397 Problem The vectors \mathbf{a}, \mathbf{b} are constant vectors. Solve the equation $\mathbf{a} \times (\mathbf{x} \times \mathbf{b}) = \mathbf{b} \times (\mathbf{x} \times \mathbf{a})$.

398 Problem The vectors $\mathbf{a}, \mathbf{b}, \mathbf{c}$ are constant vectors. Solve the system of equations

$$2\mathbf{x} + \mathbf{y} \times \mathbf{a} = \mathbf{b}, \quad 3\mathbf{y} + \mathbf{x} \times \mathbf{a} = \mathbf{c},$$

399 Problem Prove that there do not exist three unit vectors in \mathbb{R}^3 such that the angle between any two of them be $> \frac{2\pi}{3}$.

4.6 Planes and Lines in \mathbb{R}^3

400 Definition If bi-point representatives of a family of vectors in \mathbb{R}^3 lie on the same plane, we will say that the vectors are *coplanar* or parallel to the plane.

401 Lemma Let \mathbf{v}, \mathbf{w} in \mathbb{R}^3 be non-parallel vectors. Then every vector \mathbf{u} of the form

$$\mathbf{u} = \mathbf{av} + \mathbf{bw},$$

$((\mathbf{a}, \mathbf{b}) \in \mathbb{R}^2 \text{ arbitrary})$ is coplanar with both \mathbf{v} and \mathbf{w} . Conversely, any vector \mathbf{t} coplanar with both \mathbf{v} and \mathbf{w} can be uniquely expressed in the form

$$\mathbf{t} = \mathbf{pv} + \mathbf{qw}.$$

Proof: This follows at once from Corollary 349, since the operations occur on a plane, which can be identified with \mathbb{R}^2 . \square

A plane is determined by three non-collinear points. Suppose that \mathbf{A} , \mathbf{B} , and \mathbf{C} are non-collinear points on the

same plane and that $\mathbf{R} = \begin{pmatrix} x \\ y \\ z \end{pmatrix}$ is another arbitrary point on this plane. Since \mathbf{A} , \mathbf{B} , and \mathbf{C} are non-collinear, \overrightarrow{AB} and \overrightarrow{AC} , which are coplanar, are non-parallel. Since \overrightarrow{AR} also lies on the plane, we have by Lemma 401, that there exist real numbers p, q with

$$\overrightarrow{AR} = p\overrightarrow{AB} + q\overrightarrow{AC}.$$

By Chasles' Rule,

$$\overrightarrow{OR} = \overrightarrow{OA} + p(\overrightarrow{OB} - \overrightarrow{OA}) + q(\overrightarrow{OC} - \overrightarrow{OA}),$$

is the equation of a plane containing the three non-collinear points \mathbf{A} , \mathbf{B} , and \mathbf{C} . By letting $\mathbf{r} = \overrightarrow{OR}$, $\mathbf{a} = \overrightarrow{OA}$, etc., we deduce that

$$\mathbf{r} - \mathbf{a} = p(\mathbf{b} - \mathbf{a}) + q(\mathbf{c} - \mathbf{a}).$$

Thus we have the following definition.

402 Definition The *parametric equation* of a plane containing the point \mathbf{A} , and parallel to the vectors \mathbf{u} and \mathbf{v} is given by

$$\mathbf{r} - \mathbf{a} = p\mathbf{u} + q\mathbf{v}.$$

Componentwise this takes the form

$$x - a_1 = p u_1 + q v_1,$$

$$y - a_2 = p u_2 + q v_2,$$

$$z - a_3 = p u_3 + q v_3.$$

The *Cartesian equation* of a plane is an equation of the form $a_1 x + a_2 y + a_3 z = d$ with $(a_1, a_2, a_3, d) \in \mathbb{R}^4$ and $a_1^2 + a_2^2 + a_3^2 \neq 0$.

403 Example Find both the parametric equation and the Cartesian equation of the plane parallel to the vectors

$$\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \text{ and } \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \text{ and passing through the point } \begin{pmatrix} 0 \\ -1 \\ 2 \end{pmatrix}.$$

Solution: The desired parametric equation is

$$\begin{bmatrix} x \\ y+1 \\ z-2 \end{bmatrix} = s \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + t \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}.$$

This gives $s = z - 2$, $t = y + 1 - s = y + 1 - z + 2 = y - z + 3$ and $x = s + t = z - 2 + y - z + 3 = y + 1$. Hence the Cartesian equation is $x - y = 1$.

404 Theorem Let \mathbf{u} and \mathbf{v} be non-parallel vectors and let $\mathbf{r} - \mathbf{a} = p\mathbf{u} + q\mathbf{v}$ be the equation of the plane containing \mathbf{A} and parallel to the vectors \mathbf{u} and \mathbf{v} . If \mathbf{n} is simultaneously perpendicular to \mathbf{u} and \mathbf{v} then

$$(\mathbf{r} - \mathbf{a}) \cdot \mathbf{n} = 0.$$

Moreover, the vector $\begin{bmatrix} a \\ b \\ c \end{bmatrix}$ is normal to the plane with Cartesian equation $a_1 x + a_2 y + a_3 z = d$.

Proof: The first part is clear, as $\mathbf{u} \cdot \mathbf{n} = 0 = \mathbf{v} \cdot \mathbf{n}$. For the second part, recall that at least one of a, b, c is non-zero. Let us assume $a \neq 0$. The argument is similar if one of the other letters is non-zero and $a = 0$. In this case we can see that

$$x = \frac{d}{a} - \frac{b}{a}y - \frac{c}{a}z.$$

Put $y = s$ and $z = t$. Then

$$\begin{bmatrix} x - \frac{d}{a} \\ y \\ z \end{bmatrix} = s \begin{bmatrix} -\frac{b}{a} \\ 1 \\ 0 \end{bmatrix} + t \begin{bmatrix} -\frac{c}{a} \\ 0 \\ 1 \end{bmatrix}$$

is a parametric equation for the plane. \square

405 Example Find once again, by appealing to Theorem 404, the Cartesian equation of the plane parallel to the

vectors $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ and $\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$ and passing through the point $\begin{pmatrix} 0 \\ -1 \\ 2 \end{pmatrix}$.

Solution: The vector $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \times \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$ is normal to the plane. The plane has thus equation

$$\begin{bmatrix} x \\ y+1 \\ z-2 \end{bmatrix} \cdot \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix} = 0 \implies -x + y + 1 = 0 \implies x - y = 1,$$

as obtained before.

406 Theorem (Distance Between a Point and a Plane) Let $(\mathbf{r} - \mathbf{a}) \cdot \mathbf{n} = 0$ be a plane passing through the point \mathbf{A} and perpendicular to vector \mathbf{n} . If \mathbf{B} is not a point on the plane, then the distance from \mathbf{B} to the plane is

$$\frac{|(\mathbf{a} - \mathbf{b}) \cdot \mathbf{n}|}{\|\mathbf{n}\|}.$$

Proof: Let \mathbf{R}_0 be the point on the plane that is nearest to \mathbf{B} . Then $\overrightarrow{\mathbf{BR}_0} = \mathbf{r}_0 - \mathbf{b}$ is orthogonal to the plane, and the distance we seek is

$$\|\text{proj}_{\mathbf{n}}^{\mathbf{r}_0 - \mathbf{b}}\| = \left\| \frac{(\mathbf{r}_0 - \mathbf{b}) \cdot \mathbf{n}}{\|\mathbf{n}\|^2} \mathbf{n} \right\| = \frac{|(\mathbf{r}_0 - \mathbf{b}) \cdot \mathbf{n}|}{\|\mathbf{n}\|}.$$

Since \mathbf{R}_0 is on the plane, $\mathbf{r}_0 \cdot \mathbf{n} = \mathbf{a} \cdot \mathbf{n}$, and so

$$\|\text{proj}_{\mathbf{n}}^{\mathbf{r}_0 - \mathbf{b}}\| = \frac{|\mathbf{r}_0 \cdot \mathbf{n} - \mathbf{b} \cdot \mathbf{n}|}{\|\mathbf{n}\|} = \frac{|\mathbf{a} \cdot \mathbf{n} - \mathbf{b} \cdot \mathbf{n}|}{\|\mathbf{n}\|} = \frac{|(\mathbf{a} - \mathbf{b}) \cdot \mathbf{n}|}{\|\mathbf{n}\|},$$

as we wanted to shew. \square

 Given three planes in space, they may (i) be parallel (which allows for some of them to coincide), (ii) two may be parallel and the third intersect each of the other two at a line, (iii) intersect at a line, (iv) intersect at a point.

407 Definition The equation of a line passing through $\mathbf{A} \in \mathbb{R}^3$ in the direction of $\mathbf{v} \neq 0$ is given by

$$\mathbf{r} - \mathbf{a} = t\mathbf{v}, \quad t \in \mathbb{R}.$$

408 Theorem Put $\overrightarrow{\mathbf{OA}} = \mathbf{a}$, $\overrightarrow{\mathbf{OB}} = \mathbf{b}$, and $\overrightarrow{\mathbf{OC}} = \mathbf{c}$. Points $(\mathbf{A}, \mathbf{B}, \mathbf{C}) \in (\mathbb{R}^3)^3$ are collinear if and only if

$$\mathbf{a} \times \mathbf{b} + \mathbf{b} \times \mathbf{c} + \mathbf{c} \times \mathbf{a} = \mathbf{0}.$$

Proof: If the points $\mathbf{A}, \mathbf{B}, \mathbf{C}$ are collinear, then $\overrightarrow{\mathbf{AB}}$ is parallel to $\overrightarrow{\mathbf{AC}}$ and by Corollary 388, we must have

$$(\mathbf{c} - \mathbf{a}) \times (\mathbf{b} - \mathbf{a}) = \mathbf{0}.$$

Rearranging, gives

$$\mathbf{c} \times \mathbf{b} - \mathbf{c} \times \mathbf{a} - \mathbf{a} \times \mathbf{b} = \mathbf{0}.$$

Further rearranging completes the proof. \square

409 Theorem (Distance Between a Point and a Line) Let $L : \mathbf{r} = \mathbf{a} + \lambda\mathbf{v}$, $\mathbf{v} \neq 0$, be a line and let \mathbf{B} be a point not on L . Then the distance from \mathbf{B} to L is given by

$$\frac{\|(\mathbf{a} - \mathbf{b}) \times \mathbf{v}\|}{\|\mathbf{v}\|}.$$

Proof: If \mathbf{R}_0 —with position vector \mathbf{r}_0 —is the point on L that is at shortest distance from \mathbf{B} then $\overrightarrow{\mathbf{BR}_0}$ is perpendicular to the line, and so

$$\|\overrightarrow{\mathbf{BR}_0} \times \mathbf{v}\| = \|\overrightarrow{\mathbf{BR}_0}\| \|\mathbf{v}\| \sin \frac{\pi}{2} = \|\overrightarrow{\mathbf{BR}_0}\| \|\mathbf{v}\|.$$

The distance we must compute is $\|\overrightarrow{\mathbf{BR}_0}\| = \|\mathbf{r}_0 - \mathbf{b}\|$, which is then given by

$$\|\mathbf{r}_0 - \mathbf{b}\| = \frac{\|\overrightarrow{\mathbf{BR}_0} \times \mathbf{v}\|}{\|\mathbf{v}\|} = \frac{\|(\mathbf{r}_0 - \mathbf{b}) \times \mathbf{v}\|}{\|\mathbf{v}\|}.$$

Now, since \mathbf{R}_0 is on the line $\exists t_0 \in \mathbb{R}$ such that $\mathbf{r}_0 = \mathbf{a} + t_0\mathbf{v}$. Hence

$$(\mathbf{r}_0 - \mathbf{b}) \times \mathbf{v} = (\mathbf{a} - \mathbf{b}) \times \mathbf{v},$$

giving

$$\|\mathbf{r}_0 - \mathbf{b}\| = \frac{\|(\mathbf{a} - \mathbf{b}) \times \mathbf{v}\|}{\|\mathbf{v}\|},$$

proving the theorem. \square

 Given two lines in space, one of the following three situations might arise: (i) the lines intersect at a point, (ii) the lines are parallel, (iii) the lines are skew (one over the other, without intersecting).

410 Problem Find the equation of the plane passing through the points $(\mathbf{a}, 0, \mathbf{a})$, $(-\mathbf{a}, 1, 0)$, and $(0, 1, 2\mathbf{a})$ in \mathbb{R}^3 .

$(1, -1, -1)$ and containing the line $x = 2y = 3z$.

411 Problem Find the equation of plane containing the point $(1, 1, 1)$ and perpendicular to the line $x = 1 + t$, $y = -2t$, $z = 1 - t$.

413 Problem Find the equation of the plane perpendicular to the line $ax = by = cz$, $abc \neq 0$ and passing through the point $(1, 1, 1)$ in \mathbb{R}^3 .

412 Problem Find the equation of plane containing the point

414 Problem Find the equation of the line perpendicular to the plane $ax + a^2y + a^3z = 0$, $a \neq 0$ and passing through

the point $(0, 0, 1)$.

415 Problem The two planes

$$x - y - z = 1, \quad x - z = -1,$$

intersect at a line. Write the equation of this line in the form

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \mathbf{a} + t\mathbf{v}, \quad t \in \mathbb{R}.$$

416 Problem Find the equation of the plane passing through

the points $\begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$, $\begin{pmatrix} 2 \\ 1 \\ 1 \end{pmatrix}$ and parallel to the line $\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -1 \\ -2 \\ 0 \end{bmatrix} + t \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$.

417 Problem Points a, b, c in \mathbb{R}^3 are collinear and it is known that $a \times c = i - 2j$ and $a \times b = 2k - 3i$. Find $b \times c$.

418 Problem Find the equation of the plane which is equidis-

tant of the points $\begin{pmatrix} 3 \\ 2 \\ 1 \end{pmatrix}$ and $\begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}$.

419 Problem (Putnam Exam, 1980) Let S be the solid in three-dimensional space consisting of all points (x, y, z) satisfying the following system of six conditions:

$$x \geq 0, \quad y \geq 0, \quad z \geq 0,$$

$$x + y + z \leq 11,$$

$$2x + 4y + 3z \leq 36,$$

$$2x + 3z \leq 24.$$

Determine the number of vertices and the number of edges of S .

4.7 \mathbb{R}^n

As a generalisation of \mathbb{R}^2 and \mathbb{R}^3 we define \mathbb{R}^n as the set of n -tuples

$$\left\{ \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} : x_i \in \mathbb{R} \right\}.$$

The dot product of two vectors in \mathbb{R}^n is defined as

$$\mathbf{x} \cdot \mathbf{y} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \cdot \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix} = x_1 y_1 + x_2 y_2 + \cdots + x_n y_n.$$

The norm of a vector in \mathbb{R}^n is given by

$$\|\mathbf{x}\| = \sqrt{\mathbf{x} \cdot \mathbf{x}}.$$

As in the case of \mathbb{R}^2 and \mathbb{R}^3 we have

420 Theorem (Cauchy-Bunyakovsky-Schwarz Inequality) Given $(\mathbf{x}, \mathbf{y}) \in (\mathbb{R}^n)^2$ the following inequality holds

$$|\mathbf{x} \cdot \mathbf{y}| \leq \|\mathbf{x}\| \|\mathbf{y}\|.$$

Proof: Put $\mathbf{a} = \sum_{k=1}^n x_k^2$, $\mathbf{b} = \sum_{k=1}^n x_k y_k$, and $\mathbf{c} = \sum_{k=1}^n y_k^2$. Consider

$$f(t) = \sum_{k=1}^n (tx_k - y_k)^2 = t^2 \sum_{k=1}^n x_k^2 - 2t \sum_{k=1}^n x_k y_k + \sum_{k=1}^n y_k^2 = \mathbf{a}t^2 + \mathbf{b}t + \mathbf{c}.$$

This is a quadratic polynomial which is non-negative for all real t , so it must have complex roots. Its discriminant $\mathbf{b}^2 - 4\mathbf{a}\mathbf{c}$ must be non-positive, from where we gather

$$4 \left(\sum_{k=1}^n x_k y_k \right)^2 \leq 4 \left(\sum_{k=1}^n x_k^2 \right) \left(\sum_{k=1}^n y_k^2 \right).$$

This gives

$$|\mathbf{x} \bullet \mathbf{y}|^2 \leq \|\mathbf{x}\|^2 \|\mathbf{y}\|^2$$

from where we deduce the result. \square

421 Example Assume that $a_k, b_k, c_k, k = 1, \dots, n$, are positive real numbers. Shew that

$$\left(\sum_{k=1}^n a_k b_k c_k \right)^4 \leq \left(\sum_{k=1}^n a_k^4 \right) \left(\sum_{k=1}^n b_k^4 \right) \left(\sum_{k=1}^n c_k^2 \right)^2.$$

Solution: Using CBS on $\sum_{k=1}^n (a_k b_k) c_k$ once we obtain

$$\sum_{k=1}^n a_k b_k c_k \leq \left(\sum_{k=1}^n a_k^2 b_k^2 \right)^{1/2} \left(\sum_{k=1}^n c_k^2 \right)^{1/2}.$$

Using CBS again on $(\sum_{k=1}^n a_k^2 b_k^2)^{1/2}$ we obtain

$$\begin{aligned} \sum_{k=1}^n a_k b_k c_k &\leq (\sum_{k=1}^n a_k^2 b_k^2)^{1/2} (\sum_{k=1}^n c_k^2)^{1/2} \\ &\leq (\sum_{k=1}^n a_k^4)^{1/4} (\sum_{k=1}^n b_k^4)^{1/4} (\sum_{k=1}^n c_k^2)^{1/2}, \end{aligned}$$

which gives the required inequality.

422 Theorem (Triangle Inequality) Given $(\mathbf{x}, \mathbf{y}) \in (\mathbb{R}^n)^2$ the following inequality holds

$$\|\mathbf{x} + \mathbf{y}\| \leq \|\mathbf{x}\| + \|\mathbf{y}\|.$$

Proof: We have

$$\begin{aligned} \|\mathbf{a} + \mathbf{b}\|^2 &= (\mathbf{a} + \mathbf{b}) \bullet (\mathbf{a} + \mathbf{b}) \\ &= \mathbf{a} \bullet \mathbf{a} + 2\mathbf{a} \bullet \mathbf{b} + \mathbf{b} \bullet \mathbf{b} \\ &\leq \|\mathbf{a}\|^2 + 2\|\mathbf{a}\| \|\mathbf{b}\| + \|\mathbf{b}\|^2 \\ &= (\|\mathbf{a}\| + \|\mathbf{b}\|)^2, \end{aligned}$$

from where the desired result follows.

\square

We now consider a generalisation of the Euclidean norm. Given $p > 1$ and $\mathbf{x} \in \mathbb{R}^n$ we put

$$\|\mathbf{x}\|_p = \left(\sum_{k=1}^n |x_k|^p \right)^{1/p} \quad (4.20)$$

Clearly

$$\|\mathbf{x}\|_p \geq 0 \quad (4.21)$$

$$\|\mathbf{x}\|_p = 0 \Leftrightarrow \mathbf{x} = \mathbf{0} \quad (4.22)$$

$$\|\alpha \mathbf{x}\|_p = |\alpha| \|\mathbf{x}\|_p, \quad \alpha \in \mathbb{R} \quad (4.23)$$

We now prove analogues of the Cauchy-Bunyakovsky-Schwarz and the Triangle Inequality for $\|\cdot\|_p$. For this we need the following lemma.

423 Lemma (Young's Inequality) Let $p > 1$ and put $\frac{1}{p} + \frac{1}{q} = 1$. Then for $(a, b) \in ([0; +\infty[)^2$ we have

$$ab \leq \frac{a^p}{p} + \frac{b^q}{q}.$$

Proof: Let $0 < k < 1$, and consider the function

$$\begin{aligned} f : [0; +\infty[&\rightarrow \mathbb{R} \\ x &\mapsto x^k - k(x-1) \end{aligned}$$

Then $0 = f'(x) = kx^{k-1} - k \Leftrightarrow x = 1$. Since $f''(x) = k(k-1)x^{k-2} < 0$ for $0 < k < 1, x \geq 0$, $x = 1$ is a maximum point. Hence $f(x) \leq f(1)$ for $x \geq 0$, that is $x^k \leq 1 + k(x-1)$. Letting $k = \frac{1}{p}$ and

$x = \frac{a^p}{b^q}$ we deduce

$$\frac{a}{b^{q/p}} \leq 1 + \frac{1}{p} \left(\frac{a^p}{b^q} - 1 \right).$$

Rearranging gives

$$ab \leq b^{1+p/q} + \frac{a^p b^{1+p/q-p}}{p} - \frac{b^{1+p/q}}{p}$$

from where we obtain the inequality. \square

The promised generalisation of the Cauchy-Bunyakovsky-Schwarz Inequality is given in the following theorem.

424 Theorem (Hölder Inequality) Given $(x, y) \in (\mathbb{R}^n)^2$ the following inequality holds

$$|\mathbf{x} \cdot \mathbf{y}| \leq \|\mathbf{x}\|_p \|\mathbf{y}\|_q.$$

Proof: If $\|\mathbf{x}\|_p = 0$ or $\|\mathbf{y}\|_q = 0$ there is nothing to prove, so assume otherwise. From the Young Inequality we have

$$\frac{|\mathbf{x}_k|}{\|\mathbf{x}\|_p} \frac{|\mathbf{y}_k|}{\|\mathbf{y}\|_q} \leq \frac{|\mathbf{x}_k|^p}{\|\mathbf{x}\|_p^p p} + \frac{|\mathbf{y}_k|^q}{\|\mathbf{y}\|_q^q q}.$$

Adding, we deduce

$$\begin{aligned} \sum_{k=1}^n \frac{|\mathbf{x}_k|}{\|\mathbf{x}\|_p} \frac{|\mathbf{y}_k|}{\|\mathbf{y}\|_q} &\leq \frac{1}{\|\mathbf{x}\|_p^p p} \sum_{k=1}^n |\mathbf{x}_k|^p + \frac{1}{\|\mathbf{y}\|_q^q q} \sum_{k=1}^n |\mathbf{y}_k|^q \\ &= \frac{\|\mathbf{x}\|_p^p}{\|\mathbf{x}\|_p^p p} + \frac{\|\mathbf{y}\|_q^q}{\|\mathbf{y}\|_q^q q} \\ &= \frac{1}{p} + \frac{1}{q} \\ &= 1. \end{aligned}$$

This gives

$$\sum_{k=1}^n |x_k y_k| \leq \|x\|_p \|y\|_q.$$

The result follows by observing that

$$\left| \sum_{k=1}^n x_k y_k \right| \leq \sum_{k=1}^n |x_k y_k| \leq \|x\|_p \|y\|_q.$$

□

As a generalisation of the Triangle Inequality we have

425 Theorem (Minkowski Inequality) Let $p \in]1; +\infty[$. Given $(x, y) \in (\mathbb{R}^n)^2$ the following inequality holds

$$\|x + y\|_p \leq \|x\|_p + \|y\|_p.$$

Proof: From the triangle inequality for real numbers 1.6

$$|x_k + y_k|^p = |x_k + y_k| |x_k + y_k|^{p-1} \leq (|x_k| + |y_k|) |x_k + y_k|^{p-1}.$$

Adding

$$\sum_{k=1}^n |x_k + y_k|^p \leq \sum_{k=1}^n |x_k| |x_k + y_k|^{p-1} + \sum_{k=1}^n |y_k| |x_k + y_k|^{p-1}. \quad (4.24)$$

By the Hölder Inequality

$$\begin{aligned} \sum_{k=1}^n |x_k| |x_k + y_k|^{p-1} &\leq \left(\sum_{k=1}^n |x_k|^p \right)^{1/p} \left(\sum_{k=1}^n |x_k + y_k|^{(p-1)q} \right)^{1/q} \\ &= \left(\sum_{k=1}^n |x_k|^p \right)^{1/p} \left(\sum_{k=1}^n |x_k + y_k|^p \right)^{1/q} \\ &= \|x\|_p \|x + y\|_p^{p/q} \end{aligned} \quad (4.25)$$

In the same manner we deduce

$$\sum_{k=1}^n |y_k| |x_k + y_k|^{p-1} \leq \|y\|_p \|x + y\|_p^{p/q}. \quad (4.26)$$

Hence (4.24) gives

$$\|x + y\|_p^p = \sum_{k=1}^n |x_k + y_k|^p \leq \|x\|_p \|x + y\|_p^{p/q} + \|y\|_p \|x + y\|_p^{p/q},$$

from where we deduce the result. □

426 Problem Prove Lagrange's identity:

$$\begin{aligned} (\sum_{1 \leq j \leq n} a_j b_j)^2 &= (\sum_{1 \leq j \leq n} a_j^2) (\sum_{1 \leq j \leq n} b_j^2) \\ &\quad - \sum_{1 \leq k < j \leq n} (a_k b_j - a_j b_k)^2 \end{aligned}$$

and then deduce the CBS Inequality in \mathbb{R}^n .

$\sum_{i=1}^n a_i = 0$. Prove that $\sum_{1 \leq i < j \leq n} a_i \cdot a_j = -\frac{n}{2}$.

428 Problem Let $a_k > 0$. Use the CBS Inequality to show that

$$\left(\sum_{k=1}^n a_k^2 \right) \left(\sum_{k=1}^n \frac{1}{a_k^2} \right) \geq n^2.$$

427 Problem Let $a_i \in \mathbb{R}^n$ for $1 \leq i \leq n$ be unit vectors with

429 Problem Let $a_k \geq 0, 1 \leq k \leq n$ be arbitrary. Prove

that

$$\left(\sum_{k=1}^n a_k \right)^2 \leq \frac{n(n+1)(2n+1)}{6} \sum_{k=1}^n \frac{a_k^2}{k^2}.$$



Chapter 5

Vector Spaces

5.1 Vector Spaces

430 Definition A *vector space* $\langle \mathbf{V}, +, \cdot, \mathbb{F} \rangle$ over a field $\langle \mathbb{F}, +, \cdot \rangle$ is a non-empty set \mathbf{V} whose elements are called *vectors*, possessing two operations $+$ (vector addition), and \cdot (scalar multiplication) which satisfy the following axioms.

$$\forall(\mathbf{a}, \mathbf{b}, \mathbf{c}) \in \mathbf{V}^3, \forall(\alpha, \beta) \in \mathbb{F}^2,$$

VS1 Closure under vector addition :

$$\mathbf{a} + \mathbf{b} \in \mathbf{V}, \quad (5.1)$$

VS2 Closure under scalar multiplication

$$\alpha\mathbf{a} \in \mathbf{V}, \quad (5.2)$$

VS3 Commutativity

$$\mathbf{a} + \mathbf{b} = \mathbf{b} + \mathbf{a} \quad (5.3)$$

VS4 Associativity

$$(\mathbf{a} + \mathbf{b}) + \mathbf{c} = \mathbf{a} + (\mathbf{b} + \mathbf{c}) \quad (5.4)$$

VS5 Existence of an additive identity

$$\exists \mathbf{0} \in \mathbf{V} : \mathbf{a} + \mathbf{0} = \mathbf{a} + \mathbf{0} = \mathbf{a} \quad (5.5)$$

VS6 Existence of additive inverses

$$\exists -\mathbf{a} \in \mathbf{V} : \mathbf{a} + (-\mathbf{a}) = (-\mathbf{a}) + \mathbf{a} = \mathbf{0} \quad (5.6)$$

VS7 Distributive Law

$$\alpha(\mathbf{a} + \mathbf{b}) = \alpha\mathbf{a} + \alpha\mathbf{b} \quad (5.7)$$

VS8 Distributive Law

$$(\alpha + \beta)\mathbf{a} = \alpha\mathbf{a} + \beta\mathbf{a} \quad (5.8)$$

VS9

$$1_{\mathbb{F}}\mathbf{a} = \mathbf{a} \quad (5.9)$$

VS10

$$(\alpha\beta)\mathbf{a} = \alpha(\beta\mathbf{a}) \quad (5.10)$$

431 Example If n is a positive integer, then $\langle \mathbb{F}^n, +, \cdot, \mathbb{F} \rangle$ is a vector space by defining

$$(a_1, a_2, \dots, a_n) + (b_1, b_2, \dots, b_n) = (a_1 + b_1, a_2 + b_2, \dots, a_n + b_n),$$

$$\lambda(a_1, a_2, \dots, a_n) = (\lambda a_1, \lambda a_2, \dots, \lambda a_n).$$

In particular, $\langle \mathbb{Z}_2^2, +, \cdot, \mathbb{Z}_2 \rangle$ is a vector space with only four elements and we have seen the two-dimensional and tridimensional spaces $\langle \mathbb{R}^2, +, \cdot, \mathbb{R} \rangle$ and $\langle \mathbb{R}^3, +, \cdot, \mathbb{R} \rangle$.

432 Example $\langle M_{m \times n}(\mathbb{F}), +, \cdot, \mathbb{F} \rangle$ is a vector space under matrix addition and scalar multiplication of matrices.

433 Example If

$$\mathbb{F}[x] = \{a_0 + a_1x + a_2x^2 + \dots + a_nx^n : a_i \in \mathbb{F}, n \in \mathbb{N}\}$$

denotes the set of polynomials with coefficients in a field $\langle \mathbb{F}, +, \cdot \rangle$ then $\langle \mathbb{F}[x], +, \cdot, \mathbb{F} \rangle$ is a vector space, under polynomial addition and scalar multiplication of a polynomial.

434 Example If

$$\mathbb{F}_n[x] = \{a_0 + a_1x + a_2x^2 + \dots + a_kx^k : a_i \in \mathbb{F}, n \in \mathbb{N}, k \leq n\}$$

denotes the set of polynomials with coefficients in a field $\langle \mathbb{F}, +, \cdot \rangle$ and degree at most n , then $\langle \mathbb{F}_n[x], +, \cdot, \mathbb{F} \rangle$ is a vector space, under polynomial addition and scalar multiplication of a polynomial.

435 Example Let $k \in \mathbb{N}$ and let $C^k(\mathbb{R}^{[a;b]})$ denote the set of k -fold continuously differentiable real-valued functions defined on the interval $[a; b]$. Then $C^k(\mathbb{R}^{[a;b]})$ is a vector space under addition of functions and multiplication of a function by a scalar.

436 Example Let $p \in]1; +\infty[$. Consider the set of sequences $\{a_n\}_{n=0}^{\infty}$, $a_n \in \mathbb{C}$,

$$l^p = \left\{ \{a_n\}_{n=0}^{\infty} : \sum_{n=0}^{\infty} |a_n|^p < +\infty \right\}.$$

Then l^p is a vector space by defining addition as termwise addition of sequences and scalar multiplication as termwise multiplication:

$$\{a_n\}_{n=0}^{\infty} + \{b_n\}_{n=0}^{\infty} = \{(a_n + b_n)\}_{n=0}^{\infty},$$

$$\lambda\{a_n\}_{n=0}^{\infty} = \{\lambda a_n\}_{n=0}^{\infty}, \lambda \in \mathbb{C}.$$

All the axioms of a vector space follow trivially from the fact that we are adding complex numbers, except that we must prove that in l^p there is closure under addition and scalar multiplication. Since $\sum_{n=0}^{\infty} |a_n|^p < +\infty \Rightarrow \sum_{n=0}^{\infty} |\lambda a_n|^p < +\infty$ closure under scalar multiplication follows easily. To prove closure under addition, observe that if $z \in \mathbb{C}$ then $|z| \in \mathbb{R}_+$ and so by the Minkowski Inequality Theorem 425 we have

$$\begin{aligned} \left(\sum_{n=0}^{\infty} |a_n + b_n|^p \right)^{1/p} &\leq \left(\sum_{n=0}^{\infty} |a_n|^p \right)^{1/p} + \left(\sum_{n=0}^{\infty} |b_n|^p \right)^{1/p} \\ &\leq \left(\sum_{n=0}^{\infty} |a_n|^p \right)^{1/p} + \left(\sum_{n=0}^{\infty} |b_n|^p \right)^{1/p}. \end{aligned} \tag{5.11}$$

This in turn implies that the series on the left in (5.11) converges, and so we may take the limit as $N \rightarrow +\infty$ obtaining

$$\left(\sum_{n=0}^{\infty} |a_n + b_n|^p \right)^{1/p} \leq \left(\sum_{n=0}^{\infty} |a_n|^p \right)^{1/p} + \left(\sum_{n=0}^{\infty} |b_n|^p \right)^{1/p}. \tag{5.12}$$

Now (5.12) implies that the sum of two sequences in l^p is also in l^p , which demonstrates closure under addition.

437 Example The set

$$V = \{a + b\sqrt{2} + c\sqrt{3} : (a, b, c) \in \mathbb{Q}^3\}$$

with addition defined as

$$(a + b\sqrt{2} + c\sqrt{3}) + (a' + b'\sqrt{2} + c'\sqrt{3}) = (a + a') + (b + b')\sqrt{2} + (c + c')\sqrt{3},$$

and scalar multiplication defined as

$$\lambda(a + b\sqrt{2} + c\sqrt{3}) = (\lambda a) + (\lambda b)\sqrt{2} + (\lambda c)\sqrt{3},$$

constitutes a vector space over \mathbb{Q} .

438 Theorem In any vector space $\langle V, +, \cdot, \mathbb{F} \rangle$,

$$\forall \alpha \in \mathbb{F}, \quad \alpha \mathbf{0} = \mathbf{0}.$$

Proof: We have

$$\alpha \mathbf{0} = \alpha(\mathbf{0} + \mathbf{0}) = \alpha \mathbf{0} + \alpha \mathbf{0}.$$

Hence

$$\alpha \mathbf{0} - \alpha \mathbf{0} = \alpha \mathbf{0},$$

or

$$\mathbf{0} = \alpha \mathbf{0},$$

proving the theorem. \square

439 Theorem In any vector space $\langle V, +, \cdot, \mathbb{F} \rangle$,

$$\forall v \in V, \quad 0_{\mathbb{F}}v = \mathbf{0}.$$

Proof: We have

$$0_{\mathbb{F}}v = (0_{\mathbb{F}} + 0_{\mathbb{F}})v = 0_{\mathbb{F}}v + 0_{\mathbb{F}}v.$$

Therefore

$$0_{\mathbb{F}}v - 0_{\mathbb{F}}v = 0_{\mathbb{F}}v,$$

or

$$\mathbf{0} = 0_{\mathbb{F}}v,$$

proving the theorem. \square

440 Theorem In any vector space $\langle V, +, \cdot, \mathbb{F} \rangle$, $\alpha \in \mathbb{F}$, $v \in V$,

$$\alpha v = \mathbf{0} \implies \alpha = 0_{\mathbb{F}} \vee v = \mathbf{0}.$$

Proof: Assume that $\alpha \neq 0_{\mathbb{F}}$. Then α possesses a multiplicative inverse α^{-1} such that $\alpha^{-1}\alpha = 1_{\mathbb{F}}$. Thus

$$\alpha v = \mathbf{0} \implies \alpha^{-1}\alpha v = \alpha^{-1}\mathbf{0}.$$

By Theorem 439, $\alpha^{-1}\mathbf{0} = \mathbf{0}$. Hence

$$\alpha^{-1}\alpha v = \mathbf{0}.$$

Since by Axiom 5.9, we have $\alpha^{-1}\alpha v = 1_{\mathbb{F}}v = v$, and so we conclude that $v = \mathbf{0}$. \square

441 Theorem In any vector space $\langle V, +, \cdot, \mathbb{F} \rangle$,

$$\forall \alpha \in \mathbb{F}, \quad \forall v \in V, \quad (-\alpha)v = \alpha(-v) = -(\alpha v).$$

Proof: We have

$$0_{\mathbb{F}}\mathbf{v} = (\alpha + (-\alpha))\mathbf{v} = \alpha\mathbf{v} + (-\alpha)\mathbf{v},$$

whence

$$-(\alpha\mathbf{v}) + 0_{\mathbb{F}}\mathbf{v} = (-\alpha)\mathbf{v},$$

that is

$$-(\alpha\mathbf{v}) = (-\alpha)\mathbf{v}.$$

Similarly,

$$0 = \alpha(\mathbf{v} - \mathbf{v}) = \alpha\mathbf{v} + \alpha(-\mathbf{v}),$$

whence

$$-(\alpha\mathbf{v}) + 0 = \alpha(-\mathbf{v}),$$

that is

$$-(\alpha\mathbf{v}) = \alpha(-\mathbf{v}),$$

proving the theorem. \square

442 Problem Is \mathbb{R}^2 with vector addition and scalar multiplication defined as

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} x_1 + y_1 \\ x_2 + y_2 \end{bmatrix}, \quad \lambda \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} \lambda x_1 \\ 0 \end{bmatrix}$$

a vector space?

443 Problem Demonstrate that the commutativity axiom 5.3 is redundant.

444 Problem Let $V = \mathbb{R}^+ = [0; +\infty[$, the positive real numbers and $\mathbb{F} = \mathbb{R}$, the real numbers. Demonstrate that V

is a vector space over \mathbb{F} if vector addition is defined as $\mathbf{a} \oplus \mathbf{b} = \mathbf{ab}$, $(\mathbf{a}, \mathbf{b}) \in (\mathbb{R}^+)^2$ and scalar multiplication is defined as $\alpha \otimes \mathbf{a} = \mathbf{a}^\alpha$, $(\alpha, \mathbf{a}) \in (\mathbb{R}, \mathbb{R}^+)$.

445 Problem Let \mathbb{C} denote the complex numbers and \mathbb{R} denote the real numbers. Is \mathbb{C} a vector space over \mathbb{R} under ordinary addition and multiplication? Is \mathbb{R} a vector space over \mathbb{C} ?

446 Problem Construct a vector space with exactly 8 elements.

447 Problem Construct a vector space with exactly 9 elements.

5.2 Vector Subspaces

448 Definition Let $\langle V, +, \cdot, \mathbb{F} \rangle$ be a vector space. A non-empty subset $U \subseteq V$ which is also a vector space under the inherited operations of V is called a *vector subspace* of V .

449 Example Trivially, $X_1 = \{0\}$ and $X_2 = V$ are vector subspaces of V .

450 Theorem Let $\langle V, +, \cdot, \mathbb{F} \rangle$ be a vector space. Then $U \subseteq V$, $U \neq \emptyset$ is a subspace of V if and only if $\forall \alpha \in \mathbb{F}$ and $\forall (a, b) \in U^2$ it is verified that

$$a + \alpha b \in U.$$

Proof: Observe that U inherits commutativity, associativity and the distributive laws from V . Thus a non-empty $U \subseteq V$ is a vector subspace of V if (i) U is closed under scalar multiplication, that is, if $\alpha \in \mathbb{F}$ and $v \in U$, then $\alpha v \in U$; (ii) U is closed under vector addition, that is, if $(u, v) \in U^2$, then $u + v \in U$. Observe that (i) gives the existence of inverses in U , for take $\alpha = -1_{\mathbb{F}}$ and so $v \in U \implies -v \in U$. This coupled with (ii) gives the existence of the zero-vector, for $0 = v - v \in U$. Thus we need to prove that if a non-empty subset of V satisfies the property stated in the Theorem then it is closed under scalar multiplication and vector addition, and vice-versa, if a non-empty subset of V is closed under scalar multiplication and vector addition, then it satisfies the property stated in the Theorem. But this is trivial. \square

451 Example Shew that $X = \{A \in M_n(\mathbb{F}) : \text{tr}(A) = 0_{\mathbb{F}}\}$ is a subspace of $M_n(\mathbb{F})$.

Solution: Take $\mathbf{A}, \mathbf{B} \in X, \alpha \in \mathbb{R}$. Then

$$\text{tr}(\mathbf{A} + \alpha\mathbf{B}) = \text{tr}(\mathbf{A}) + \alpha\text{tr}(\mathbf{B}) = 0_{\mathbb{F}} + \alpha(0_{\mathbb{F}}) = 0_{\mathbb{F}}.$$

Hence $\mathbf{A} + \alpha\mathbf{B} \in X$, meaning that X is a subspace of $M_n(\mathbb{F})$.

452 Example Let $\mathbf{U} \in M_n(\mathbb{F})$ be an arbitrary but fixed. Shew that

$$\mathcal{C}_{\mathbf{U}} = \{\mathbf{A} \in M_n(\mathbb{F}) : \mathbf{AU} = \mathbf{UA}\}$$

is a subspace of $M_n(\mathbb{F})$.

Solution: Take $(\mathbf{A}, \mathbf{B}) \in (\mathcal{C}_{\mathbf{U}})^2$. Then $\mathbf{AU} = \mathbf{UA}$ and $\mathbf{BU} = \mathbf{UB}$. Now

$$(\mathbf{A} + \alpha\mathbf{B})\mathbf{U} = \mathbf{AU} + \alpha\mathbf{BU} = \mathbf{UA} + \alpha\mathbf{UB} = \mathbf{U}(\mathbf{A} + \alpha\mathbf{B}),$$

meaning that $\mathbf{A} + \alpha\mathbf{B} \in \mathcal{C}_{\mathbf{U}}$. Hence $\mathcal{C}_{\mathbf{U}}$ is a subspace of $M_n(\mathbb{F})$. $\mathcal{C}_{\mathbf{U}}$ is called the *commutator* of \mathbf{U} .

453 Theorem Let $X \subseteq V, Y \subseteq V$ be vector subspaces of a vector space $\langle V, +, \cdot, \mathbb{F} \rangle$. Then their intersection $X \cap Y$ is also a vector subspace of V .

Proof: Let $\alpha \in \mathbb{F}$ and $(\mathbf{a}, \mathbf{b}) \in (X \cap Y)^2$. Then clearly $(\mathbf{a}, \mathbf{b}) \in X$ and $(\mathbf{a}, \mathbf{b}) \in Y$. Since X is a vector subspace, $\mathbf{a} + \alpha\mathbf{b} \in X$ and since Y is a vector subspace, $\mathbf{a} + \alpha\mathbf{b} \in Y$. Thus

$$\mathbf{a} + \alpha\mathbf{b} \in X \cap Y$$

and so $X \cap Y$ is a vector subspace of V by virtue of Theorem 450. \square

 We will soon see that the only vector subspaces of $\langle \mathbb{R}^2, +, \cdot, \mathbb{R} \rangle$ are the set containing the zero-vector, any line through the origin, and \mathbb{R}^2 itself. The only vector subspaces of $\langle \mathbb{R}^3, +, \cdot, \mathbb{R} \rangle$ are the set containing the zero-vector, any line through the origin, any plane containing the origin and \mathbb{R}^3 itself.

454 Problem Prove that

$$X = \left\{ \begin{bmatrix} \mathbf{a} \\ \mathbf{b} \\ \mathbf{c} \\ \mathbf{d} \end{bmatrix} \in \mathbb{R}^4 : \mathbf{a} - \mathbf{b} - 3\mathbf{d} = 0 \right\}$$

is a vector subspace of \mathbb{R}^4 .

455 Problem Prove that

$$X = \left\{ \begin{bmatrix} \mathbf{a} \\ 2\mathbf{a} - 3\mathbf{b} \\ 5\mathbf{b} \\ \mathbf{a} + 2\mathbf{b} \\ \mathbf{a} \end{bmatrix} : \mathbf{a}, \mathbf{b} \in \mathbb{R} \right\}$$

is a vector subspace of \mathbb{R}^5 .

456 Problem Let $\mathbf{a} \in \mathbb{R}^n$ be a fixed vector. Demonstrate that

$$X = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a} \cdot \mathbf{x} = 0\}$$

is a subspace of \mathbb{R}^n .

457 Problem Let $\mathbf{a} \in \mathbb{R}^3$ be a fixed vector. Demonstrate that

$$X = \{\mathbf{x} \in \mathbb{R}^3 : \mathbf{a} \times \mathbf{x} = 0\}$$

is a subspace of \mathbb{R}^3 .

458 Problem Let $\mathbf{A} \in M_{m \times n}(\mathbb{F})$ be a fixed matrix. Demonstrate that

$$S = \{X \in M_{n \times 1}(\mathbb{F}) : \mathbf{AX} = 0_{m \times 1}\}$$

is a subspace of $M_{n \times 1}(\mathbb{F})$.

459 Problem Prove that the set $X \subseteq M_n(\mathbb{F})$ of upper triangular matrices is a subspace of $M_n(\mathbb{F})$.

460 Problem Prove that the set $X \subseteq M_n(\mathbb{F})$ of symmetric matrices is a subspace of $M_n(\mathbb{F})$.

461 Problem Prove that the set $X \subseteq M_n(\mathbb{F})$ of skew-symmetric matrices is a subspace of $M_n(\mathbb{F})$.

462 Problem Prove that the following subsets are not subspaces of the given vector space. Here you must say which of the axioms for a vector space fail.

$$\textcircled{1} \quad \left\{ \begin{bmatrix} a \\ b \\ 0 \end{bmatrix} : a, b \in \mathbb{R}, a^2 + b^2 = 1 \right\} \subseteq \mathbb{R}^3$$

$$\textcircled{2} \quad \left\{ \begin{bmatrix} a \\ b \\ 0 \end{bmatrix} : a, b \in \mathbb{R}^2, ab = 0 \right\} \subseteq \mathbb{R}^3$$

$$\textcircled{3} \quad \left\{ \begin{bmatrix} a & b \\ 0 & 0 \end{bmatrix} : (a, b) \in \mathbb{R}^2, a + b^2 = 0 \right\} \subseteq M_{2 \times 2}(\mathbb{R})$$

463 Problem Let $\langle V, +, \cdot, \mathbb{F} \rangle$ be a vector space, and let $U_1 \subseteq V$ and $U_2 \subseteq V$ be vector subspaces. Prove that if $U_1 \cup U_2$ is a vector subspace of V , then either $U_1 \subseteq U_2$ or $U_2 \subseteq U_1$.

464 Problem Let V a vector space over a field \mathbb{F} . If \mathbb{F} is infinite, show that V is not the set-theoretic union of a finite number of proper subspaces.

465 Problem Give an example of a finite vector space V over a finite field \mathbb{F} such that

$$V = V_1 \cup V_2 \cup V_3,$$

where the V_k are proper subspaces.

5.3 Linear Independence

466 Definition Let $(\lambda_1, \lambda_2, \dots, \lambda_n) \in \mathbb{F}^n$. Then the vectorial sum

$$\sum_{j=1}^n \lambda_j a_j$$

is said to be a *linear combination* of the vectors $a_i \in V, 1 \leq i \leq n$.

467 Example Any matrix $\begin{bmatrix} a & b \\ c & d \end{bmatrix} \in M_2(\mathbb{R})$ can be written as a linear combination of the matrices

$$\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix},$$

for

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} = a \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + b \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} + c \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} + d \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}.$$

468 Example Any polynomial of degree at most 2, say $a + bx + cx^2 \in \mathbb{R}_2[x]$ can be written as a linear combination of $1, x - 1$, and $x^2 - x + 2$, for

$$a + bx + cx^2 = (a - c)(1) + (b + c)(x - 1) + c(x^2 - x + 2).$$

Generalising the notion of two parallel vectors, we have

469 Definition The vectors $a_i \in V, 1 \leq i \leq n$, are *linearly dependent* or *tied* if

$$\exists (\lambda_1, \lambda_2, \dots, \lambda_n) \in \mathbb{F}^n \setminus \{0\} \text{ such that } \sum_{j=1}^n \lambda_j a_j = 0,$$

that is, if there is a non-trivial linear combination of them adding to the zero vector.

470 Definition The vectors $\mathbf{a}_i \in V, 1 \leq i \leq n$, are *linearly independent* or *free* if they are not linearly dependent. That is, if $(\lambda_1, \lambda_2, \dots, \lambda_n) \in \mathbb{F}^n$ then

$$\sum_{j=1}^n \lambda_j \mathbf{a}_j = \mathbf{0} \implies \lambda_1 = \lambda_2 = \dots = \lambda_n = 0_{\mathbb{F}}.$$

 A family of vectors is linearly independent if and only if the only linear combination of them giving the zero-vector is the trivial linear combination.

471 Example

$$\left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix}, \begin{bmatrix} 7 \\ 8 \\ 9 \end{bmatrix} \right\}$$

is a tied family of vectors in \mathbb{R}^3 , since

$$(1) \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + (-2) \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} + (1) \begin{bmatrix} 7 \\ 8 \\ 9 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}.$$

472 Example Let \mathbf{u}, \mathbf{v} be linearly independent vectors in some vector space over a field \mathbb{F} with characteristic different from 2. Show that the two new vectors $\mathbf{x} = \mathbf{u} - \mathbf{v}$ and $\mathbf{y} = \mathbf{u} + \mathbf{v}$ are also linearly independent.

Solution: Assume that $a(\mathbf{u} - \mathbf{v}) + b(\mathbf{u} + \mathbf{v}) = \mathbf{0}$. Then

$$(a + b)\mathbf{u} + (a - b)\mathbf{v} = \mathbf{0}.$$

Since \mathbf{u}, \mathbf{v} are linearly independent, the above coefficients must be 0, that is, $a + b = 0_{\mathbb{F}}$ and $a - b = 0_{\mathbb{F}}$. But this gives $2a = 2b = 0_{\mathbb{F}}$, which implies $a = b = 0_{\mathbb{F}}$, if the characteristic of the field is not 2. This proves the linear independence of $\mathbf{u} - \mathbf{v}$ and $\mathbf{u} + \mathbf{v}$.

473 Theorem Let $\mathbf{A} \in M_{m \times n}(\mathbb{F})$. Then the columns of \mathbf{A} are linearly independent if and only the only solution to the system $\mathbf{AX} = \mathbf{0}_m$ is the trivial solution.

Proof: Let $\mathbf{A}_1, \dots, \mathbf{A}_n$ be the columns of \mathbf{A} . Since

$$x_1 \mathbf{A}_1 + x_2 \mathbf{A}_2 + \dots + x_n \mathbf{A}_n = \mathbf{AX},$$

the result follows. \square

474 Theorem Any family

$$\{\mathbf{0}, \mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k\}$$

containing the zero-vector is linearly dependent.

Proof: This follows at once by observing that

$$1_{\mathbb{F}} \mathbf{0} + 0_{\mathbb{F}} \mathbf{u}_1 + 0_{\mathbb{F}} \mathbf{u}_2 + \dots + 0_{\mathbb{F}} \mathbf{u}_k = \mathbf{0}$$

is a non-trivial linear combination of these vectors equalling the zero-vector. \square

475 Problem Shew that

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right\}$$

forms a free family of vectors in \mathbb{R}^3 .

476 Problem Prove that the set

$$\left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ -1 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \\ 1 \end{bmatrix} \right\}$$

is a linearly independent set of vectors in \mathbb{R}^4 and shew that

$X = \begin{bmatrix} 1 \\ 2 \\ 1 \\ 1 \end{bmatrix}$ can be written as a linear combination of these vectors.

477 Problem Let $(a, b) \in (\mathbb{R}^3)^2$ and assume that $a \cdot b = 0$ and that a and b are linearly independent. Prove that $a, b, a \times b$ are linearly independent.

478 Problem Let $a_i \in \mathbb{R}^n$, $1 \leq i \leq k$ ($k \leq n$) be k non-zero vectors such that $a_i \cdot a_j = 0$ for $i \neq j$. Prove that these k vectors are linearly independent.

479 Problem Let $(u, v) \in (\mathbb{R}^n)^2$. Prove that $|u \cdot v| = \|u\| \|v\|$ if and only if u and v are linearly dependent.

480 Problem Prove that

$$\left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \right\}$$

is a linearly independent family over \mathbb{R} . Write $\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$ as a linear combination of these matrices.

481 Problem Let $\{v_1, v_2, v_3, v_4\}$ be a linearly independent family of vectors. Prove that the family

$$\{v_1 + v_2, v_2 + v_3, v_3 + v_4, v_4 + v_1\}$$

is not linearly independent.

482 Problem Let $\{v_1, v_2, v_3\}$ be linearly independent vectors in \mathbb{R}^5 . Are the vectors

$$b_1 = 3v_1 + 2v_2 + 4v_3,$$

$$b_2 = v_1 + 4v_2 + 2v_3,$$

$$b_3 = 9v_1 + 4v_2 + 3v_3,$$

$$b_4 = v_1 + 2v_2 + 5v_3,$$

linearly independent? Prove or disprove!

483 Problem Is the family $\{1, \sqrt{2}\}$ linearly independent over \mathbb{Q} ?

484 Problem Is the family $\{1, \sqrt{2}\}$ linearly independent over \mathbb{R} ?

485 Problem Consider the vector space

$$V = \{a + b\sqrt{2} + c\sqrt{3} : (a, b, c) \in \mathbb{Q}^3\}.$$

1. Shew that $\{1, \sqrt{2}, \sqrt{3}\}$ are linearly independent over \mathbb{Q} .

2. Express

$$\frac{1}{1 - \sqrt{2}} + \frac{2}{\sqrt{12} - 2}$$

as a linear combination of $\{1, \sqrt{2}, \sqrt{3}\}$.

486 Problem Let f, g, h belong to $C^\infty(\mathbb{R})$ (the space of infinitely continuously differentiable real-valued functions defined on the real line) and be given by

$$f(x) = e^x, g(x) = e^{2x}, h(x) = e^{3x}.$$

Shew that f, g, h are linearly independent over \mathbb{R} .

487 Problem Let f, g, h belong to $C^\infty(\mathbb{R})$ be given by

$$f(x) = \cos^2 x, g(x) = \sin^2 x, h(x) = \cos 2x.$$

Shew that f, g, h are linearly dependent over \mathbb{R} .

5.4 Spanning Sets

488 Definition A family $\{u_1, u_2, \dots, u_k, \dots\} \subseteq V$ is said to *span* or *generate* V if every $v \in V$ can be written as a linear combination of the u_j 's.

489 Theorem If $\{u_1, u_2, \dots, u_k, \dots\} \subseteq V$ spans V , then any superset

$$\{w, u_1, u_2, \dots, u_k, \dots\} \subseteq V$$

also spans V .

Proof: This follows at once from

$$\sum_{i=1}^l \lambda_i u_i = 0_F w + \sum_{i=1}^l \lambda_i u_i.$$

□

490 Example The family of vectors

$$\left\{ i = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, j = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, k = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$$

spans \mathbb{R}^3 since given $\begin{bmatrix} a \\ b \\ c \end{bmatrix} \in \mathbb{R}^3$ we may write

$$\begin{bmatrix} a \\ b \\ c \end{bmatrix} = ai + bj + ck.$$

491 Example Prove that the family of vectors

$$\left\{ t_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, t_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, t_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right\}$$

spans \mathbb{R}^3 .

Solution: This follows from the identity

$$\begin{bmatrix} a \\ b \\ c \end{bmatrix} = (a - b) \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + (b - c) \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + c \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = (a - b)t_1 + (b - c)t_2 + ct_3.$$

492 Example Since

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} = a \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + b \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} + c \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} + d \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$$

the set of matrices $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$ is a spanning set for $M_2(\mathbb{R})$

493 Example The set

$$\{1, x, x^2, x^3, \dots, x^n, \dots\}$$

spans $\mathbb{R}[x]$, the set of polynomials with real coefficients and indeterminate x .

494 Definition The *span* of a family of vectors $\{u_1, u_2, \dots, u_k, \dots\}$ is the set of all finite linear combinations obtained from the u_i 's. We denote the span of $\{u_1, u_2, \dots, u_k, \dots\}$ by

$$\text{span}(u_1, u_2, \dots, u_k, \dots).$$

495 Theorem Let $\langle V, +, \cdot, \mathbb{F} \rangle$ be a vector space. Then

$$\text{span}(u_1, u_2, \dots, u_k, \dots) \subseteq V$$

is a vector subspace of V .

Proof: Let $\alpha \in \mathbb{F}$ and let

$$x = \sum_{k=1}^l a_k u_k, \quad y = \sum_{k=1}^l b_k u_k,$$

be in $\text{span}(u_1, u_2, \dots, u_k, \dots)$ (some of the coefficients might be $0_{\mathbb{F}}$). Then

$$x + \alpha y = \sum_{k=1}^l (a_k + \alpha b_k) u_k \in \text{span}(u_1, u_2, \dots, u_k, \dots),$$

and so $\text{span}(u_1, u_2, \dots, u_k, \dots)$ is a subspace of V . \square

496 Corollary $\text{span}(u_1, u_2, \dots, u_k, \dots) \subseteq V$ is the smallest vector subspace of V (in the sense of set inclusion) containing the set

$$\{u_1, u_2, \dots, u_k, \dots\}.$$

Proof: If $W \subseteq V$ is a vector subspace of V containing the set

$$\{u_1, u_2, \dots, u_k, \dots\}$$

then it contains every finite linear combination of them, and hence, it contains $\text{span}(u_1, u_2, \dots, u_k, \dots)$.

\square

497 Example If $A \in M_2(\mathbb{R})$, $A \in \text{span}\left(\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}\right)$ then A has the form

$$a \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + b \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} + c \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} a & c \\ c & b \end{bmatrix},$$

i.e., this family spans the set of all symmetric 2×2 matrices over \mathbb{R} .

498 Theorem Let \mathbf{V} be a vector space over a field \mathbb{F} and let $(\mathbf{v}, \mathbf{w}) \in \mathbf{V}^2$, $\gamma \in \mathbb{F} \setminus \{0_{\mathbb{F}}\}$. Then

$$\text{span}(\mathbf{v}, \mathbf{w}) = \text{span}(\mathbf{v}, \gamma\mathbf{w}).$$

Proof: *The equality*

$$\mathbf{av} + \mathbf{bw} = \mathbf{av} + (\mathbf{b}\gamma^{-1})(\gamma\mathbf{w}),$$

proves the statement. \square

499 Theorem Let \mathbf{V} be a vector space over a field \mathbb{F} and let $(\mathbf{v}, \mathbf{w}) \in \mathbf{V}^2$, $\gamma \in \mathbb{F}$. Then

$$\text{span}(\mathbf{v}, \mathbf{w}) = \text{span}(\mathbf{w}, \mathbf{v} + \gamma\mathbf{w}).$$

Proof: *This follows from the equality*

$$\mathbf{av} + \mathbf{bw} = \mathbf{a}(\mathbf{v} + \gamma\mathbf{w}) + (\mathbf{b} - \mathbf{a}\gamma)\mathbf{w}.$$

\square

500 Problem Let $\mathbb{R}_3[x]$ denote the set of polynomials with degree at most 3 and real coefficients. Prove that the set

$$\{1, 1+x, (1+x)^2, (1+x)^3\}$$

spans $\mathbb{R}_3[x]$.

501 Problem Shew that $\begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix} \notin \text{span} \left(\begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix} \right)$.

502 Problem What is $\text{span} \left(\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \right)$?

503 Problem Prove that

$$\text{span} \left(\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \right) = M_2(\mathbb{R}).$$

504 Problem For the vectors in \mathbb{R}^3 ,

$$\mathbf{a} = \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} 1 \\ 3 \\ 2 \end{bmatrix}, \quad \mathbf{c} = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \quad \mathbf{d} = \begin{bmatrix} 3 \\ 8 \\ 5 \end{bmatrix},$$

prove that $\text{span}(\mathbf{a}, \mathbf{b}) = \text{span}(\mathbf{c}, \mathbf{d})$.

5.5 Bases

505 Definition A family $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k, \dots\} \subseteq \mathbf{V}$ is said to be a *basis* of \mathbf{V} if (i) they are linearly independent, (ii) they span \mathbf{V} .

506 Example The family

$$\mathbf{e}_i = \begin{bmatrix} 0_{\mathbb{F}} \\ \vdots \\ 0_{\mathbb{F}} \\ 1_{\mathbb{F}} \\ 0_{\mathbb{F}} \\ \vdots \\ 0_{\mathbb{F}} \end{bmatrix},$$

where there is a $1_{\mathbb{F}}$ on the i -th slot and $0_{\mathbb{F}}$'s on the other $n - 1$ positions, is a basis for \mathbb{F}^n .

507 Theorem Let $\langle V, +, \cdot, \mathbb{F} \rangle$ be a vector space and let

$$\mathbf{U} = \{u_1, u_2, \dots, u_k, \dots\} \subseteq V$$

be a family of linearly independent vectors in V which is maximal in the sense that if \mathbf{U}' is any other family of vectors of V properly containing \mathbf{U} then \mathbf{U}' is a dependent family. Then \mathbf{U} forms a basis for V .

Proof: Since \mathbf{U} is a linearly independent family, we need only to prove that it spans V . Take $v \in V$. If $v \in \mathbf{U}$ then there is nothing to prove, so assume that $v \in V \setminus \mathbf{U}$. Consider the set $\mathbf{U}' = \mathbf{U} \cup \{v\}$. This set properly contains \mathbf{U} , and so, by assumption, it forms a dependent family. There exists scalars $\alpha_0, \alpha_1, \dots, \alpha_n$ such that

$$\alpha_0 v + \alpha_1 u_1 + \cdots + \alpha_n u_n = 0.$$

Now, $\alpha_0 \neq 0_{\mathbb{F}}$, otherwise the u_i would be linearly dependent. Hence α_0^{-1} exists and we have

$$v = -\alpha_0^{-1}(\alpha_1 u_1 + \cdots + \alpha_n u_n),$$

and so the u_i span V . \square

 From Theorem 507 it follows that to shew that a vector space has a basis it is enough to shew that it has a maximal linearly independent set of vectors. Such a proof requires something called Zörn's Lemma, and it is beyond our scope. We dodge the whole business by taking as an axiom that every vector space possesses a basis.

508 Theorem (Steinitz Replacement Theorem) Let $\langle V, +, \cdot, \mathbb{F} \rangle$ be a vector space and let $\mathbf{U} = \{u_1, u_2, \dots\} \subseteq V$. Let $\mathbf{W} = \{w_1, w_2, \dots, w_k\}$ be an independent family of vectors in $\text{span}(\mathbf{U})$. Then there exist k of the u_i 's, say $\{u_1, u_2, \dots, u_k\}$ which may be replaced by the w_i 's in such a way that

$$\text{span}(w_1, w_2, \dots, w_k, u_{k+1}, \dots) = \text{span}(\mathbf{U}).$$

Proof: We prove this by induction on k . If $k = 1$, then

$$w_1 = \alpha_1 u_1 + \alpha_2 u_2 + \cdots + \alpha_n u_n$$

for some n and scalars α_i . There is an $\alpha_i \neq 0_{\mathbb{F}}$, since otherwise $w_1 = 0$ contrary to the assumption that the w_i are linearly independent. After reordering, we may assume that $\alpha_1 \neq 0_{\mathbb{F}}$. Hence

$$u_1 = \alpha_1^{-1}(w_1 - (\alpha_2 u_2 + \cdots + \alpha_n u_n)),$$

and so $\mathbf{u}_1 \in \text{span}(\mathbf{w}_1, \mathbf{u}_2, \dots)$ and

$$\text{span}(\mathbf{w}_1, \mathbf{u}_2, \dots) = \text{span}(\mathbf{u}_1, \mathbf{u}_2, \dots).$$

Assume now that the theorem is true for any set of fewer than k independent vectors. We may thus assume that that $\{\mathbf{u}_1, \dots\}$ has more than $k - 1$ vectors and that

$$\text{span}(\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_{k-1}, \mathbf{u}_k, \dots) = \text{span}(\mathbf{U}).$$

Since $\mathbf{w}_k \in \mathbf{U}$ we have

$$\mathbf{w}_k = \beta_1 \mathbf{w}_1 + \beta_2 \mathbf{w}_2 + \dots + \beta_{k-1} \mathbf{w}_{k-1} + \gamma_k \mathbf{u}_k + \gamma_{k+1} \mathbf{u}_{k+1} + \dots + \gamma_m \mathbf{u}_m.$$

If all the $\gamma_i = 0_F$, then the $\{\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_k\}$ would be linearly dependent, contrary to assumption. Thus there is a $\gamma_i \neq 0_F$, and after reordering, we may assume that $\gamma_k \neq 0_F$. We have therefore

$$\mathbf{u}_k = \gamma_k^{-1} (\mathbf{w}_k - (\beta_1 \mathbf{w}_1 + \beta_2 \mathbf{w}_2 + \dots + \beta_{k-1} \mathbf{w}_{k-1} + \gamma_{k+1} \mathbf{u}_{k+1} + \dots + \gamma_m \mathbf{u}_m)).$$

But this means that

$$\text{span}(\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_k, \mathbf{u}_{k+1}, \dots) = \text{span}(\mathbf{U}).$$

This finishes the proof. \square

509 Corollary Let $\{\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_n\}$ be an independent family of vectors with $\mathbf{V} = \text{span}(\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_n)$. If we also have $\mathbf{V} = \text{span}(\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_v)$, then

1. $n \leq v$,
2. $n = v$ if and only if the $\{y_1, y_2, \dots, y_v\}$ are a linearly independent family.
3. Any basis for \mathbf{V} has exactly n elements.

Proof:

1. In the Steinitz Replacement Theorem 508 replace the first n \mathbf{u}_i 's by the \mathbf{w}_i 's and $n \leq v$ follows.
2. If $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_v\}$ are a linearly independent family, then we may interchange the rôle of the \mathbf{w}_i and \mathbf{u}_i obtaining $v \leq n$. Conversely, if $v = n$ and if the \mathbf{u}_i are dependent, we could express some \mathbf{u}_i as a linear combination of the remaining $v - 1$ vectors, and thus we would have shewn that some $v - 1$ vectors span \mathbf{V} . From (1) in this corollary we would conclude that $n \leq v - 1$, contradicting $n = v$.
3. This follows from the definition of what a basis is and from (2) of this corollary.

\square

510 Definition The *dimension* of a vector space $\langle \mathbf{V}, +, \cdot, \mathbb{F} \rangle$ is the number of elements of any of its bases, and we denote it by $\dim \mathbf{V}$.

511 Theorem Any linearly independent family of vectors

$$\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$$

in a vector space \mathbf{V} can be completed into a family

$$\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k, \mathbf{y}_{k+1}, \mathbf{y}_{k+2}, \dots\}$$

so that this latter family become a basis for \mathbf{V} .

Proof: Take any basis $\{\mathbf{u}_1, \dots, \mathbf{u}_k, \mathbf{u}_{k+1}, \dots\}$ and use Steinitz Replacement Theorem 508. \square

512 Corollary If $\mathbf{U} \subseteq \mathbf{V}$ is a vector subspace of a finite dimensional vector space \mathbf{V} then $\dim \mathbf{U} \leq \dim \mathbf{V}$.

Proof: Since any basis of \mathbf{U} can be extended to a basis of \mathbf{V} , it follows that the number of elements of the basis of \mathbf{U} is at most as large as that for \mathbf{V} . \square

513 Example Find a basis and the dimension of the space generated by the set of symmetric matrices in $M_n(\mathbb{R})$.

Solution: Let $E_{ij} \in M_n(\mathbb{R})$ be the $n \times n$ matrix with a 1 on the ij -th position and 0's everywhere else. For $1 \leq i < j \leq n$, consider the $\binom{n}{2} = \frac{n(n-1)}{2}$ matrices $A_{ij} = E_{ij} + E_{ji}$. The A_{ij} have a 1 on the ij -th and ji -th position and 0's everywhere else. They, together with the n matrices E_{ii} , $1 \leq i \leq n$ constitute a basis for the space of symmetric matrices. The dimension of this space is thus

$$\frac{n(n-1)}{2} + n = \frac{n(n+1)}{2}.$$

514 Theorem Let $\{u_1, \dots, u_n\}$ be vectors in \mathbb{R}^n . Then the u 's form a basis if and only if the $n \times n$ matrix \mathbf{A} formed by taking the u 's as the columns of \mathbf{A} is invertible.

Proof: Since we have the right number of vectors, it is enough to prove that the u 's are linearly independent. But if $X = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$, then

$$x_1 u_1 + \cdots + x_n u_n = \mathbf{A}X.$$

If \mathbf{A} is invertible, then $\mathbf{A}X = \mathbf{0}_n \implies X = \mathbf{A}^{-1}\mathbf{0}_n = \mathbf{0}_n$, meaning that $x_1 = x_2 = \cdots = x_n = 0$, so the u 's are linearly independent.

Conversely, assume that the u 's are linearly independent. Then the equation $\mathbf{A}X = \mathbf{0}_n$ has a unique solution. Let $r = \text{rank } (\mathbf{A})$ and let $(P, Q) \in (GL_n(\mathbb{R}))^2$ be matrices such that $\mathbf{A} = P^{-1}D_{n,n,r}Q^{-1}$, where $D_{n,n,r}$ is the Hermite normal form of \mathbf{A} . Thus

$$\mathbf{A}X = \mathbf{0}_n \implies P^{-1}D_{n,n,r}Q^{-1}X = \mathbf{0}_n \implies D_{n,n,r}Q^{-1}X = \mathbf{0}_n.$$

Put $Q^{-1}X = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix}$. Then

$$D_{n,n,r}Q^{-1}X = \mathbf{0}_n \implies y_1 e_1 + \cdots + y_r e_r = \mathbf{0}_n,$$

where e_j is the n -dimensional column vector with a 1 on the j -th slot and 0's everywhere else. If $r < n$ then y_{r+1}, \dots, y_n can be taken arbitrarily and so there would not be a unique solution, a contradiction. Hence $r = n$ and \mathbf{A} is invertible. \square

515 Problem In problem 455 we saw that

$$X = \left\{ \begin{bmatrix} a \\ 2a - 3b \\ 5b \\ a + 2b \\ a \end{bmatrix} : a, b \in \mathbb{R} \right\}$$

is a vector subspace of \mathbb{R}^5 . Find a basis for this subspace.

516 Problem Let $\{v_1, v_2, v_3, v_4, v_5\}$ be a basis for a vector space V over a field \mathbb{F} . Prove that

$$\{v_1 + v_2, v_2 + v_3, v_3 + v_4, v_4 + v_5, v_5 + v_1\}$$

is also a basis for V .

517 Problem Find a basis for the solution space of the system of $n+1$ linear equations of $2n$ unknowns

$$x_1 + x_2 + \cdots + x_n = 0,$$

$$x_2 + x_3 + \cdots + x_{n+1} = 0,$$

⋮

$$x_{n+1} + x_{n+2} + \cdots + x_{2n} = 0.$$

518 Problem Prove that the set

$$X = \{(a, b, c, d) | b + 2c = 0\} \subseteq \mathbb{R}^4$$

is a vector subspace of \mathbb{R}^4 . Find its dimension and a basis for X .

5.6 Coordinates

524 Theorem Let $\{v_1, v_2, \dots, v_n\}$ be a basis for a vector space V . Then any $v \in V$ has a unique representation

$$v = a_1 v_1 + a_2 v_2 + \cdots + a_n v_n.$$

Proof: Let

$$v = b_1 v_1 + b_2 v_2 + \cdots + b_n v_n$$

be another representation of v . Then

$$0 = (a_1 - b_1)v_1 + (a_2 - b_2)v_2 + \cdots + (a_n - b_n)v_n.$$

Since $\{v_1, v_2, \dots, v_n\}$ forms a basis for V , they are a linearly independent family. Thus we must have

$$a_1 - b_1 = a_2 - b_2 = \cdots = a_n - b_n = 0_{\mathbb{F}},$$

that is

$$a_1 = b_1; a_2 = b_2; \dots; a_n = b_n,$$

proving uniqueness. \square

519 Problem Prove that the dimension of the vector subspace of lower triangular $n \times n$ matrices is $\frac{n(n+1)}{2}$ and find a basis for this space.

520 Problem Find a basis and the dimension of

$$X = \text{span} \left(v_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, v_2 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 0 \end{bmatrix}, v_3 = \begin{bmatrix} 2 \\ 2 \\ 2 \\ 1 \end{bmatrix} \right).$$

521 Problem Find a basis and the dimension of

$$X = \text{span} \left(v_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, v_2 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 0 \end{bmatrix}, v_3 = \begin{bmatrix} 2 \\ 2 \\ 2 \\ 2 \end{bmatrix} \right).$$

522 Problem Find a basis and the dimension of

$$X = \text{span} \left(v_1 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, v_2 = \begin{bmatrix} 1 & 0 \\ 2 & 0 \end{bmatrix}, v_3 = \begin{bmatrix} 0 & 1 \\ 2 & 0 \end{bmatrix}, v_4 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right).$$

523 Problem Let $(a, b) \in \mathbb{R}^3 \times \mathbb{R}^3$ be fixed. Solve the equation

$$a \times x = b,$$

for x .

525 Definition An *ordered basis* $\{v_1, v_2, \dots, v_n\}$ of a vector space V is a basis where the order of the v_k has been fixed. Given an ordered basis $\{v_1, v_2, \dots, v_n\}$ of a vector space V , Theorem 524 ensures that there are unique $(a_1, a_2, \dots, a_n) \in \mathbb{F}^n$ such that

$$v = a_1 v_1 + a_2 v_2 + \cdots + a_n v_n.$$

The a_k 's are called the *coordinates* of the vector v .

526 Example The standard ordered basis for \mathbb{R}^3 is $\mathcal{S} = \{i, j, k\}$. The vector $\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \in \mathbb{R}^3$ for example, has coordinates $(1, 2, 3)_{\mathcal{S}}$.

If the order of the basis were changed to the ordered basis $\mathcal{S}_1 = \{i, k, j\}$, then $\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \in \mathbb{R}^3$ would have coordinates $(1, 3, 2)_{\mathcal{S}_1}$.



Usually, when we give a coordinate representation for a vector $v \in \mathbb{R}^n$, we assume that we are using the standard basis.

527 Example Consider the vector $\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \in \mathbb{R}^3$ (given in standard representation). Since

$$\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = -1 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} - 1 \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + 3 \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix},$$

under the ordered basis $\mathcal{B}_1 = \left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right\}$, $\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$ has coordinates $(-1, -1, 3)_{\mathcal{B}_1}$. We write

$$\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = \begin{bmatrix} -1 \\ -1 \\ 3 \end{bmatrix}_{\mathcal{B}_1}.$$

528 Example The vectors of

$$\mathcal{B}_1 = \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \end{bmatrix} \right\}$$

are non-parallel, and so form a basis for \mathbb{R}^2 . So do the vectors

$$\mathcal{B}_2 = \left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\}.$$

Find the coordinates of $\begin{bmatrix} 3 \\ 4 \end{bmatrix}_{\mathcal{B}_1}$ in the base \mathcal{B}_2 .

Solution: We are seeking x, y such that

$$3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 4 \begin{bmatrix} 1 \\ 2 \end{bmatrix} = x \begin{bmatrix} 2 \\ 1 \end{bmatrix} + y \begin{bmatrix} 1 \\ -1 \end{bmatrix} \implies \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 3 \\ 4 \end{bmatrix} = \begin{bmatrix} 2 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}_{\mathcal{B}_2}.$$

Thus

$$\begin{aligned} \begin{bmatrix} x \\ y \end{bmatrix}_{\mathcal{B}_2} &= \begin{bmatrix} 2 & 1 \\ 1 & -1 \end{bmatrix}^{-1} \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 3 \\ 4 \end{bmatrix} \\ &= \begin{bmatrix} \frac{1}{3} & \frac{1}{3} \\ \frac{1}{3} & -\frac{2}{3} \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 3 \\ 4 \end{bmatrix} \\ &= \begin{bmatrix} \frac{2}{3} & 1 \\ -\frac{1}{3} & -1 \end{bmatrix} \begin{bmatrix} 3 \\ 4 \end{bmatrix} \\ &= \begin{bmatrix} 6 \\ -5 \end{bmatrix}_{\mathcal{B}_2}. \end{aligned}$$

Let us check by expressing both vectors in the standard basis of \mathbb{R}^2 :

$$\begin{bmatrix} 3 \\ 4 \end{bmatrix}_{\mathcal{B}_1} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 4 \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 7 \\ 11 \end{bmatrix},$$

$$\begin{bmatrix} 6 \\ -5 \end{bmatrix}_{\mathcal{B}_2} = 6 \begin{bmatrix} 2 \\ 1 \end{bmatrix} - 5 \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 7 \\ 11 \end{bmatrix}.$$

In general let us consider bases $\mathcal{B}_1, \mathcal{B}_2$ for the same vector space V . We want to convert $X_{\mathcal{B}_1}$ to $Y_{\mathcal{B}_2}$. We let A be the matrix formed with the column vectors of \mathcal{B}_1 in the given order and B be the matrix formed with the column vectors of \mathcal{B}_2 in the given order. Both A and B are invertible matrices since the \mathcal{B} 's are bases, in view of Theorem 514. Then we must have

$$AX_{\mathcal{B}_1} = BY_{\mathcal{B}_2} \implies Y_{\mathcal{B}_2} = B^{-1}AX_{\mathcal{B}_1}.$$

Also,

$$X_{\mathcal{B}_1} = A^{-1}BY_{\mathcal{B}_2}.$$

This prompts the following definition.

529 Definition Let $\mathcal{B}_1 = \{u_1, u_2, \dots, u_n\}$ and $\mathcal{B}_2 = \{v_1, v_2, \dots, v_n\}$ be two ordered bases for a vector space V . Let $A \in M_n(\mathbb{F})$ be the matrix having the u 's as its columns and let $B \in M_n(\mathbb{F})$ be the matrix having the v 's as its columns. The matrix $P = B^{-1}A$ is called the *transition matrix* from \mathcal{B}_1 to \mathcal{B}_2 and the matrix $P^{-1} = A^{-1}B$ is called the *transition matrix* from \mathcal{B}_2 to \mathcal{B}_1 .

530 Example Consider the bases of \mathbb{R}^3

$$\mathcal{B}_1 = \left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right\},$$

$$\mathcal{B}_2 = \left\{ \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ 0 \end{bmatrix} \right\}.$$

Find the transition matrix from \mathcal{B}_1 to \mathcal{B}_2 and also the transition matrix from \mathcal{B}_2 to \mathcal{B}_1 . Also find the coordinates

of $\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}_{\mathcal{B}_1}$ in terms of \mathcal{B}_2 .

Solution: Let

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 1 & 2 \\ 1 & -1 & 0 \\ -1 & 0 & 0 \end{bmatrix}.$$

The transition matrix from \mathcal{B}_1 to \mathcal{B}_2 is

$$\begin{aligned} \mathbf{P} &= \mathbf{B}^{-1}\mathbf{A} \\ &= \begin{bmatrix} 1 & 1 & 2 \\ 1 & -1 & 0 \\ -1 & 0 & 0 \end{bmatrix}^{-1} \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 0 & 0 & -1 \\ 0 & -1 & -1 \\ \frac{1}{2} & \frac{1}{2} & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} \\ &= \begin{bmatrix} -1 & 0 & 0 \\ -2 & -1 & -0 \\ 2 & 1 & \frac{1}{2} \end{bmatrix}. \end{aligned}$$

The transition matrix from \mathcal{B}_2 to \mathcal{B}_1 is

$$\mathbf{P}^{-1} = \begin{bmatrix} -1 & 0 & 0 \\ -2 & -1 & 0 \\ 2 & 1 & \frac{1}{2} \end{bmatrix}^{-1} = \begin{bmatrix} -1 & 0 & 0 \\ 2 & -1 & 0 \\ 0 & 2 & 2 \end{bmatrix}.$$

Now,

$$\mathbf{Y}_{\mathcal{B}_2} = \begin{bmatrix} -1 & 0 & 0 \\ -2 & -1 & 0 \\ 2 & 1 & \frac{1}{2} \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}_{\mathcal{B}_1} = \begin{bmatrix} -1 \\ -4 \\ \frac{11}{2} \end{bmatrix}_{\mathcal{B}_2}.$$

As a check, observe that in the standard basis for \mathbb{R}^3

$$\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}_{\mathcal{B}_1} = 1 \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + 2 \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + 3 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 6 \\ 3 \\ 1 \end{bmatrix},$$

$$\begin{bmatrix} -1 \\ -4 \\ \frac{11}{2} \end{bmatrix}_{\mathcal{B}_2} = -1 \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix} - 4 \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + \frac{11}{2} \begin{bmatrix} 2 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 6 \\ 3 \\ 1 \end{bmatrix}.$$

531 Problem 1. Prove that the following vectors are linearly independent in \mathbb{R}^4

$$\mathbf{a}_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \quad \mathbf{a}_2 = \begin{bmatrix} 1 \\ 1 \\ -1 \\ -1 \end{bmatrix}, \quad \mathbf{a}_3 = \begin{bmatrix} 1 \\ -1 \\ 1 \\ -1 \end{bmatrix}, \quad \mathbf{a}_4 = \begin{bmatrix} 1 \\ -1 \\ -1 \\ 1 \end{bmatrix}.$$

2. Find the coordinates of $\begin{bmatrix} 1 \\ 2 \\ 1 \\ 1 \end{bmatrix}$ under the ordered basis $(\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3, \mathbf{a}_4)$.

3. Find the coordinates of $\begin{bmatrix} 1 \\ 2 \\ 1 \\ 1 \end{bmatrix}$ under the ordered basis $(\mathbf{a}_1, \mathbf{a}_3, \mathbf{a}_2, \mathbf{a}_4)$.

532 Problem Consider the matrix

$$\mathbf{A}(\alpha) = \begin{bmatrix} \alpha & 1 & 1 & 1 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & \alpha & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix}.$$

- ① Determine all α for which $\mathbf{A}(\alpha)$ is not invertible.
- ② Find the inverse of $\mathbf{A}(\alpha)$ when $\mathbf{A}(\alpha)$ is invertible.
- ③ Find the transition matrix from the basis

$$\mathcal{B}_1 = \left[\begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right].$$

to the basis

$$\mathcal{B}_2 = \left[\begin{bmatrix} \alpha \\ 0 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ \alpha \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} \right].$$

Chapter 6

Linear Transformations

6.1 Linear Transformations

533 Definition Let $\langle V, +, \cdot, \mathbb{F} \rangle$ and $\langle W, +, \cdot, \mathbb{F} \rangle$ be vector spaces over the same field \mathbb{F} . A *linear transformation* or *homomorphism*

$$\begin{aligned} L : \quad &V \rightarrow W \\ &a \mapsto L(a) \end{aligned},$$

is a function which is

- **Linear:** $L(a + b) = L(a) + L(b)$,
- **Homogeneous:** $L(\alpha a) = \alpha L(a)$, for $\alpha \in \mathbb{F}$.

 It is clear that the above two conditions can be summarised conveniently into

$$L(a + \alpha b) = L(a) + \alpha L(b).$$

534 Example Let

$$\begin{aligned} L : \quad &M_n(\mathbb{R}) \rightarrow \mathbb{R} \\ &A \mapsto \text{tr}(A) \end{aligned}.$$

Then L is linear, for if $(A, B) \in M_n(\mathbb{R})$, then

$$L(A + \alpha B) = \text{tr}(A + \alpha B) = \text{tr}(A) + \alpha \text{tr}(B) = L(A) + \alpha L(B).$$

535 Example Let

$$\begin{aligned} L : \quad &M_n(\mathbb{R}) \rightarrow M_n(\mathbb{R}) \\ &A \mapsto A^T \end{aligned}.$$

Then L is linear, for if $(A, B) \in M_n(\mathbb{R})$, then

$$L(A + \alpha B) = (A + \alpha B)^T = A^T + \alpha B^T = L(A) + \alpha L(B).$$

536 Example For a point $(x, y) \in \mathbb{R}^2$, its reflexion about the y -axis is $(-x, y)$. Prove that

$$\begin{aligned}\mathbf{R} : \quad & \mathbb{R}^2 \rightarrow \mathbb{R}^2 \\ & (x, y) \mapsto (-x, y)\end{aligned}$$

is linear.

Solution: Let $(x_1, y_1) \in \mathbb{R}^2$, $(x_2, y_2) \in \mathbb{R}^2$, and $\alpha \in \mathbb{R}$. Then

$$\begin{aligned}\mathbf{R}((x_1, y_1) + \alpha(x_2, y_2)) &= \mathbf{R}((x_1 + \alpha x_2, y_1 + \alpha y_2)) \\ &= ((-x_1 - \alpha x_2), y_1 + \alpha y_2) \\ &= (-x_1, y_1) + \alpha(-x_2, y_2) \\ &= \mathbf{R}((x_1, y_1)) + \alpha\mathbf{R}((x_2, y_2)),\end{aligned}$$

whence \mathbf{R} is linear.

537 Example Let $\mathbf{L} : \mathbb{R}^2 \rightarrow \mathbb{R}^4$ be a linear transformation with

$$\mathbf{L} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 3 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \\ 2 \\ 3 \end{bmatrix}; \quad \mathbf{L} \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 0 \\ 2 \\ 3 \end{bmatrix}.$$

$$\text{Find } \mathbf{L} \begin{bmatrix} 5 \\ 3 \end{bmatrix}.$$

Solution: Since

$$\begin{bmatrix} 5 \\ 3 \end{bmatrix} = 4 \begin{bmatrix} 1 \\ 1 \end{bmatrix} - \begin{bmatrix} -1 \\ 1 \end{bmatrix},$$

we have

$$\mathbf{L} \begin{bmatrix} 5 \\ 3 \end{bmatrix} = 4\mathbf{L} \begin{bmatrix} 1 \\ 1 \end{bmatrix} - \mathbf{L} \begin{bmatrix} -1 \\ 1 \end{bmatrix} = 4 \begin{bmatrix} -1 \\ 1 \\ 2 \\ 3 \end{bmatrix} - \begin{bmatrix} 2 \\ 0 \\ 2 \\ 3 \end{bmatrix} = \begin{bmatrix} -6 \\ 4 \\ 6 \\ 9 \end{bmatrix}.$$

538 Theorem Let $\langle V, +, \cdot, \mathbb{F} \rangle$ and $\langle W, +, \cdot, \mathbb{F} \rangle$ be vector spaces over the same field \mathbb{F} , and let $\mathbf{L} : V \rightarrow W$ be a linear transformation. Then

- $\mathbf{L}(\mathbf{0}_V) = \mathbf{0}_W$.
- $\forall \mathbf{x} \in V, \mathbf{L}(-\mathbf{x}) = -\mathbf{L}(\mathbf{x})$.

Proof: We have

$$\mathbf{L}(\mathbf{0}_V) = \mathbf{L}(\mathbf{0}_V + \mathbf{0}_V) = \mathbf{L}(\mathbf{0}_V) + \mathbf{L}(\mathbf{0}_V),$$

hence

$$\mathbf{L}(\mathbf{0}_V) - \mathbf{L}(\mathbf{0}_V) = \mathbf{L}(\mathbf{0}_V).$$

Since

$$\mathbf{L}(\mathbf{0}_V) - \mathbf{L}(\mathbf{0}_V) = \mathbf{0}_W,$$

we obtain the first result.

Now

$$\mathbf{0}_W = \mathbf{L}(\mathbf{0}_V) = \mathbf{L}(\mathbf{x} + (-\mathbf{x})) = \mathbf{L}(\mathbf{x}) + \mathbf{L}(-\mathbf{x}),$$

from where the second result follows. \square

539 Problem Consider $\mathbf{L} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$,

$$\mathbf{L} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} x - y - z \\ x + y + z \\ z \end{bmatrix}.$$

Prove that \mathbf{L} is linear.

540 Problem Let \mathbf{h}, \mathbf{k} be fixed vectors in \mathbb{R}^3 . Prove that

$$\begin{aligned} \mathbf{L} : \quad \mathbb{R}^3 \times \mathbb{R}^3 &\rightarrow \mathbb{R}^3 \\ (\mathbf{x}, \mathbf{y}) &\mapsto \mathbf{x} \times \mathbf{k} + \mathbf{h} \times \mathbf{y} \end{aligned}$$

is a linear transformation.

541 Problem Let $\mathbf{A} \in \mathrm{GL}_n(\mathbb{R})$ be a fixed matrix. Prove that

$$\begin{aligned} \mathbf{L} : \quad \mathrm{GL}_n(\mathbb{R}) &\rightarrow \mathrm{GL}_n(\mathbb{R}) \\ \mathbf{H} &\mapsto -\mathbf{A}^{-1}\mathbf{H}\mathbf{A}^{-1} \end{aligned}$$

is a linear transformation.

542 Problem Let V be a vector space and let $S \subseteq V$. The set S is said to be *convex* if $\forall \alpha \in [0; 1], \forall \mathbf{x}, \mathbf{y} \in S$, $(1 - \alpha)\mathbf{x} + \alpha\mathbf{y} \in S$, that is, for any two points in S , the straight line joining them also belongs to S . Let $T : V \rightarrow W$ be a linear transformation from the vector space V to the vector space W . Prove that T maps convex sets into convex sets.

6.2 Kernel and Image of a Linear Transformation

543 Definition Let $\langle V, +, \cdot, \mathbb{F} \rangle$ and $\langle W, +, \cdot, \mathbb{F} \rangle$ be vector spaces over the same field \mathbb{F} , and

$$\begin{aligned} T : \quad V &\rightarrow W \\ v &\mapsto T(v) \end{aligned}$$

be a linear transformation. The *kernel* of T is the set

$$\ker(T) = \{v \in V : T(v) = \mathbf{0}_W\}.$$

The *image* of T is the set

$$\mathrm{Im}(T) = \{w \in W : \exists v \in V \text{ such that } T(v) = w\} = T(V).$$



Since $T(\mathbf{0}_V) = \mathbf{0}_W$ by Theorem 538, we have $\mathbf{0}_V \in \ker(T)$ and $\mathbf{0}_W \in \mathrm{Im}(T)$.

544 Theorem Let $\langle V, +, \cdot, \mathbb{F} \rangle$ and $\langle W, +, \cdot, \mathbb{F} \rangle$ be vector spaces over the same field \mathbb{F} , and

$$\begin{aligned} T : \quad &V \rightarrow W \\ &\mathbf{v} \mapsto T(\mathbf{v}) \end{aligned}$$

be a linear transformation. Then $\ker(T)$ is a vector subspace of V and $\text{Im}(T)$ is a vector subspace of W .

Proof: Let $(\mathbf{v}_1, \mathbf{v}_2) \in (\ker(T))^2$ and $\alpha \in \mathbb{F}$. Then $T(\mathbf{v}_1) = T(\mathbf{v}_2) = \mathbf{0}_V$. We must prove that $\mathbf{v}_1 + \alpha\mathbf{v}_2 \in \ker(T)$, that is, that $T(\mathbf{v}_1 + \alpha\mathbf{v}_2) = \mathbf{0}_W$. But

$$T(\mathbf{v}_1 + \alpha\mathbf{v}_2) = T(\mathbf{v}_1) + \alpha T(\mathbf{v}_2) = \mathbf{0}_V + \alpha\mathbf{0}_V = \mathbf{0}_V$$

proving that $\ker(T)$ is a subspace of V .

Now, let $(\mathbf{w}_1, \mathbf{w}_2) \in (\text{Im}(T))^2$ and $\alpha \in \mathbb{F}$. Then $\exists (\mathbf{v}_1, \mathbf{v}_2) \in V^2$ such that $T(\mathbf{v}_1) = \mathbf{w}_1$ and $T(\mathbf{v}_2) = \mathbf{w}_2$. We must prove that $\mathbf{w}_1 + \alpha\mathbf{w}_2 \in \text{Im}(T)$, that is, that $\exists \mathbf{v}$ such that $T(\mathbf{v}) = \mathbf{w}_1 + \alpha\mathbf{w}_2$. But

$$\mathbf{w}_1 + \alpha\mathbf{w}_2 = T(\mathbf{v}_1) + \alpha T(\mathbf{v}_2) = T(\mathbf{v}_1 + \alpha\mathbf{v}_2),$$

and so we may take $\mathbf{v} = \mathbf{v}_1 + \alpha\mathbf{v}_2$. This proves that $\text{Im}(T)$ is a subspace of W .

□

545 Theorem Let $\langle V, +, \cdot, \mathbb{F} \rangle$ and $\langle W, +, \cdot, \mathbb{F} \rangle$ be vector spaces over the same field \mathbb{F} , and

$$\begin{aligned} T : \quad &V \rightarrow W \\ &\mathbf{v} \mapsto T(\mathbf{v}) \end{aligned}$$

be a linear transformation. Then T is injective if and only if $\ker(T) = \mathbf{0}_V$.

Proof: Assume that T is injective. Then there is a unique $\mathbf{x} \in V$ mapping to $\mathbf{0}_W$:

$$T(\mathbf{x}) = \mathbf{0}_W.$$

By Theorem 538, $T(\mathbf{0}_V) = \mathbf{0}_W$, i.e., a linear transformation takes the zero vector of one space to the zero vector of the target space, and so we must have $\mathbf{x} = \mathbf{0}_V$.

Conversely, assume that $\ker(T) = \{\mathbf{0}_V\}$, and that $T(\mathbf{x}) = T(\mathbf{y})$. We must prove that $\mathbf{x} = \mathbf{y}$. But

$$\begin{aligned} T(\mathbf{x}) = T(\mathbf{y}) &\implies T(\mathbf{x}) - T(\mathbf{y}) = \mathbf{0}_W \\ &\implies T(\mathbf{x} - \mathbf{y}) = \mathbf{0}_W \\ &\implies (\mathbf{x} - \mathbf{y}) \in \ker(T) \\ &\implies \mathbf{x} - \mathbf{y} = \mathbf{0}_V \\ &\implies \mathbf{x} = \mathbf{y}, \end{aligned}$$

as we wanted to show. □

546 Theorem (Dimension Theorem) Let $\langle V, +, \cdot, \mathbb{F} \rangle$ and $\langle W, +, \cdot, \mathbb{F} \rangle$ be vector spaces of finite dimension over the same field \mathbb{F} , and

$$\begin{aligned} T : \quad & V \rightarrow W \\ & v \mapsto T(v) \end{aligned}$$

be a linear transformation. Then

$$\dim \ker(T) + \dim \text{Im}(T) = \dim V.$$

Proof: Let $\{v_1, v_2, \dots, v_k\}$ be a basis for $\ker(T)$. By virtue of Theorem 511, we may extend this to a basis $\mathcal{A} = \{v_1, v_2, \dots, v_k, v_{k+1}, v_{k+2}, \dots, v_n\}$ of V . Here $n = \dim V$. We will now show that $\mathcal{B} = \{T(v_{k+1}), T(v_{k+2}), \dots, T(v_n)\}$ is a basis for $\text{Im}(T)$. We prove that (i) \mathcal{B} spans $\text{Im}(T)$, and (ii) \mathcal{B} is a linearly independent family.

Let $w \in \text{Im}(T)$. Then $\exists v \in V$ such that $T(v) = w$. Now since \mathcal{A} is a basis for V we can write

$$v = \sum_{i=1}^n \alpha_i v_i.$$

Hence

$$w = T(v) = \sum_{i=1}^n \alpha_i T(v_i) = \sum_{i=k+1}^n \alpha_i T(v_i),$$

since $T(v_i) = 0_V$ for $1 \leq i \leq k$. Thus \mathcal{B} spans $\text{Im}(T)$.

To prove the linear independence of the \mathcal{B} assume that

$$0_W = \sum_{i=k+1}^n \beta_i T(v_i) = T \left(\sum_{i=k+1}^n \beta_i v_i \right).$$

This means that $\sum_{i=k+1}^n \beta_i v_i \in \ker(T)$, which is impossible unless $\beta_{k+1} = \beta_{k+2} = \dots = \beta_n = 0_{\mathbb{F}}$.

□

547 Corollary If $\dim V = \dim W < +\infty$, then T is injective if and only if it is surjective.

Proof: Simply observe that if T is injective then $\dim \ker(T) = 0$, and if T is surjective $\text{Im}(T) = T(V) = W$ and $\dim \text{Im}(T) = \dim W$. □

548 Example Let

$$\begin{aligned} L : \quad & M_2(\mathbb{R}) \rightarrow M_2(\mathbb{R}) \\ & A \mapsto A^T - A \end{aligned}$$

Observe that L is linear. Determine $\ker(L)$ and $\text{Im}(L)$.

Solution: Put $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ and assume $L(A) = 0_2$. Then

$$\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = L(A) = \begin{bmatrix} a & c \\ b & d \end{bmatrix} - \begin{bmatrix} a & b \\ c & d \end{bmatrix} = (c - b) \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}.$$

This means that $\mathbf{c} = \mathbf{b}$. Thus

$$\ker(L) = \left\{ \begin{bmatrix} \mathbf{a} & \mathbf{b} \\ \mathbf{b} & \mathbf{d} \end{bmatrix} : (\mathbf{a}, \mathbf{b}, \mathbf{d}) \in \mathbb{R}^3 \right\},$$

$$\text{Im}(L) = \left\{ \begin{bmatrix} 0 & k \\ -k & 0 \end{bmatrix} : k \in \mathbb{R} \right\}.$$

This means that $\dim \ker(L) = 3$, and so $\dim \text{Im}(L) = 4 - 3 = 1$.

549 Example Consider the linear transformation $L : M_2(\mathbb{R}) \rightarrow \mathbb{R}_3[X]$ given by

$$L \begin{bmatrix} \mathbf{a} & \mathbf{b} \\ \mathbf{c} & \mathbf{d} \end{bmatrix} = (\mathbf{a} + \mathbf{b})X^2 + (\mathbf{a} - \mathbf{b})X^3.$$

Determine $\ker(L)$ and $\text{Im}(L)$.

Solution: We have

$$0 = L \begin{bmatrix} \mathbf{a} & \mathbf{b} \\ \mathbf{c} & \mathbf{d} \end{bmatrix} = (\mathbf{a} + \mathbf{b})X^2 + (\mathbf{a} - \mathbf{b})X^3 \implies \mathbf{a} + \mathbf{b} = 0, \mathbf{a} - \mathbf{b} = 0, \implies \mathbf{a} = \mathbf{b} = 0.$$

Thus

$$\ker(L) = \left\{ \begin{bmatrix} 0 & 0 \\ \mathbf{c} & \mathbf{d} \end{bmatrix} : (\mathbf{c}, \mathbf{d}) \in \mathbb{R}^2 \right\}.$$

Thus $\dim \ker(L) = 2$ and hence $\dim \text{Im}(L) = 2$. Now

$$(\mathbf{a} + \mathbf{b})X^2 + (\mathbf{a} - \mathbf{b})X^3 \implies \mathbf{a}(X^2 + X^3) + \mathbf{b}(X^2 - X^3).$$

Clearly $X^2 + X^3$, and $X^2 - X^3$ are linearly independent and span $\text{Im}(L)$. Thus

$$\text{Im}(L) = \text{span}(X^2 + X^3, X^2 - X^3).$$

550 Problem In problem 539 we saw that $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$,

$$L \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} x - y - z \\ x + y + z \\ z \end{bmatrix}$$

is linear. Determine $\ker(L)$ and $\text{Im}(L)$.

satisfy

$$L \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}; \quad L \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \\ 0 \end{bmatrix}; \quad L \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}.$$

551 Problem Let

$$L : \begin{array}{ccc} \mathbb{R}^3 & \rightarrow & \mathbb{R}^4 \\ \mathbf{a} & \mapsto & L(\mathbf{a}) \end{array}$$

Determine $\ker(L)$ and $\text{Im}(L)$.

552 Problem It is easy to see that $L : \mathbb{R}^2 \rightarrow \mathbb{R}^3$,

$$L \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x + 2y \\ x + 2y \\ 0 \end{bmatrix}$$

is linear. Determine $\ker(L)$ and $\text{Im}(L)$.

553 Problem It is easy to see that $L : \mathbb{R}^2 \rightarrow \mathbb{R}^3$,

$$L \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x - y \\ x + y \\ 0 \end{bmatrix}$$

is linear. Determine $\ker(T)$ and $\text{Im}(T)$.

554 Problem It is easy to see that $L : \mathbb{R}^3 \rightarrow \mathbb{R}^2$,

$$L \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} x - y - z \\ y - 2z \end{bmatrix}$$

is linear. Determine $\ker(L)$ and $\text{Im}(L)$.

555 Problem Let

$$\begin{aligned} L : \quad M_2(\mathbb{R}) &\rightarrow \quad \mathbb{R} \\ A &\mapsto \text{tr}(A) \end{aligned} .$$

Determine $\ker(L)$ and $\text{Im}(L)$.

556 Problem 1. Demonstrate that

$$\begin{aligned} L : \quad M_2(\mathbb{R}) &\rightarrow \quad M_2(\mathbb{R}) \\ A &\mapsto A^T + A \end{aligned}$$

is a linear transformation.

2. Find a basis for $\ker(L)$ and find $\dim \ker(L)$.
3. Find a basis for $\text{Im}(L)$ and find $\dim \text{Im}(L)$.

557 Problem Let V be an n -dimensional vector space, where the characteristic of the underlying field is different from 2. A linear transformation $T : V \rightarrow V$ is *idempotent* if $T^2 = T$. Prove that if T is idempotent, then

- ① $I - T$ is idempotent (I is the identity function).
- ② $I + T$ is invertible.
- ③ $\ker(T) = \text{Im}(I - T)$

6.3 Matrix Representation

Let V, W be two vector spaces over the same field \mathbb{F} . Assume that $\dim V = m$ and $\{v_i\}_{i \in [1; m]}$ is an ordered basis for V , and that $\dim W = n$ and $\mathcal{A} = \{w_i\}_{i \in [1; n]}$ an ordered basis for W . Then

$$\begin{aligned}
 L(v_1) &= a_{11}w_1 + a_{21}w_2 + \cdots + a_{n1}w_n = \begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{n1} \end{bmatrix}_{\mathcal{A}} \\
 L(v_2) &= a_{12}w_1 + a_{22}w_2 + \cdots + a_{n2}w_n = \begin{bmatrix} a_{12} \\ a_{22} \\ \vdots \\ a_{n2} \end{bmatrix}_{\mathcal{A}} \\
 &\vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \\
 L(v_m) &= a_{1m}w_1 + a_{2m}w_2 + \cdots + a_{nm}w_n = \begin{bmatrix} a_{1n} \\ a_{2n} \\ \vdots \\ a_{nm} \end{bmatrix}_{\mathcal{A}}
 \end{aligned}$$

558 Definition The $n \times m$ matrix

$$M_L = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{12} & \cdots & a_{2n} \\ \vdots & \vdots & \vdots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nm} \end{bmatrix}$$

formed by the column vectors above is called the *matrix representation of the linear map L with respect to the bases $\{v_i\}_{i \in [1; m]}, \{w_i\}_{i \in [1; n]}$* .

559 Example Consider $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$,

$$L \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} x - y - z \\ x + y + z \\ z \end{bmatrix}.$$

Clearly L is a linear transformation.

- Find the matrix corresponding to L under the standard ordered basis.

2. Find the matrix corresponding to \mathbf{L} under the ordered basis $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$, for both the domain and the image of \mathbf{L} .

Solution:

1. Solution: The matrix will be a 3×3 matrix. We have $\mathbf{L} \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$, $\mathbf{L} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$, and $\mathbf{L} \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$,

whence the desired matrix is

$$\begin{bmatrix} 1 & -1 & -1 \\ 1 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}.$$

2. Call this basis \mathcal{A} . We have

$$\mathbf{L} \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = 0 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 1 \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + 0 \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}_{\mathcal{A}},$$

$$\mathbf{L} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \\ 0 \end{bmatrix} = -2 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 2 \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + 0 \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} -2 \\ 2 \\ 0 \end{bmatrix}_{\mathcal{A}},$$

and

$$\mathbf{L} \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \\ 1 \end{bmatrix} = -3 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 2 \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + 1 \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} -3 \\ 2 \\ 1 \end{bmatrix}_{\mathcal{A}},$$

whence the desired matrix is

$$\begin{bmatrix} 0 & -2 & -3 \\ 1 & 2 & 2 \\ 0 & 0 & 1 \end{bmatrix}.$$

560 Example Let $\mathbb{R}_n[x]$ denote the set of polynomials with real coefficients with degree at most n .

1. Prove that

$$\begin{aligned} \mathbf{L} : \quad \mathbb{R}_3[x] &\rightarrow \mathbb{R}_1[x] \\ p(x) &\mapsto p''(x) \end{aligned}$$

is a linear transformation. Here $p''(x)$ denotes the second derivative of $p(x)$ with respect to x .

2. Find the matrix of \mathbf{L} using the ordered bases $\{1, x, x^2, x^3\}$ for $\mathbb{R}_3[x]$ and $\{1, x\}$ for $\mathbb{R}_2[x]$.
3. Find the matrix of \mathbf{L} using the ordered bases $\{1, x, x^2, x^3\}$ for $\mathbb{R}_3[x]$ and $\{1, x + 2\}$ for $\mathbb{R}_1[x]$.
4. Find a basis for $\ker(\mathbf{L})$ and find $\dim \ker(\mathbf{L})$.
5. Find a basis for $\text{Im } (\mathbf{L})$ and find $\dim \text{Im } (\mathbf{L})$.

Solution:

1. Let $(p(x), q(x)) \in (\mathbb{R}_3[x])^2$ and $\alpha \in \mathbb{R}$. Then

$$\mathbf{L}(p(x) + \alpha q(x)) = (p(x) + \alpha q(x))'' = p''(x) + \alpha q''(x) = \mathbf{L}(p(x)) + \alpha \mathbf{L}(q(x)),$$

whence \mathbf{L} is linear.

2. We have

$$\begin{aligned} \mathbf{L}(1) &= \frac{d^2}{dx^2} 1 = 0 = 0(1) + 0(x) = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \\ \mathbf{L}(x) &= \frac{d^2}{dx^2} x = 0 = 0(1) + 0(x) = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \\ \mathbf{L}(x^2) &= \frac{d^2}{dx^2} x^2 = 2 = 2(1) + 0(x) = \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \\ \mathbf{L}(x^3) &= \frac{d^2}{dx^2} x^3 = 6x = 0(1) + 6(x) = \begin{bmatrix} 0 \\ 6 \end{bmatrix}, \end{aligned}$$

whence the matrix representation of \mathbf{L} under the standard basis is

$$\begin{bmatrix} 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 6 \end{bmatrix}.$$

3. We have

$$\begin{aligned} L(1) &= \frac{d^2}{dx^2}1 = 0 = 0(1) + 0(x+2) = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \\ L(x) &= \frac{d^2}{dx^2}x = 0 = 0(1) + 0(x+2) = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \\ L(x^2) &= \frac{d^2}{dx^2}x^2 = 2 = 2(1) + 0(x+2) = \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \\ L(x^3) &= \frac{d^2}{dx^2}x^3 = 6x = -12(1) + 6(x+2) = \begin{bmatrix} -12 \\ 6 \end{bmatrix}, \end{aligned}$$

whence the matrix representation of L under the standard basis is

$$\begin{bmatrix} 0 & 0 & 2 & -12 \\ 0 & 0 & 0 & 6 \end{bmatrix}.$$

4. Assume that $p(x) = a + bx + cx^2 + dx^3 \in \ker(L)$. Then

$$0 = L(p(x)) = 2c + 6dx, \quad \forall x \in \mathbb{R}.$$

This means that $c = d = 0$. Thus a, b are free and

$$\ker(L) = \{a + bx : (a, b) \in \mathbb{R}^2\}.$$

Hence $\dim \ker(L) = 2$.

5. By the Dimension Theorem, $\dim \text{Im}(L) = 4 - 2 = 2$. Put $q(x) = \alpha + \beta x + \gamma x^2 + \delta x^3$. Then

$$L(q(x)) = 2\gamma + 6\delta(x) = (2\gamma)(1) + (6\delta)(x).$$

Clearly $\{1, x\}$ are linearly independent and $\text{span } \text{Im}(L)$. Hence

$$\text{Im}(L) = \text{span}(1, x) = \mathbb{R}_1[x].$$

561 Example 1. A linear transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is such that

$$T(i) = \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix}; \quad T(j) = \begin{bmatrix} 3 \\ 0 \\ -1 \end{bmatrix}.$$

It is known that

$$\text{Im}(T) = \text{span}(T(i), T(j))$$

and that

$$\ker(T) = \text{span} \left(\begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix} \right).$$

Argue that there must be λ and μ such that

$$T(k) = \lambda T(i) + \mu T(j).$$

2. Find λ and μ , and hence, the matrix representing T under the standard ordered basis.

Solution:

1. Since $T(k) \in \text{Im}(T)$ and $\text{Im}(T)$ is generated by $T(i)$ and $T(k)$ there must be $(\lambda, \mu) \in \mathbb{R}^2$ with

$$T(k) = \lambda T(i) + \mu T(j) = \lambda \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} + \mu \begin{bmatrix} 3 \\ 0 \\ -1 \end{bmatrix} = \begin{bmatrix} 2\lambda + 3\mu \\ \lambda \\ \lambda - \mu \end{bmatrix}.$$

2. The matrix of T is

$$\begin{bmatrix} T(i) & T(j) & T(k) \end{bmatrix} = \begin{bmatrix} 2 & 3 & 2\lambda + 3\mu \\ 1 & 0 & \lambda \\ 1 & -1 & \lambda - \mu \end{bmatrix}.$$

Since $\begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix} \in \ker(T)$ we must have

$$\begin{bmatrix} 2 & 3 & 2\lambda + 3\mu \\ 1 & 0 & \lambda \\ 1 & -1 & \lambda - \mu \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}.$$

Solving the resulting system of linear equations we obtain $\lambda = 1, \mu = 2$. The required matrix is thus

$$\begin{bmatrix} 2 & 3 & 8 \\ 1 & 0 & 1 \\ 1 & -1 & -1 \end{bmatrix}.$$

 If the linear mapping $L : V \rightarrow W$, $\dim V = n, \dim W = m$ has matrix representation $A \in M_{m \times n}(\mathbb{F})$, then $\dim \text{Im}(L) = \text{rank}(A)$.

562 Problem 1. A linear transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ has as image the plane with equation $x + y + z = 0$ and as

kernel the line $x = y = z$. If

$$T \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix} = \begin{bmatrix} a \\ 0 \\ 1 \end{bmatrix}, \quad T \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 \\ b \\ -5 \end{bmatrix}, \quad T \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \\ c \end{bmatrix}.$$

Find $\mathbf{a}, \mathbf{b}, \mathbf{c}$.

2. Find the matrix representation of \mathbf{T} under the standard basis.

563 Problem 1. Prove that $\mathbf{T} : \mathbb{R}^2 \rightarrow \mathbb{R}^3$

$$\mathbf{T} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x+y \\ x-y \\ 2x+3y \end{bmatrix}$$

is a linear transformation.

2. Find a basis for $\ker(\mathbf{T})$ and find $\dim \ker(\mathbf{T})$

3. Find a basis for $\text{Im } (\mathbf{T})$ and find $\dim \text{Im } (\mathbf{T})$.

4. Find the matrix of \mathbf{T} under the ordered bases $\mathcal{A} = \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 3 \end{bmatrix} \right\}$ of \mathbb{R}^2 and $\mathcal{B} = \left\{ \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \right\}$ of \mathbb{R}^3 .

564 Problem Let

$$\begin{aligned} \mathbf{L} : \mathbb{R}^3 &\rightarrow \mathbb{R}^2 \\ \mathbf{a} &\mapsto \mathbf{L}(\mathbf{a}) \end{aligned}$$

where

$$\mathbf{L} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} x+2y \\ 3x-z \end{bmatrix}.$$

Clearly \mathbf{L} is linear. Find a matrix representation for \mathbf{L} if

1. The bases for both \mathbb{R}^3 and \mathbb{R}^2 are both the standard ordered bases.

2. The ordered basis for \mathbb{R}^3 is $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ and \mathbb{R}^2 has the standard ordered basis.

3. The ordered basis for \mathbb{R}^3 is $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ and the ordered basis for \mathbb{R}^2 is $\mathcal{A} = \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$.

565 Problem A linear transformation $\mathbf{T} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ satisfies $\ker(\mathbf{T}) = \text{Im } (\mathbf{T})$, and $\mathbf{T} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$. Find the matrix representing \mathbf{T} under the standard ordered basis.

566 Problem Find the matrix representation for the linear map

$$\begin{aligned} \mathbf{L} : \mathbf{M}_2(\mathbb{R}) &\rightarrow \mathbb{R} \\ \mathbf{A} &\mapsto \text{tr}(\mathbf{A}) \end{aligned}$$

under the standard basis

$$\mathcal{A} = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}$$

for $\mathbf{M}_2(\mathbb{R})$.

567 Problem Let $\mathbf{A} \in \mathbf{M}_{n \times p}(\mathbb{R})$, $\mathbf{B} \in \mathbf{M}_{p \times q}(\mathbb{R})$, and $\mathbf{C} \in \mathbf{M}_{q \times r}(\mathbb{R})$, be such that $\text{rank } (\mathbf{B}) = \text{rank } (\mathbf{AB})$. Show that

$$\text{rank } (\mathbf{BC}) = \text{rank } (\mathbf{ABC}).$$

Determinants

7.1 Permutations

568 Definition Let S be a finite set with $n \geq 1$ elements. A *permutation* is a bijective function $\tau : S \rightarrow S$. It is easy to see that there are $n!$ permutations from S onto itself.

Since we are mostly concerned with the *action* that τ exerts on S rather than with the particular names of the elements of S , we will take S to be the set $S = \{1, 2, 3, \dots, n\}$. We indicate a permutation τ by means of the following convenient diagram

$$\tau = \begin{bmatrix} 1 & 2 & \dots & n \\ \tau(1) & \tau(2) & \dots & \tau(n) \end{bmatrix}.$$

569 Definition The notation S_n will denote the set of all permutations on $\{1, 2, 3, \dots, n\}$. Under this notation, the composition of two permutations $(\tau, \sigma) \in S_n^2$ is

$$\begin{aligned} \tau \circ \sigma &= \begin{bmatrix} 1 & 2 & \dots & n \\ \tau(1) & \tau(2) & \dots & \tau(n) \end{bmatrix} \circ \begin{bmatrix} 1 & 2 & \dots & n \\ \sigma(1) & \sigma(2) & \dots & \sigma(n) \end{bmatrix} \\ &= \begin{bmatrix} 1 & & 2 & & \dots & & n \\ (\tau \circ \sigma)(1) & (\tau \circ \sigma)(2) & \dots & (\tau \circ \sigma)(n) \end{bmatrix}. \end{aligned}$$

The k -fold composition of τ is

$$\underbrace{\tau \circ \dots \circ \tau}_{k \text{ compositions}} = \tau^k.$$

 We usually do away with the \circ and write $\tau \circ \sigma$ simply as $\tau\sigma$. This “product of permutations” is thus simply function composition.

Given a permutation $\tau : S \rightarrow S$, since τ is bijective,

$$\tau^{-1} : S \rightarrow S$$

exists and is also a permutation. In fact if

$$\tau = \begin{bmatrix} 1 & 2 & \dots & n \\ \tau(1) & \tau(2) & \dots & \tau(n) \end{bmatrix},$$

then

$$\tau^{-1} = \begin{bmatrix} \tau(1) & \tau(2) & \cdots & \tau(n) \\ 1 & 2 & \cdots & n \end{bmatrix}.$$




Figure 7.1: S_3 are rotations and reflexions.

570 Example The set S_3 has $3! = 6$ elements, which are given below.

1. $\text{Id} : \{1, 2, 3\} \rightarrow \{1, 2, 3\}$ with

$$\text{Id} = \begin{bmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \end{bmatrix}.$$

2. $\tau_1 : \{1, 2, 3\} \rightarrow \{1, 2, 3\}$ with

$$\tau_1 = \begin{bmatrix} 1 & 2 & 3 \\ 1 & 3 & 2 \end{bmatrix}.$$

3. $\tau_2 : \{1, 2, 3\} \rightarrow \{1, 2, 3\}$ with

$$\tau_2 = \begin{bmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{bmatrix}.$$

4. $\tau_3 : \{1, 2, 3\} \rightarrow \{1, 2, 3\}$ with

$$\tau_3 = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{bmatrix}.$$

5. $\sigma_1 : \{1, 2, 3\} \rightarrow \{1, 2, 3\}$ with

$$\sigma_1 = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{bmatrix}.$$

6. $\sigma_2 : \{1, 2, 3\} \rightarrow \{1, 2, 3\}$ with

$$\sigma_2 = \begin{bmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{bmatrix}.$$

571 Example The compositions $\tau_1 \circ \sigma_1$ and $\sigma_1 \circ \tau_1$ can be found as follows.

$$\tau_1 \circ \sigma_1 = \begin{bmatrix} 1 & 2 & 3 \\ 1 & 3 & 2 \end{bmatrix} \circ \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{bmatrix} = \tau_2.$$

(We read from right to left $1 \rightarrow 2 \rightarrow 3$ ("1 goes to 2, 2 goes to 3, so 1 goes to 3"), etc. Similarly

$$\sigma_1 \circ \tau_1 = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{bmatrix} \circ \begin{bmatrix} 1 & 2 & 3 \\ 1 & 3 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{bmatrix} = \tau_3.$$

Observe in particular that $\sigma_1 \circ \tau_1 \neq \tau_1 \circ \sigma_1$. Finding all the other products we deduce the following "multiplication table" (where the "multiplication" operation is really composition of functions).

\circ	Id	τ_1	τ_2	τ_3	σ_1	σ_2
Id	Id	τ_1	τ_2	τ_3	σ_1	σ_2
τ_1	τ_1	Id	σ_1	σ_2	τ_2	τ_3
τ_2	τ_2	σ_2	Id	σ_1	τ_3	τ_1
τ_3	τ_3	σ_1	σ_2	Id	τ_1	τ_2
σ_2	σ_2	τ_2	τ_3	τ_1	Id	σ_1
σ_1	σ_1	τ_3	τ_1	τ_2	σ_2	Id

The permutations in example 570 can be conveniently interpreted as follows. Consider an equilateral triangle with vertices labelled 1, 2 and 3, as in figure 7.1. Each τ_a is a reflexion ("flipping") about the line joining the vertex a with the midpoint of the side opposite a . For example τ_1 fixes 1 and flips 2 and 3. Observe that two successive flips return the vertices to their original position and so $(\forall a \in \{1, 2, 3\})(\tau_a^2 = \text{Id})$. Similarly, σ_1 is a rotation of the vertices by an angle of 120° . Three successive rotations restore the vertices to their original position and so $\sigma_1^3 = \text{Id}$.

572 Example To find τ_1^{-1} take the representation of τ_1 and exchange the rows:

$$\tau_1^{-1} = \begin{bmatrix} 1 & 3 & 2 \\ 1 & 2 & 3 \end{bmatrix}.$$

This is more naturally written as

$$\tau_1^{-1} = \begin{bmatrix} 1 & 2 & 3 \\ 1 & 3 & 2 \end{bmatrix}.$$

Observe that $\tau_1^{-1} = \tau_1$.

573 Example To find σ_1^{-1} take the representation of σ_1 and exchange the rows:

$$\sigma_1^{-1} = \begin{bmatrix} 2 & 3 & 1 \\ 1 & 2 & 3 \end{bmatrix}.$$

This is more naturally written as

$$\sigma_1^{-1} = \begin{bmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{bmatrix}.$$

Observe that $\sigma_1^{-1} = \sigma_2$.

7.2 Cycle Notation

We now present a shorthand notation for permutations by introducing the idea of a *cycle*. Consider in S_9 , the permutation

$$\tau = \begin{bmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ 2 & 1 & 3 & 6 & 9 & 7 & 8 & 4 & 5 \end{bmatrix}.$$

We start with 1. Since 1 goes to 2 and 2 goes back to 1, we write (12). Now we continue with 3. Since 3 goes to 3, we write (3). We continue with 4. As 4 goes 6, 6 goes to 7, 7 goes 8, and 8 goes back to 4, we write (4678). We consider now 5 which goes to 9 and 9 goes back to 5, so we write (59). We have written τ as a product of disjoint cycles

$$\tau = (12)(3)(4678)(59).$$

This prompts the following definition.

574 Definition Let $l \geq 1$ and let $i_1, \dots, i_l \in \{1, 2, \dots, n\}$ be distinct. We write $(i_1 \ i_2 \ \dots \ i_l)$ for the element $\sigma \in S_n$ such that $\sigma(i_r) = i_{r+1}$, $1 \leq r < l$, $\sigma(i_l) = i_1$ and $\sigma(i) = i$ for $i \notin \{i_1, \dots, i_l\}$. We say that $(i_1 \ i_2 \ \dots \ i_l)$ is a *cycle of length l*. The *order* of a cycle is its length. Observe that if τ has order l then $\tau^l = \text{Id}$.

 Observe that $(i_2 \ \dots \ i_l \ i_1) = (i_1 \ \dots \ i_l)$ etc., and that $(1) = (2) = \dots = (n) = \text{Id}$. In fact, we have

$$(i_1 \ \dots \ i_l) = (j_1 \ \dots \ j_m)$$

if and only if (1) $l = m$ and if (2) $l > 1$: $\exists a$ such that $\forall k: i_k = j_{k+a \bmod l}$. Two cycles (i_1, \dots, i_l) and (j_1, \dots, j_m) are disjoint if $\{i_1, \dots, i_l\} \cap \{j_1, \dots, j_m\} = \emptyset$. Disjoint cycles commute and if $\tau = \sigma_1 \sigma_2 \dots \sigma_t$ is the product of disjoint cycles of length l_1, l_2, \dots, l_t respectively, then τ has order

$$\text{lcm}(l_1, l_2, \dots, l_t).$$

575 Example A cycle decomposition for $\alpha \in S_9$,

$$\alpha = \begin{bmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ 1 & 8 & 7 & 6 & 2 & 3 & 4 & 5 & 9 \end{bmatrix}$$

is

$$(285)(3746).$$

The order of α is $\text{lcm}(3, 4) = 12$.

576 Example The cycle decomposition $\beta = (123)(567)$ in S_9 arises from the permutation

$$\beta = \begin{bmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ 2 & 3 & 1 & 4 & 6 & 7 & 5 & 8 & 9 \end{bmatrix}.$$

Its order is $\text{lcm}(3, 3) = 3$.

577 Example Find a shuffle of a deck of 13 cards that requires 42 repeats to return the cards to their original order.

Solution: Here is one (of many possible ones). Observe that $7 + 6 = 13$ and $7 \times 6 = 42$. We take the permutation

$$(1\ 2\ 3\ 4\ 5\ 6\ 7)(8\ 9\ 10\ 11\ 12\ 13)$$

which has order 42. This corresponds to the following shuffle: For

$$i \in \{1, 2, 3, 4, 5, 6, 8, 9, 10, 11, 12\},$$

take the i th card to the $(i+1)$ th place, take the 7th card to the first position and the 13th card to the 8th position.
Query: Of all possible shuffles of 13 cards, which one takes the longest to restore the cards to their original position?

578 Example Let a shuffle of a deck of 10 cards be made as follows: The top card is put at the bottom, the deck is cut in half, the bottom half is placed on top of the top half, and then the resulting bottom card is put on top. How many times must this shuffle be repeated to get the cards in the initial order? Explain.

Solution: Putting the top card at the bottom corresponds to

$$\begin{bmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 \\ 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 & 1 \end{bmatrix}.$$

Cutting this new arrangement in half and putting the lower half on top corresponds to

$$\begin{bmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 \\ 7 & 8 & 9 & 10 & 1 & 2 & 3 & 4 & 5 & 6 \end{bmatrix}.$$

Putting the bottom card of this new arrangement on top corresponds to

$$\begin{bmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 \\ 6 & 7 & 8 & 9 & 10 & 1 & 2 & 3 & 4 & 5 \end{bmatrix} = (1\ 6)(2\ 7)(3\ 8)(4\ 9)(5\ 10).$$

The order of this permutation is $\text{lcm}(2, 2, 2, 2, 2) = 2$, so in 2 shuffles the cards are restored to their original position.

The above examples illustrate the general case, given in the following theorem.

579 Theorem Every permutation in S_n can be written as a product of disjoint cycles.

Proof: Let $\tau \in S_n, a_1 \in \{1, 2, \dots, n\}$. Put $\tau^k(a_1) = a_{k+1}, k \geq 0$. Let a_1, a_2, \dots, a_s be the longest chain with no repeats. Then we have $\tau(a_s) = a_1$. If the $\{a_1, a_2, \dots, a_s\}$ exhaust $\{1, 2, \dots, n\}$ then we have $\tau = (a_1\ a_2\ \dots\ a_s)$. If not, there exist $b_1 \in \{1, 2, \dots, n\} \setminus \{a_1, a_2, \dots, a_s\}$. Again, we find the longest chain of distinct b_1, b_2, \dots, b_t such that $\tau(b_k) = b_{k+1}, k = 1, \dots, t-1$ and $\tau(b_t) = b_1$. If the $\{a_1, a_2, \dots, a_s, b_1, b_2, \dots, b_t\}$ exhaust all the

$\{1, 2, \dots, n\}$ we have $\tau = (a_1 \ a_2 \ \dots \ a_s)(b_1 \ b_2 \ \dots \ b_t)$. If not we continue the process and find

$$\tau = (a_1 \ a_2 \ \dots \ a_s)(b_1 \ b_2 \ \dots \ b_t)(c_1 \ \dots) \dots$$

This process stops because we have only n elements. \square

580 Definition A transposition is a cycle of length 2.¹

581 Example The cycle (23468) can be written as a product of transpositions as follows

$$(23468) = (28)(26)(24)(23).$$

Notice that this decomposition as the product of transpositions is not unique. Another decomposition is

$$(23468) = (23)(34)(46)(68).$$

582 Lemma Every permutation is the product of transpositions.

Proof: It is enough to observe that

$$(a_1 \ a_2 \ \dots \ a_s) = (a_1 \ a_s)(a_1 \ a_{s-1}) \cdots (a_1 \ a_2)$$

and appeal to Theorem 579. \square

Let $\sigma \in S_n$ and let $(i, j) \in \{1, 2, \dots, n\}^2$, $i \neq j$. Since σ is a permutation, $\exists (a, b) \in \{1, 2, \dots, n\}^2$, $a \neq b$, such that $\sigma(j) - \sigma(i) = b - a$. This means that

$$\left| \prod_{1 \leq i < j \leq n} \frac{\sigma(i) - \sigma(j)}{i - j} \right| = 1.$$

583 Definition Let $\sigma \in S_n$. We define the sign $\text{sgn}(\sigma)$ of σ as

$$\text{sgn}(\sigma) = \prod_{1 \leq i < j \leq n} \frac{\sigma(i) - \sigma(j)}{i - j} = (-1)^{I(\sigma)}.$$

If $\text{sgn}(\sigma) = 1$, then we say that σ is an even permutation, and if $\text{sgn}(\sigma) = -1$ we say that σ is an odd permutation.

 Notice that in fact

$$\text{sgn}(\sigma) = (-1)^{I(\sigma)},$$

where $I(\sigma) = \#\{(i, j) \mid 1 \leq i < j \leq n \text{ and } \sigma(i) > \sigma(j)\}$, i.e., $I(\sigma)$ is the number of inversions that σ effects to the identity permutation Id .

584 Example The transposition $(1 \ 2)$ has one inversion.

585 Lemma For any transposition $(k \ l)$ we have $\text{sgn}((k \ l)) = -1$.

Proof: Let τ be transposition that exchanges k and l , and assume that $k < l$:

$$\tau = \begin{bmatrix} 1 & 2 & \dots & k-1 & k & k+1 & \dots & l-1 & l & l+1 & \dots & n \\ 1 & 2 & \dots & k-1 & l & k+1 & \dots & l-1 & k & l+1 & \dots & n \end{bmatrix}$$

Let us count the number of inversions of τ :

¹A cycle of length 2 should more appropriately be called a bicycle.

- The pairs (i, j) with $i \in \{1, 2, \dots, k-1\} \cup \{l, l+1, \dots, n\}$ and $i < j$ do not suffer an inversion;
- The pair (k, j) with $k < j$ suffers an inversion if and only if $j \in \{k+1, k+2, \dots, l\}$, making $l-k$ inversions;
- If $i \in \{k+1, k+2, \dots, l-1\}$ and $i < j$, (i, j) suffers an inversion if and only if $j = l$, giving $l-1-k$ inversions.

This gives a total of $I(\tau) = (l-k) + (l-1-k) = 2(l-k-1) + 1$ inversions when $k < l$. Since this number is odd, we have $\text{sgn}(\tau) = (-1)^{I(\tau)} = -1$. In general we see that the transposition $(k \ l)$ has $2|k-l|-1$ inversions. \square

586 Theorem Let $(\sigma, \tau) \in S_n^2$. Then

$$\text{sgn}(\tau\sigma) = \text{sgn}(\tau)\text{sgn}(\sigma).$$

Proof: We have

$$\begin{aligned} \text{sgn}(\sigma\tau) &= \prod_{1 \leq i < j \leq n} \frac{(\sigma\tau)(i) - (\sigma\tau)(j)}{i-j} \\ &= \left(\prod_{1 \leq i < j \leq n} \frac{\sigma(\tau(i)) - \sigma(\tau(j))}{\tau(i) - \tau(j)} \right) \cdot \left(\prod_{1 \leq i < j \leq n} \frac{\tau(i) - \tau(j)}{i-j} \right). \end{aligned}$$

The second factor on this last equality is clearly $\text{sgn}(\tau)$, we must shew that the first factor is $\text{sgn}(\sigma)$. Observe now that for $1 \leq a < b \leq n$ we have

$$\frac{\sigma(a) - \sigma(b)}{a - b} = \frac{\sigma(b) - \sigma(a)}{b - a}.$$

Since σ and τ are permutations, $\exists b \neq a, \tau(i) = a, \tau(j) = b$ and so $\sigma\tau(i) = \sigma(a), \sigma\tau(j) = b$. Thus

$$\frac{\sigma(\tau(i)) - \sigma(\tau(j))}{\tau(i) - \tau(j)} = \frac{\sigma(a) - \sigma(b)}{a - b}$$

and so

$$\prod_{1 \leq i < j \leq n} \frac{\sigma(\tau(i)) - \sigma(\tau(j))}{\tau(i) - \tau(j)} = \prod_{1 \leq a < b \leq n} \frac{\sigma(a) - \sigma(b)}{a - b} = \text{sgn}(\sigma).$$

\square

587 Corollary The identity permutation is even. If $\tau \in S_n$, then $\text{sgn}(\tau) = \text{sgn}(\tau^{-1})$.

Proof: Since there are no inversions in Id , we have $\text{sgn}(\text{Id}) = (-1)^0 = 1$. Since $\tau\tau^{-1} = \text{Id}$, we must have $1 = \text{sgn}(\text{Id}) = \text{sgn}(\tau\tau^{-1}) = \text{sgn}(\tau)\text{sgn}(\tau^{-1}) = (-1)^{\tau}(-1)^{\tau^{-1}}$ by Theorem 586. Since the values on the righthand of this last equality are ± 1 , we must have $\text{sgn}(\tau) = \text{sgn}(\tau^{-1})$. \square

588 Lemma We have $\text{sgn}(1 \ 2 \ \dots \ l) = (-1)^{l-1}$.

Proof: Simply observe that the number of inversions of $(1 \ 2 \ \dots \ l)$ is $l-1$. \square

589 Lemma Let $(\tau, (i_1 \ \dots \ i_l)) \in S_n^2$. Then

$$\tau(i_1 \ \dots \ i_l)\tau^{-1} = (\tau(i_1) \ \dots \ \tau(i_l)),$$

and if $\sigma \in S_n$ is a cycle of length l then

$$\text{sgn}(\sigma) = (-1)^{l-1}$$

Proof: For $1 \leq k < l$ we have $(\tau(i_1 \dots i_l)\tau^{-1})(\tau(i_k)) = \tau((i_1 \dots i_l)(i_k)) = \tau(i_{k+1})$. On a $(\tau(i_1 \dots i_l)\tau^{-1})(\tau(i_l)) = \tau((i_1 \dots i_l)(i_l)) = \tau(i_1)$. For $i \notin \{\tau(i_1) \dots \tau(i_l)\}$ we have $\tau^{-1}(i) \notin \{i_1 \dots i_l\}$ whence $(i_1 \dots i_l)(\tau^{-1}(i)) = \tau^{-1}(i)$ etc.

Furthermore, write $\sigma = (i_1 \dots i_l)$. Let $\tau \in S_n$ be such that $\tau(k) = i_k$ for $1 \leq k \leq l$. Then $\sigma = \tau(1 2 \dots l)\tau^{-1}$ and so we must have $\text{sgn}(\sigma) = \text{sgn}(\tau)\text{sgn}((1 2 \dots l))\text{sgn}(\tau^{-1})$, which equals $\text{sgn}((1 2 \dots l))$ by virtue of Theorem 586 and Corollary 587. The result now follows by appealing to Lemma 588 \square

590 Corollary Let $\sigma = \sigma_1\sigma_2 \dots \sigma_r$ be a product of disjoint cycles, each of length l_1, \dots, l_r , respectively. Then

$$\text{sgn}(\sigma) = (-1)^{\sum_{i=1}^r (l_i - 1)}.$$

Hence, the product of two even permutations is even, the product of two odd permutations is even, and the product of an even permutation and an odd permutation is odd.

Proof: This follows at once from Theorem 586 and Lemma 589. \square

591 Example The cycle (4678) is an odd cycle; the cycle (1) is an even cycle; the cycle (12345) is an even cycle.

592 Corollary Every permutation can be decomposed as a product of transpositions. This decomposition is not necessarily unique, but its parity is unique.

Proof: This follows from Theorem 579, Lemma 582, and Corollary 590. \square

593 Example (The 15 puzzle) Consider a grid with 16 squares, as shewn in (7.1), where 15 squares are numbered 1 through 15 and the 16th slot is empty.

1	2	3	4
5	6	7	8
9	10	11	12
13	14	15	

(7.1)

In this grid we may successively exchange the empty slot with any of its neighbours, as for example

1	2	3	4
5	6	7	8
9	10	11	12
13	14		15

(7.2)

We ask whether through a series of valid moves we may arrive at the following position.

1	2	3	4
5	6	7	8
9	10	11	12
13	15	14	

(7.3)

Solution: Let us shew that this is impossible. Each time we move a square to the empty position, we make transpositions on the set $\{1, 2, \dots, 16\}$. Thus at each move, the permutation is multiplied by a transposition and hence it changes sign. Observe that the permutation corresponding to the square in (7.3) is $(14\ 15)$ (the positions 14th and 15th are transposed) and hence it is an odd permutation. But we claim that the empty slot can only return to its original position after an even permutation. To see this paint the grid as a checkerboard:

B	R	B	R
R	B	R	B
B	R	B	R
R	B	R	B

(7.4)

We see that after each move, the empty square changes from black to red, and thus after an odd number of moves the empty slot is on a red square. Thus the empty slot cannot return to its original position in an odd number of moves. This completes the proof.

594 Problem Decompose the permutation

$$\begin{bmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ 2 & 3 & 4 & 1 & 5 & 8 & 6 & 7 & 9 \end{bmatrix}$$

as a product of disjoint cycles and find its order.

7.3 Determinants

There are many ways of developing the theory of determinants. We will choose a way that will allow us to deduce the properties of determinants with ease, but has the drawback of being computationally cumbersome. In the next section we will shew that our way of defining determinants is equivalent to a more computationally friendly one.

It may be pertinent here to quickly review some properties of permutations. Recall that if $\sigma \in S_n$ is a cycle of length l , then its signum $\text{sgn}(\sigma) = \pm 1$ depending on the parity of $l - 1$. For example, in S_7 ,

$$\sigma = (1\ 3\ 4\ 7\ 6)$$

has length 5, and the parity of $5 - 1 = 4$ is even, and so we write $\text{sgn}(\sigma) = +1$. On the other hand,

$$\tau = (1\ 3\ 4\ 7\ 6\ 5)$$

has length 6, and the parity of $6 - 1 = 5$ is odd, and so we write $\text{sgn}(\tau) = -1$.

Recall also that if $(\sigma, \tau) \in S_n^2$, then

$$\text{sgn}(\tau\sigma) = \text{sgn}(\tau)\text{sgn}(\sigma).$$

Thus from the above two examples

$$\sigma\tau = (1\ 3\ 4\ 7\ 6)(1\ 3\ 4\ 7\ 6\ 5)$$

has signum $\text{sgn}(\sigma)\text{sgn}(\tau) = (+1)(-1) = -1$. Observe in particular that for the identity permutation $\text{Id} \in S_n$ we have $\text{sgn}(\text{Id}) = +1$.

595 Definition Let $A \in M_n(\mathbb{F})$, $A = [a_{ij}]$ be a square matrix. The *determinant of A* is defined and denoted by the sum

$$\det A = \sum_{\sigma \in S_n} \text{sgn}(\sigma)a_{1\sigma(1)}a_{2\sigma(2)} \cdots a_{n\sigma(n)}.$$

 The determinantal sum has $n!$ summands.

596 Example If $n = 1$, then S_1 has only one member, Id , where $\text{Id}(1) = 1$. Since Id is an even permutation, $\text{sgn}(\text{Id}) = (+1)$. Thus if $A = (a_{11})$, then

$$\det A = a_{11}$$

597 Example If $n = 2$, then S_2 has $2! = 2$ members, Id and $\sigma = (1\ 2)$. Observe that $\text{sgn}(\sigma) = -1$. Thus if

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$$

then

$$\det A = \text{sgn}(\text{Id})a_{11}a_{22} + \text{sgn}(\sigma)a_{12}a_{21} = a_{11}a_{22} - a_{12}a_{21}.$$

598 Example From the above formula for 2×2 matrices it follows that

$$\begin{aligned} \det A &= \det \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \\ &= (1)(4) - (3)(2) = -2, \end{aligned}$$

$$\begin{aligned} \det B &= \det \begin{bmatrix} -1 & 2 \\ 3 & 4 \end{bmatrix} (-1)(4) - (3)(2) \\ &= -10, \end{aligned}$$

and

$$\det(A + B) = \det \begin{bmatrix} 0 & 4 \\ 6 & 8 \end{bmatrix} = (0)(8) - (6)(4) = -24.$$

Observe in particular that $\det(A + B) \neq \det A + \det B$.

599 Example If $n = 3$, then S_3 has $3! = 6$ members:

$$\text{Id}, \tau_1 = (2\ 3), \tau_2 = (1\ 3), \tau_3 = (1\ 2), \sigma_1 = (1\ 2\ 3), \sigma_2 = (1\ 3\ 2).$$

Observe that $\text{Id}, \sigma_1, \sigma_2$ are even, and τ_1, τ_2, τ_3 are odd. Thus if

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

then

$$\begin{aligned}
 \det \mathbf{A} &= \operatorname{sgn}(\text{Id}) a_{1\text{Id}(1)} a_{2\text{Id}(2)} a_{3\text{Id}(3)} + \operatorname{sgn}(\tau_1) a_{1\tau_1(1)} a_{2\tau_1(2)} a_{3\tau_1(3)} \\
 &\quad + \operatorname{sgn}(\tau_2) a_{1\tau_2(1)} a_{2\tau_2(2)} a_{3\tau_2(3)} + \operatorname{sgn}(\tau_3) a_{1\tau_3(1)} a_{2\tau_3(2)} a_{3\tau_3(3)} \\
 &\quad + \operatorname{sgn}(\sigma_1) a_{1\sigma_1(1)} a_{2\sigma_1(2)} a_{3\sigma_1(3)} + \operatorname{sgn}(\sigma_2) a_{1\sigma_2(1)} a_{2\sigma_2(2)} a_{3\sigma_2(3)} \\
 &= a_{11} a_{22} a_{33} - a_{11} a_{23} a_{32} - a_{13} a_{22} a_{31} \\
 &\quad - a_{13} a_{21} a_{33} + a_{12} a_{23} a_{31} + a_{13} a_{21} a_{32}.
 \end{aligned}$$

600 Theorem (Row-Alternancy of Determinants) Let $\mathbf{A} \in M_n(\mathbb{F})$, $\mathbf{A} = [a_{ij}]$. If $\mathbf{B} \in M_n(\mathbb{F})$, $\mathbf{B} = [b_{ij}]$ is the matrix obtained by interchanging the s -th row of \mathbf{A} with its t -th row, then $\det \mathbf{B} = -\det \mathbf{A}$.

Proof: Let τ be the transposition

$$\tau = \begin{bmatrix} s & t \\ \tau(t) & \tau(s) \end{bmatrix}.$$

Then $\sigma\tau(a) = \sigma(a)$ for $a \in \{1, 2, \dots, n\} \setminus \{s, t\}$. Also, $\operatorname{sgn}(\sigma\tau) = \operatorname{sgn}(\sigma)\operatorname{sgn}(\tau) = -\operatorname{sgn}(\sigma)$. As σ ranges through all permutations of S_n , so does $\sigma\tau$, hence

$$\begin{aligned}
 \det \mathbf{B} &= \sum_{\sigma \in S_n} \operatorname{sgn}(\sigma) b_{1\sigma(1)} b_{2\sigma(2)} \cdots b_{s\sigma(s)} \cdots b_{t\sigma(t)} \cdots b_{n\sigma(n)} \\
 &= \sum_{\sigma \in S_n} \operatorname{sgn}(\sigma) a_{1\sigma(1)} a_{2\sigma(2)} \cdots a_{s\sigma(s)} \cdots a_{t\sigma(t)} \cdots a_{n\sigma(n)} \\
 &= - \sum_{\sigma \in S_n} \operatorname{sgn}(\sigma\tau) a_{1\sigma\tau(1)} a_{2\sigma\tau(2)} \cdots a_{s\sigma\tau(s)} \cdots a_{t\sigma\tau(t)} \cdots a_{n\sigma\tau(n)} \\
 &= - \sum_{\lambda \in S_n} \operatorname{sgn}(\lambda) a_{1\lambda(1)} a_{2\lambda(2)} \cdots a_{n\lambda(n)} \\
 &= -\det \mathbf{A}.
 \end{aligned}$$

□

601 Corollary If $\mathbf{A}_{(r:k)}$, $1 \leq k \leq n$ denote the rows of \mathbf{A} and $\sigma \in S_n$, then

$$\det \begin{bmatrix} \mathbf{A}_{(r:\sigma(1))} \\ \mathbf{A}_{(r:\sigma(2))} \\ \vdots \\ \mathbf{A}_{(r:\sigma(n))} \end{bmatrix} = (\operatorname{sgn}(\sigma)) \det \mathbf{A}.$$

An analogous result holds for columns.

Proof: Apply the result of Theorem 600 multiple times. □

602 Theorem Let $\mathbf{A} \in M_n(\mathbb{F})$, $\mathbf{A} = [a_{ij}]$. Then

$$\det \mathbf{A}^T = \det \mathbf{A}.$$

Proof: Let $\mathbf{C} = \mathbf{A}^T$. By definition

$$\begin{aligned}\det \mathbf{A}^T &= \det \mathbf{C} \\ &= \sum_{\sigma \in S_n} \operatorname{sgn}(\sigma) c_{1\sigma(1)} c_{2\sigma(2)} \cdots c_{n\sigma(n)} \\ &= \sum_{\sigma \in S_n} \operatorname{sgn}(\sigma) a_{\sigma(1)1} a_{\sigma(2)2} \cdots a_{\sigma(n)n}.\end{aligned}$$

But the product $a_{\sigma(1)1} a_{\sigma(2)2} \cdots a_{\sigma(n)n}$ also appears in $\det \mathbf{A}$ with the same signum $\operatorname{sgn}(\sigma)$, since the permutation

$$\begin{bmatrix} \sigma(1) & \sigma(2) & \cdots & \sigma(n) \\ 1 & 2 & \cdots & n \end{bmatrix}$$

is the inverse of the permutation

$$\begin{bmatrix} 1 & 2 & \cdots & n \\ \sigma(1) & \sigma(2) & \cdots & \sigma(n) \end{bmatrix}.$$

□

603 Corollary (Column-Alternacy of Determinants) Let $\mathbf{A} \in M_n(\mathbb{F})$, $\mathbf{A} = [a_{ij}]$. If $\mathbf{C} \in M_n(\mathbb{F})$, $\mathbf{C} = [c_{ij}]$ is the matrix obtained by interchanging the s -th column of \mathbf{A} with its t -th column, then $\det \mathbf{C} = -\det \mathbf{A}$.

Proof: This follows upon combining Theorem 600 and Theorem 602. □

604 Theorem (Row Homogeneity of Determinants) Let $\mathbf{A} \in M_n(\mathbb{F})$, $\mathbf{A} = [a_{ij}]$ and $\alpha \in \mathbb{F}$. If $\mathbf{B} \in M_n(\mathbb{F})$, $\mathbf{B} = [b_{ij}]$ is the matrix obtained by multiplying the s -th row of \mathbf{A} by α , then

$$\det \mathbf{B} = \alpha \det \mathbf{A}.$$

Proof: Simply observe that

$$\operatorname{sgn}(\sigma) a_{1\sigma(1)} a_{2\sigma(2)} \cdots \alpha a_{s\sigma(s)} \cdots a_{n\sigma(n)} = \alpha \operatorname{sgn}(\sigma) a_{1\sigma(1)} a_{2\sigma(2)} \cdots a_{s\sigma(s)} \cdots a_{n\sigma(n)}.$$

□

605 Corollary (Column Homogeneity of Determinants) If $\mathbf{C} \in M_n(\mathbb{F})$, $\mathbf{C} = (C_{ij})$ is the matrix obtained by multiplying the s -th column of \mathbf{A} by α , then

$$\det \mathbf{C} = \alpha \det \mathbf{A}.$$

Proof: This follows upon using Theorem 602 and Theorem 604. □

 It follows from Theorem 604 and Corollary 605 that if a row (or column) of a matrix consists of 0_Fs only, then the determinant of this matrix is 0_F.

606 Example

$$\det \begin{bmatrix} x & 1 & a \\ x^2 & 1 & b \\ x^3 & 1 & c \end{bmatrix} = x \det \begin{bmatrix} 1 & 1 & a \\ x & 1 & b \\ x^2 & 1 & c \end{bmatrix}.$$

607 Corollary

$$\det(\alpha A) = \alpha^n \det A.$$

Proof: Since there are n columns, we are able to pull out one factor of α from each one. \square

608 Example Recall that a matrix A is *skew-symmetric* if $A = -A^T$. Let $A \in M_{2001}(\mathbb{R})$ be skew-symmetric. Prove that $\det A = 0$.

Solution: We have

$$\det A = \det(-A^T) = (-1)^{2001} \det A^T = -\det A,$$

and so $2 \det A = 0$, from where $\det A = 0$.

609 Lemma (Row-Linearity and Column-Linearity of Determinants) Let $A \in M_n(\mathbb{F})$, $A = [a_{ij}]$. For a fixed row s , suppose that $a_{sj} = b_{sj} + c_{sj}$ for each $j \in [1; n]$. Then

$$\begin{aligned} & \det \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{(s-1)1} & a_{(s-1)2} & \cdots & a_{(s-1)n} \\ b_{s1} + c_{s1} & b_{s2} + c_{s2} & \cdots & b_{sn} + c_{sn} \\ a_{(s+1)1} & a_{(s+1)2} & \cdots & a_{(s+1)n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{bmatrix} \\ &= \det \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{(s-1)1} & a_{(s-1)2} & \cdots & a_{(s-1)n} \\ b_{s1} & b_{s2} & \cdots & b_{sn} \\ a_{(s+1)1} & a_{(s+1)2} & \cdots & a_{(s+1)n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{bmatrix} \\ &+ \det \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{(s-1)1} & a_{(s-1)2} & \cdots & a_{(s-1)n} \\ c_{s1} & c_{s2} & \cdots & c_{sn} \\ a_{(s+1)1} & a_{(s+1)2} & \cdots & a_{(s+1)n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{bmatrix}. \end{aligned}$$

An analogous result holds for columns.

Proof: Put

$$S = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{(s-1)1} & a_{(s-1)2} & \cdots & a_{(s-1)n} \\ b_{s1} + c_{s1} & b_{s2} + c_{s2} & \cdots & b_{sn} + c_{sn} \\ a_{(s+1)1} & a_{(s+1)2} & \cdots & a_{(s+1)n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{bmatrix},$$

$$\mathbf{T} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{(s-1)1} & a_{(s-1)2} & \cdots & a_{(s-1)n} \\ b_{s1} & b_{s2} & \cdots & b_{sn} \\ a_{(s+1)1} & a_{(s+1)2} & \cdots & a_{(s+1)n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{bmatrix}$$

and

$$\mathbf{U} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{(s-1)1} & a_{(s-1)2} & \cdots & a_{(s-1)n} \\ c_{s1} & c_{s2} & \cdots & c_{sn} \\ a_{(s+1)1} & a_{(s+1)2} & \cdots & a_{(s+1)n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{bmatrix}.$$

Then

$$\begin{aligned} \det \mathbf{S} &= \sum_{\sigma \in S_n} \operatorname{sgn}(\sigma) a_{1\sigma(1)} a_{2\sigma(2)} \cdots a_{(s-1)\sigma(s-1)} (b_{s\sigma(s)} \\ &\quad + c_{s\sigma(s)}) a_{(s+1)\sigma(s+1)} \cdots a_{n\sigma(n)} \\ &= \sum_{\sigma \in S_n} \operatorname{sgn}(\sigma) a_{1\sigma(1)} a_{2\sigma(2)} \cdots a_{(s-1)\sigma(s-1)} b_{s\sigma(s)} a_{(s+1)\sigma(s+1)} \cdots a_{n\sigma(n)} \\ &\quad + \sum_{\sigma \in S_n} \operatorname{sgn}(\sigma) a_{1\sigma(1)} a_{2\sigma(2)} \cdots a_{(s-1)\sigma(s-1)} c_{s\sigma(s)} a_{(s+1)\sigma(s+1)} \cdots a_{n\sigma(n)} \\ &= \det \mathbf{T} + \det \mathbf{U}. \end{aligned}$$

By applying the above argument to \mathbf{A}^T , we obtain the result for columns.

□

610 Lemma If two rows or two columns of $\mathbf{A} \in M_n(\mathbb{F})$, $\mathbf{A} = [a_{ij}]$ are identical, then $\det \mathbf{A} = 0_{\mathbb{F}}$.

Proof: Suppose $a_{sj} = a_{tj}$ for $s \neq t$ and for all $j \in [1; n]$. In particular, this means that for any $\sigma \in S_n$ we have $a_{s\sigma(t)} = a_{t\sigma(t)}$ and $a_{t\sigma(s)} = a_{s\sigma(s)}$. Let τ be the transposition

$$\tau = \begin{bmatrix} s & t \\ \tau(t) & \tau(s) \end{bmatrix}.$$

Then $\sigma\tau(a) = \sigma(a)$ for $a \in \{1, 2, \dots, n\} \setminus \{s, t\}$. Also, $\operatorname{sgn}(\sigma\tau) = \operatorname{sgn}(\sigma)\operatorname{sgn}(\tau) = -\operatorname{sgn}(\sigma)$. As σ runs through all even permutations, $\sigma\tau$ runs through all odd permutations, and viceversa.

Therefore

$$\begin{aligned}
 \det \mathbf{A} &= \sum_{\sigma \in S_n} \operatorname{sgn}(\sigma) a_{1\sigma(1)} a_{2\sigma(2)} \cdots a_{s\sigma(s)} \cdots a_{t\sigma(t)} \cdots a_{n\sigma(n)} \\
 &= \sum_{\substack{\sigma \in S_n \\ \operatorname{sgn}(\sigma)=1}} (\operatorname{sgn}(\sigma) a_{1\sigma(1)} a_{2\sigma(2)} \cdots a_{s\sigma(s)} \cdots a_{t\sigma(t)} \cdots a_{n\sigma(n)}) \\
 &\quad + \operatorname{sgn}(\sigma\tau) a_{1\sigma\tau(1)} a_{2\sigma\tau(2)} \cdots a_{s\sigma\tau(s)} \cdots a_{t\sigma\tau(t)} \cdots a_{n\sigma\tau(n)}) \\
 &= \sum_{\substack{\sigma \in S_n \\ \operatorname{sgn}(\sigma)=1}} \operatorname{sgn}(\sigma) (a_{1\sigma(1)} a_{2\sigma(2)} \cdots a_{s\sigma(s)} \cdots a_{t\sigma(t)} \cdots a_{n\sigma(n)} \\
 &\quad - a_{1\sigma(1)} a_{2\sigma(2)} \cdots a_{s\sigma(t)} \cdots a_{t\sigma(s)} \cdots a_{n\sigma(n)}) \\
 &= \sum_{\substack{\sigma \in S_n \\ \operatorname{sgn}(\sigma)=1}} \operatorname{sgn}(\sigma) (a_{1\sigma(1)} a_{2\sigma(2)} \cdots a_{s\sigma(s)} \cdots a_{t\sigma(t)} \cdots a_{n\sigma(n)} \\
 &\quad - a_{1\sigma(1)} a_{2\sigma(2)} \cdots a_{t\sigma(t)} \cdots a_{s\sigma(s)} \cdots a_{n\sigma(n)}) \\
 &= 0_F.
 \end{aligned}$$

Arguing on \mathbf{A}^T will yield the analogous result for the columns. \square

611 Corollary If two rows or two columns of $\mathbf{A} \in M_n(\mathbb{F})$, $\mathbf{A} = [a_{ij}]$ are proportional, then $\det \mathbf{A} = 0_F$.

Proof: Suppose $a_{sj} = \alpha a_{tj}$ for $s \neq t$ and for all $j \in [1; n]$. If \mathbf{B} is the matrix obtained by pulling out the factor α from the s -th row then $\det \mathbf{A} = \alpha \det \mathbf{B}$. But now the s -th and the t -th rows in \mathbf{B} are identical, and so $\det \mathbf{B} = 0_F$. Arguing on \mathbf{A}^T will yield the analogous result for the columns.

\square

612 Example

$$\det \begin{bmatrix} 1 & a & b \\ 1 & a & c \\ 1 & a & d \end{bmatrix} = a \det \begin{bmatrix} 1 & 1 & b \\ 1 & 1 & c \\ 1 & 1 & d \end{bmatrix} = 0,$$

since on the last determinant the first two columns are identical.

613 Theorem (Multilinearity of Determinants) Let $\mathbf{A} \in M_n(\mathbb{F})$, $\mathbf{A} = [a_{ij}]$ and $\alpha \in \mathbb{F}$. If $\mathbf{X} \in M_n(\mathbb{F})$, $\mathbf{X} = (x_{ij})$ is the matrix obtained by the row transvection $R_s + \alpha R_t \rightarrow R_s$ then $\det \mathbf{X} = \det \mathbf{A}$. Similarly, if $\mathbf{Y} \in M_n(\mathbb{F})$, $\mathbf{Y} = (y_{ij})$ is the matrix obtained by the column transvection $C_s + \alpha C_t \rightarrow C_s$ then $\det \mathbf{Y} = \det \mathbf{A}$.

Proof: For the row transvection it suffices to take $b_{sj} = a_{sj}$, $c_{sj} = \alpha a_{tj}$ for $j \in [1; n]$ in Lemma 609. With the same notation as in the lemma, $\mathbf{T} = \mathbf{A}$, and so,

$$\det \mathbf{X} = \det \mathbf{T} + \det \mathbf{U} = \det \mathbf{A} + \det \mathbf{U}.$$

But \mathbf{U} has its s -th and t -th rows proportional ($s \neq t$), and so by Corollary 611 $\det \mathbf{U} = 0_F$. Hence $\det \mathbf{X} = \det \mathbf{A}$. To obtain the result for column transvections it suffices now to also apply Theorem 602. \square

614 Example Demonstrate, without actually calculating the determinant that

$$\det \begin{bmatrix} 2 & 9 & 9 \\ 4 & 6 & 8 \\ 7 & 4 & 1 \end{bmatrix}$$

is divisible by 13.

Solution: Observe that 299, 468 and 741 are all divisible by 13. Thus

$$\det \begin{bmatrix} 2 & 9 & 9 \\ 4 & 6 & 8 \\ 7 & 4 & 1 \end{bmatrix} \stackrel{C_3 + 10C_2 + 100C_1 \rightarrow C_3}{=} \det \begin{bmatrix} 2 & 9 & 299 \\ 4 & 6 & 468 \\ 7 & 4 & 741 \end{bmatrix} = 13 \det \begin{bmatrix} 2 & 9 & 23 \\ 4 & 6 & 36 \\ 7 & 4 & 57 \end{bmatrix},$$

which shews that the determinant is divisible by 13.

615 Theorem The determinant of a triangular matrix (upper or lower) is the product of its diagonal elements.

Proof: Let $A \in M_n(\mathbb{F})$, $A = [a_{ij}]$ be a triangular matrix. Observe that if $\sigma \neq \text{Id}$ then $a_{i\sigma(i)}a_{\sigma(i)\sigma^2(i)} = 0_{\mathbb{F}}$ occurs in the product

$$a_{1\sigma(1)}a_{2\sigma(2)} \cdots a_{n\sigma(n)}.$$

Thus

$$\begin{aligned} \det A &= \sum_{\sigma \in S_n} \text{sgn}(\sigma) a_{1\sigma(1)}a_{2\sigma(2)} \cdots a_{n\sigma(n)} \\ &= \text{sgn}(\text{Id}) a_{1\text{Id}(1)}a_{2\text{Id}(2)} \cdots a_{n\text{Id}(n)} = a_{11}a_{22} \cdots a_{nn}. \end{aligned}$$

□

616 Example The determinant of the $n \times n$ identity matrix I_n over a field \mathbb{F} is

$$\det I_n = 1_{\mathbb{F}}.$$

617 Example Find

$$\det \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}.$$

Solution: We have

$$\begin{aligned}
 & \det \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} \xrightarrow[\sim]{C_3 - 3C_1 \rightarrow C_3} \det \begin{bmatrix} 1 & 0 & 0 \\ 4 & -3 & -6 \\ 7 & -6 & -12 \end{bmatrix} \\
 & = (-3)(-6) \det \begin{bmatrix} 1 & 0 & 0 \\ 4 & 1 & 1 \\ 7 & 2 & 2 \end{bmatrix} \\
 & = 0,
 \end{aligned}$$

since in this last matrix the second and third columns are identical and so Lemma 610 applies.

618 Theorem Let $(\mathbf{A}, \mathbf{B}) \in (\mathbf{M}_n(\mathbb{F}))^2$. Then

$$\det(\mathbf{AB}) = (\det \mathbf{A})(\det \mathbf{B}).$$

Proof: Put $\mathbf{D} = \mathbf{AB}$, $\mathbf{d}_{ij} = \sum_{k=1}^n a_{ik} b_{kj}$. If $\mathbf{A}_{(c:k)}$, $\mathbf{D}_{(c:k)}$, $1 \leq k \leq n$ denote the columns of \mathbf{A} and \mathbf{D} , respectively, observe that

$$\mathbf{D}_{(c:k)} = \sum_{l=1}^n b_{lk} \mathbf{A}_{(c:l)}, \quad 1 \leq k \leq n.$$

Applying Corollary 605 and Lemma 609 multiple times, we obtain

$$\begin{aligned}
 \det \mathbf{D} &= \det(\mathbf{D}_{(c:1)}, \mathbf{D}_{(c:2)}, \dots, \mathbf{D}_{(c:n)}) \\
 &= \sum_{j_1=1}^n \sum_{j_2=1}^n \cdots \sum_{j_n=1}^n b_{1j_1} b_{2j_2} \cdots b_{nj_n} \\
 &\quad \cdot \det(\mathbf{A}_{(c:j_1)}, \mathbf{A}_{(c:j_2)}, \dots, \mathbf{A}_{(c:j_n)}).
 \end{aligned}$$

By Lemma 610, if any two of the $\mathbf{A}_{(c:j_l)}$ are identical, the determinant on the right vanishes. So each one of the j_l is different in the non-vanishing terms and so the map

$$\begin{array}{ccc}
 \sigma : & \{1, 2, \dots, n\} & \rightarrow \{1, 2, \dots, n\} \\
 & l & \mapsto j_l
 \end{array}$$

is a permutation. Here $j_l = \sigma(l)$. Therefore, for the non-vanishing

$$\det(\mathbf{A}_{(c:j_1)}, \mathbf{A}_{(c:j_2)}, \dots, \mathbf{A}_{(c:j_n)})$$

we have in view of Corollary 601,

$$\begin{aligned}
 \det(\mathbf{A}_{(c:j_1)}, \mathbf{A}_{(c:j_2)}, \dots, \mathbf{A}_{(c:j_n)}) &= (\text{sgn}(\sigma)) \det(\mathbf{A}_{(c:1)}, \mathbf{A}_{(c:2)}, \dots, \mathbf{A}_{(c:n)}) \\
 &= (\text{sgn}(\sigma)) \det \mathbf{A}.
 \end{aligned}$$

We deduce that

$$\begin{aligned}
 \det(\mathbf{AB}) &= \det \mathbf{D} \\
 &= \sum_{j_1=1}^n b_{1j_1} b_{2j_2} \cdots b_{nj_n} \det(\mathbf{A}_{(c:j_1)}, \mathbf{A}_{(c:j_2)}, \dots, \mathbf{A}_{(c:j_n)}) \\
 &= (\det \mathbf{A}) \sum_{\sigma \in S_n} (\text{sgn}(\sigma)) b_{1\sigma(1)} b_{2\sigma(2)} \cdots b_{n\sigma(n)} \\
 &= (\det \mathbf{A})(\det \mathbf{B}),
 \end{aligned}$$

as we wanted to shew. \square

By applying the preceding theorem multiple times we obtain

619 Corollary If $\mathbf{A} \in M_n(\mathbb{F})$ and if k is a positive integer then

$$\det \mathbf{A}^k = (\det \mathbf{A})^k.$$

620 Corollary If $\mathbf{A} \in GL_n(\mathbb{F})$ and if k is a positive integer then $\det \mathbf{A} \neq 0_{\mathbb{F}}$ and

$$\det \mathbf{A}^{-k} = (\det \mathbf{A})^{-k}.$$

Proof: We have $\mathbf{AA}^{-1} = I_n$ and so by Theorem 618 $(\det \mathbf{A})(\det \mathbf{A}^{-1}) = 1_{\mathbb{F}}$, from where the result follows. \square

621 Problem Let

$$\Omega = \det \begin{bmatrix} bc & ca & ab \\ a & b & c \\ a^2 & b^2 & c^2 \end{bmatrix}.$$

Without expanding either determinant, prove that

$$\Omega = \det \begin{bmatrix} 1 & 1 & 1 \\ a^2 & b^2 & c^2 \\ a^3 & b^3 & c^3 \end{bmatrix}.$$

622 Problem Demonstrate that

$$\Omega = \det \begin{bmatrix} a-b-c & 2a & 2a \\ 2b & b-c-a & 2b \\ 2c & 2c & c-a-b \end{bmatrix} = (a+b+c).$$

623 Problem After the indicated column operations on a 3×3 matrix \mathbf{A} with $\det \mathbf{A} = -540$, matrices $\mathbf{A}_1, \mathbf{A}_2, \dots, \mathbf{A}_5$ are successively obtained:

$$\mathbf{A} \xrightarrow{C_1 + 3C_2 \rightarrow C_1} \mathbf{A}_1 \xrightarrow{C_2 \leftrightarrow C_3} \mathbf{A}_2 \xrightarrow{3C_2 - C_1 \rightarrow C_2} \mathbf{A}_3 \xrightarrow{C_1 - 3C_2 \rightarrow C_1} \mathbf{A}_4 \xrightarrow{2C_1 \rightarrow C_1} \mathbf{A}_5$$

Determine the numerical values of $\det \mathbf{A}_1, \det \mathbf{A}_2, \det \mathbf{A}_3, \det \mathbf{A}_4$ and $\det \mathbf{A}_5$.

624 Problem Let $\mathbf{A}, \mathbf{B}, \mathbf{C}$ be 3×3 matrices with $\det \mathbf{A} = 3, \det \mathbf{B}^3 = -8, \det \mathbf{C} = 2$. Compute (i) $\det \mathbf{ABC}$, (ii) $\det 5\mathbf{AC}$, (iii) $\det \mathbf{A}^3 \mathbf{B}^{-3} \mathbf{C}^{-1}$. Express your answers as fractions.

625 Problem Shew that $\forall \mathbf{A} \in M_n(\mathbb{R})$,

$$\exists (\mathbf{X}, \mathbf{Y}) \in (M_n(\mathbb{R}))^2, (\det \mathbf{X})(\det \mathbf{Y}) \neq 0$$

such that

$$\mathbf{A} = \mathbf{X} + \mathbf{Y}.$$

That is, any square matrix over \mathbb{R} can be written as a sum of two matrices whose determinant is not zero.

626 Problem Prove or disprove! The set $X = \{\mathbf{A} \in M_n(\mathbb{F}) : \det \mathbf{A} = 0_{\mathbb{F}}\}$ is a vector subspace of $M_n(\mathbb{F})$.

7.4 Laplace Expansion

We now develop a more computationally convenient approach to determinants.

Put

$$C_{ij} = \sum_{\substack{\sigma \in S_n \\ \sigma(i)=j}} (\text{sgn}(\sigma)) a_{1\sigma(1)} a_{2\sigma(2)} \cdots a_{n\sigma(n)}.$$

Then

$$\begin{aligned}
 \det A &= \sum_{\sigma \in S_n} (\text{sgn}(\sigma)) a_{1\sigma(1)} a_{2\sigma(2)} \cdots a_{n\sigma(n)} \\
 &= \sum_{i=1}^n a_{ij} \sum_{\substack{\sigma \in S_n \\ \sigma(i)=j}} (\text{sgn}(\sigma)) a_{1\sigma(1)} a_{2\sigma(2)} \\
 &\quad \cdots a_{(i-1)\sigma(i-1)} a_{(i+1)\sigma(i+1)} \cdots a_{n\sigma(n)} \\
 &= \sum_{i=1}^n a_{ij} C_{ij},
 \end{aligned} \tag{7.5}$$

is the expansion of $\det A$ along the j -th column. Similarly,

$$\begin{aligned}
 \det A &= \sum_{\sigma \in S_n} (\text{sgn}(\sigma)) a_{1\sigma(1)} a_{2\sigma(2)} \cdots a_{n\sigma(n)} \\
 &= \sum_{j=1}^n a_{ij} \sum_{\substack{\sigma \in S_n \\ \sigma(i)=j}} (\text{sgn}(\sigma)) a_{1\sigma(1)} a_{2\sigma(2)} \\
 &\quad \cdots a_{(i-1)\sigma(i-1)} a_{(i+1)\sigma(i+1)} \cdots a_{n\sigma(n)} \\
 &= \sum_{j=1}^n a_{ij} C_{ij},
 \end{aligned}$$

is the expansion of $\det A$ along the i -th row.

627 Definition Let $A \in M_n(\mathbb{F})$, $A = [a_{ij}]$. The ij -th minor $A_{ij} \in M_{n-1}(\mathbb{R})$ is the $(n-1) \times (n-1)$ matrix obtained by deleting the i -th row and the j -th column from A .

628 Example If

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$$

then, for example,

$$A_{11} = \begin{bmatrix} 5 & 6 \\ 8 & 9 \end{bmatrix}, \quad A_{12} = \begin{bmatrix} 4 & 6 \\ 7 & 9 \end{bmatrix}, \quad A_{21} = \begin{bmatrix} 2 & 3 \\ 8 & 9 \end{bmatrix}, \quad A_{22} = \begin{bmatrix} 1 & 3 \\ 7 & 9 \end{bmatrix}, \quad A_{33} = \begin{bmatrix} 1 & 2 \\ 4 & 5 \end{bmatrix}.$$

629 Theorem Let $A \in M_n(\mathbb{F})$. Then

$$\det A = \sum_{i=1}^n a_{ij} (-1)^{i+j} \det A_{ij} = \sum_{j=1}^n a_{ij} (-1)^{i+j} \det A_{ij}.$$

Proof: It is enough to shew, in view of 7.5 that

$$(-1)^{i+j} \det A_{ij} = C_{ij}.$$

Now,

$$\begin{aligned}
 C_{nn} &= \sum_{\substack{\sigma \in S_n \\ \sigma(n)=n}} \text{sgn}(\sigma) a_{1\sigma(1)} a_{2\sigma(2)} \cdots a_{(n-1)\sigma(n-1)} \\
 &= \sum_{\tau \in S_{n-1}} \text{sgn}(\tau) a_{1\tau(1)} a_{2\tau(2)} \cdots a_{(n-1)\tau(n-1)} \\
 &= \det A_{nn},
 \end{aligned}$$

since the second sum shewn is the determinant of the submatrix obtained by deleting the last row and last column from \mathbf{A} .

To find C_{ij} for general ij we perform some row and column interchanges to \mathbf{A} in order to bring a_{ij} to the n -th position. We thus bring the i -th row to the n -th row by a series of transpositions, first swapping the i -th and the $(i+1)$ -th row, then swapping the new $(i+1)$ -th row and the $(i+2)$ -th row, and so forth until the original i -th row makes it to the n -th row. We have made thereby $n-i$ interchanges. To this new matrix we perform analogous interchanges to the j -th column, thereby making $n-j$ interchanges. We have made a total of $2n-i-j$ interchanges. Observe that $(-1)^{2n-i-j} = (-1)^{i+j}$. Call the analogous quantities in the resulting matrix \mathbf{A}' , C'_{nn} , A'_{nn} . Then

$$C_{ij} = C'_{nn} = \det A'_{nn} = (-1)^{i+j} \det A_{ij},$$

by virtue of Corollary 601.

□

 It is irrelevant which row or column we choose to expand a determinant of a square matrix. We always obtain the same result. The sign pattern is given by

$$\begin{bmatrix} + & - & + & - & \dots \\ - & + & - & + & \vdots \\ + & - & + & - & \vdots \\ \vdots & \vdots & \vdots & \vdots & \vdots \end{bmatrix}$$

630 Example

Find

$$\det \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$$

by expanding along the first row.

Solution: We have

$$\begin{aligned} \det \mathbf{A} &= 1(-1)^{1+1} \det \begin{bmatrix} 5 & 6 \\ 8 & 9 \end{bmatrix} + 2(-1)^{1+2} \det \begin{bmatrix} 4 & 6 \\ 7 & 9 \end{bmatrix} + 3(-1)^{1+3} \det \begin{bmatrix} 4 & 5 \\ 7 & 8 \end{bmatrix} \\ &= 1(45 - 48) - 2(36 - 42) + 3(32 - 35) = 0. \end{aligned}$$

631 Example

Evaluate the Vandermonde determinant

$$\det \begin{bmatrix} 1 & 1 & 1 \\ a & b & c \\ a^2 & b^2 & c^2 \end{bmatrix}.$$

Solution:

$$\begin{aligned}
 \det \begin{bmatrix} 1 & 1 & 1 \\ a & b & c \\ a^2 & b^2 & c^2 \end{bmatrix} &= \det \begin{bmatrix} 1 & 0 & 0 \\ a & b-a & c-a \\ a^2 & b^2-a^2 & c^2-a^2 \end{bmatrix} \\
 &= \det \begin{bmatrix} b-a & c-a \\ b^2-c^2 & c^2-a^2 \end{bmatrix} \\
 &= (b-a)(c-a) \det \begin{bmatrix} 1 & 1 \\ b+a & c+a \end{bmatrix} \\
 &= (b-a)(c-a)(c-b).
 \end{aligned}$$

632 Example Evaluate the determinant

$$\det A = \det \begin{bmatrix} 1 & 2 & 3 & 4 & \cdots & 2000 \\ 2 & 1 & 2 & 3 & \cdots & 1999 \\ 3 & 2 & 1 & 2 & \cdots & 1998 \\ 4 & 3 & 2 & 1 & \cdots & 1997 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 2000 & 1999 & 1998 & 1997 & \cdots & 1 \end{bmatrix}.$$

Solution: Applying $R_n - R_{n+1} \rightarrow R_n$ for $1 \leq n \leq 1999$, the determinant becomes

$$\det \begin{bmatrix} -1 & 1 & 1 & 1 & \cdots & 1 & 1 \\ -1 & -1 & 1 & 1 & \cdots & 1 & 1 \\ -1 & -1 & -1 & 1 & \cdots & 1 & 1 \\ -1 & -1 & -1 & -1 & \cdots & 1 & 1 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ -1 & -1 & -1 & -1 & \cdots & -1 & 1 \\ 2000 & 1999 & 1998 & 1997 & \cdots & 2 & 1 \end{bmatrix}.$$

Applying now $C_n + C_{2000} \rightarrow C_n$ for $1 \leq n \leq 1999$, we obtain

$$\det \begin{bmatrix} 0 & 2 & 2 & 2 & \cdots & 2 & 1 \\ 0 & 0 & 2 & 2 & \cdots & 2 & 1 \\ 0 & 0 & 0 & 2 & \cdots & 2 & 1 \\ 0 & 0 & 0 & 0 & \cdots & 2 & 1 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & 0 & 0 & \cdots & 0 & 1 \\ 2001 & 2000 & 1999 & 1998 & \cdots & 3 & 1 \end{bmatrix}.$$

This last determinant we expand along the first column. We have

$$-2001 \det \begin{bmatrix} 2 & 2 & 2 & \cdots & 2 & 1 \\ 0 & 2 & 2 & \cdots & 2 & 1 \\ 0 & 0 & 2 & \cdots & 2 & 1 \\ 0 & 0 & 0 & \cdots & 2 & 1 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & 0 & \cdots & 0 & 1 \end{bmatrix} = -2001(2^{1998}).$$

633 Definition Let $A \in M_n(\mathbb{F})$. The *classical adjoint* or *adjugate* of A is the $n \times n$ matrix $\text{adj}(A)$ whose entries are given by

$$[\text{adj}(A)]_{ij} = (-1)^{i+j} \det A_{ji},$$

where A_{ji} is the ji -th minor of A .

634 Theorem Let $A \in M_n(\mathbb{F})$. Then

$$(A \text{adj}(A)) = A(A \text{adj}(A)) = (\det A)I_n.$$

Proof: We have

$$\begin{aligned} [A(A \text{adj}(A))]_{ij} &= \sum_{k=1}^n a_{ik} [\text{adj}(A)]_{kj} \\ &= \sum_{k=1}^n a_{ik} (-1)^{i+k} \det A_{jk}. \end{aligned}$$

Now, this last sum is $\det A$ if $i = j$ by virtue of Theorem 629. If $i \neq j$ it is 0, since then the j -th row is identical to the i -th row and this determinant is $0_{\mathbb{F}}$ by virtue of Lemma 610. Thus on the diagonal entries we get $\det A$ and the off-diagonal entries are $0_{\mathbb{F}}$. This proves the theorem. \square

The next corollary follows immediately.

635 Corollary Let $A \in M_n(\mathbb{F})$. Then A is invertible if and only if $\det A \neq 0_{\mathbb{F}}$ and

$$A^{-1} = \frac{\text{adj}(A)}{\det A}.$$

636 Problem Find

$$\det \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$$

by expanding along the second column.

637 Problem Compute the determinant

$$\det \begin{bmatrix} 1 & 0 & -1 & 1 \\ 2 & 0 & 0 & 1 \\ 666 & -3 & -1 & 1000000 \\ 1 & 0 & 0 & 1 \end{bmatrix}.$$

638 Problem If

$$\det \begin{bmatrix} 1 & 1 & 1 & 1 \\ x & a & 0 & 0 \\ x & 0 & b & 0 \\ x & 0 & 0 & c \end{bmatrix} = 0,$$

and $xabc \neq 0$, prove that

$$\frac{1}{x} = \frac{1}{a} + \frac{1}{b} + \frac{1}{c}.$$

639 Problem Prove that

$$\det \begin{bmatrix} 0 & a & b & 0 \\ a & 0 & b & 0 \\ 0 & a & 0 & b \\ 1 & 1 & 1 & 1 \end{bmatrix} = 2ab(a-b).$$

640 Problem Demonstrate that

$$\det \begin{bmatrix} a & 0 & b & 0 \\ 0 & a & 0 & b \\ c & 0 & d & 0 \\ 0 & c & 0 & d \end{bmatrix} = (ad - bc)^2.$$

641 Problem Use induction to show that

$$\det \begin{bmatrix} 1 & 1 & 1 & \cdots & 1 & 1 \\ 1 & 0 & 0 & \vdots & 0 & 0 \\ 0 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \cdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & 0 \end{bmatrix} = (-1)^{n+1}.$$

642 Problem Let

$$\mathbf{A} = \begin{bmatrix} 1 & n & n & n & \cdots & n \\ n & 2 & n & n & \vdots & n \\ n & n & 3 & n & \cdots & n \\ n & n & n & 4 & \cdots & n \\ \vdots & \vdots & \vdots & \cdots & \vdots & \vdots \\ n & n & n & n & n & n \end{bmatrix},$$

that is, $\mathbf{A} \in M_n(\mathbb{R})$, $\mathbf{A} = [a_{ij}]$ is a matrix such that $a_{kk} = n$ and $a_{ij} = n$ when $i \neq j$. Find $\det \mathbf{A}$.

643 Problem Let $n \in \mathbb{N}$, $n > 1$ be an odd integer. Recall that the binomial coefficients $\binom{n}{k}$ satisfy $\binom{n}{n} = \binom{n}{0} = 1$ and that for $1 \leq k \leq n$,

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}.$$

Prove that

$$\det \begin{bmatrix} 1 & \binom{n}{1} & \binom{n}{2} & \cdots & \binom{n}{n-1} & 1 \\ 1 & 1 & \binom{n}{1} & \cdots & \binom{n}{n-2} & \binom{n}{n-1} \\ \binom{n}{n-1} & 1 & 1 & \cdots & \binom{n}{n-3} & \binom{n}{n-2} \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ \binom{n}{1} & \binom{n}{2} & \binom{n}{3} & \cdots & 1 & 1 \end{bmatrix} = (1 + (-1)^n)^n.$$

644 Problem Let $\mathbf{A} \in GL_n(\mathbb{F})$, $n > 1$. Prove that $\det(\text{adj}(\mathbf{A})) = (\det \mathbf{A})^{n-1}$.

645 Problem Let $(\mathbf{A}, \mathbf{B}, \mathbf{S}) \in (GL_n(\mathbb{F}))^3$. Prove that

$$\textcircled{1} \quad \text{adj}(\text{adj}(\mathbf{A})) = (\det \mathbf{A})^{n-2} \mathbf{A}.$$

$$\textcircled{2} \quad \text{adj}(\mathbf{AB}) = \text{adj}(\mathbf{A}) \text{adj}(\mathbf{B}).$$

$$\textcircled{3} \quad \text{adj}(\mathbf{SAS}^{-1}) = \mathbf{S}(\text{adj}(\mathbf{A}))\mathbf{S}^{-1}.$$

646 Problem If $\mathbf{A} \in GL_2(\mathbb{F})$, and let k be a positive integer. Prove that $\det(\text{adj} \cdots \text{adj}(\mathbf{A})) = \det \mathbf{A}$.

7.5 Determinants and Linear Systems

647 Theorem Let $A \in M_n(\mathbb{F})$. The following are all equivalent

- ① $\det A \neq 0_{\mathbb{F}}$.
- ② A is invertible.
- ③ There exists a unique solution $X \in M_{n \times 1}(\mathbb{F})$ to the equation $AX = Y$.
- ④ If $AX = 0_{n \times 1}$ then $X = 0_{n \times 1}$.

Proof: We prove the implications in sequence:

- ① \implies ②: follows from Corollary 635
- ② \implies ③: If A is invertible and $AX = Y$ then $X = A^{-1}Y$ is the unique solution of this equation.
- ③ \implies ④: follows by putting $Y = 0_{n \times 1}$
- ④ \implies ①: Let R be the row echelon form of A . Since $RX = 0_{n \times 1}$ has only $X = 0_{n \times 1}$ as a solution, every entry on the diagonal of R must be non-zero, R must be triangular, and hence $\det R \neq 0_{\mathbb{F}}$. Since $A = PR$ where P is an invertible $n \times n$ matrix, we deduce that $\det A = \det P \det R \neq 0_{\mathbb{F}}$.

□

The contrapositive form of the implications ① and ④ will be used later. Here it is for future reference.

648 Corollary Let $A \in M_n(\mathbb{F})$. If there is $X \neq 0_{n \times 1}$ such that $AX = 0_{n \times 1}$ then $\det A = 0_{\mathbb{F}}$.

Chapter 8

Eigenvalues and Eigenvectors

8.1 Similar Matrices

649 Definition We say that $A \in M_n(\mathbb{F})$ is *similar* to $B \in M_n(\mathbb{F})$ if there exist a matrix $P \in GL_n(\mathbb{F})$ such that

$$B = PAP^{-1}.$$

650 Theorem Similarity is an equivalence relation.

Proof: Let $A \in M_n(\mathbb{F})$. Then $A = I_n A I_n^{-1}$, so similarity is reflexive. If $B = PAP^{-1}$ ($P \in GL_n(\mathbb{F})$) then $A = P^{-1}BP$ so similarity is symmetric. Finally, if $B = PAP^{-1}$ and $C = QBQ^{-1}$ ($P \in GL_n(\mathbb{F})$, $Q \in GL_n(\mathbb{F})$) then $C = QPAP^{-1}Q^{-1} = QPA(QP)^{-1}$ and so similarity is transitive. \square

Since similarity is an equivalence relation, it partitions the set of $n \times n$ matrices into equivalence classes by Theorem 37.

651 Definition A matrix is said to be *diagonalisable* if it is similar to a diagonal matrix.

Suppose that

$$A = \begin{bmatrix} \lambda_1 & 0 & 0 & \cdots & 0 \\ 0 & \lambda_2 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ 0 & 0 & 0 & \cdots & \lambda_n \end{bmatrix}.$$

Then if K is a positive integer

$$A^K = \begin{bmatrix} \lambda_1^K & 0 & 0 & \cdots & 0 \\ 0 & \lambda_2^K & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ 0 & 0 & 0 & \cdots & \lambda_n^K \end{bmatrix}.$$

In particular, if \mathbf{B} is similar to \mathbf{A} then

$$\mathbf{B}^K = \underbrace{(\mathbf{P}\mathbf{A}\mathbf{P}^{-1})(\mathbf{P}\mathbf{A}\mathbf{P}^{-1}) \cdots (\mathbf{P}\mathbf{A}\mathbf{P}^{-1})}_{K \text{ factors}} = \mathbf{P}\mathbf{A}^K\mathbf{P}^{-1} = \mathbf{P} \begin{bmatrix} \lambda_1^K & 0 & 0 & \cdots & 0 \\ 0 & \lambda_2^K & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \lambda_n^K \end{bmatrix} \mathbf{P}^{-1},$$

so we have a simpler way of computing \mathbf{B}^K . Our task will now be to establish when a particular square matrix is diagonalisable.

8.2 Eigenvalues and Eigenvectors

Let $\mathbf{A} \in M_n(\mathbb{F})$ be a square diagonalisable matrix. Then there exist $\mathbf{P} \in GL_n(\mathbb{F})$ and a diagonal matrix $\mathbf{D} \in M_n(\mathbb{F})$ such that $\mathbf{P}^{-1}\mathbf{A}\mathbf{P} = \mathbf{D}$, whence $\mathbf{A}\mathbf{P} = \mathbf{D}\mathbf{P}$. Put

$$\mathbf{D} = \begin{bmatrix} \lambda_1 & 0 & 0 & \cdots & 0 \\ 0 & \lambda_2 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \lambda_n \end{bmatrix}, \quad \mathbf{P} = [\mathbf{P}_1; \mathbf{P}_2; \cdots; \mathbf{P}_n],$$

where the \mathbf{P}_k are the columns of \mathbf{P} . Then

$$\mathbf{A}\mathbf{P} = \mathbf{D}\mathbf{P} \implies [\mathbf{A}\mathbf{P}_1; \mathbf{A}\mathbf{P}_2; \cdots; \mathbf{A}\mathbf{P}_n] = [\lambda_1\mathbf{P}_1; \lambda_2\mathbf{P}_2; \cdots; \lambda_n\mathbf{P}_n],$$

from where it follows that $\mathbf{A}\mathbf{P}_k = \lambda_k\mathbf{P}_k$. This motivates the following definition.

652 Definition Let V be a finite-dimensional vector space over a field \mathbb{F} and let $T : V \rightarrow V$ be a linear transformation. A scalar $\lambda \in \mathbb{F}$ is called an *eigenvalue* of T if there is a $\mathbf{v} \neq \mathbf{0}$ (called an *eigenvector*) such that $T(\mathbf{v}) = \lambda\mathbf{v}$.

653 Example Show that if λ is an eigenvalue of $T : V \rightarrow V$, then λ^k is an eigenvalue of $T^k : V \rightarrow V$, for $k \in \mathbb{N} \setminus \{0\}$.

Solution: Assume that $T(\mathbf{v}) = \lambda\mathbf{v}$. Then

$$T^2(\mathbf{v}) = T(T(\mathbf{v})) = T(\lambda\mathbf{v}) = \lambda T(\mathbf{v}) = \lambda(\lambda\mathbf{v}) = \lambda^2\mathbf{v}.$$

Continuing the iterations we obtain $T^k(\mathbf{v}) = \lambda^k\mathbf{v}$, which is what we want.

654 Theorem Let $\mathbf{A} \in M_n(\mathbb{F})$ be the matrix representation of $T : V \rightarrow V$. Then $\lambda \in \mathbb{F}$ is an eigenvalue of T if and only if $\det(\lambda I_n - \mathbf{A}) = 0_{\mathbb{F}}$.

Proof: λ is an eigenvalue of $\mathbf{A} \iff$ there is $\mathbf{v} \neq \mathbf{0}$ such that $\mathbf{A}\mathbf{v} = \lambda\mathbf{v} \iff \lambda\mathbf{v} - \mathbf{A}\mathbf{v} = \mathbf{0} \iff \lambda I_n \mathbf{v} - \mathbf{A}\mathbf{v} = \mathbf{0} \iff \det(\lambda I_n - \mathbf{A}) = 0_{\mathbb{F}}$ by Corollary 648. \square

655 Definition The equation

$$\det(\lambda I_n - \mathbf{A}) = 0_{\mathbb{F}}$$

is called the *characteristic equation of \mathbf{A}* or *secular equation of \mathbf{A}* . The polynomial $p(\lambda) = \det(\lambda I_n - \mathbf{A})$ is the characteristic polynomial of \mathbf{A} .

656 Example Let $\mathbf{A} = \begin{bmatrix} 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 \end{bmatrix}$. Find

① The characteristic polynomial of \mathbf{A} .

② The eigenvalues of \mathbf{A} .

③ The corresponding eigenvectors.

Solution: We have

①

$$\begin{aligned} \det(\lambda \mathbf{I}_4 - \mathbf{A}) &= \det \begin{bmatrix} \lambda - 1 & -1 & 0 & 0 \\ -1 & \lambda - 1 & 0 & 0 \\ 0 & 0 & \lambda - 1 & -1 \\ 0 & 0 & -1 & \lambda - 1 \end{bmatrix} \\ &= (\lambda - 1) \det \begin{bmatrix} \lambda - 1 & 0 & 0 \\ 0 & \lambda - 1 & -1 \\ 0 & -1 & \lambda - 1 \end{bmatrix} + \det \begin{bmatrix} -1 & 0 & 0 \\ 0 & \lambda - 1 & -1 \\ 0 & -1 & \lambda - 1 \end{bmatrix} \\ &= (\lambda - 1)((\lambda - 1)((\lambda - 1)^2 - 1)) + (-((\lambda - 1)^2 - 1)) \\ &= (\lambda - 1)((\lambda - 1)(\lambda - 2)(\lambda)) - (\lambda - 2)(\lambda) \\ &= (\lambda - 2)(\lambda)((\lambda - 1)^2 - 1) \\ &= (\lambda - 2)^2(\lambda)^2 \end{aligned}$$

② The eigenvalues are clearly $\lambda = 0$ and $\lambda = 2$.

③ If $\lambda = 0$, then

$$0\mathbf{I}_4 - \mathbf{A} = \begin{bmatrix} -1 & -1 & 0 & 0 \\ -1 & -1 & 0 & 0 \\ 0 & 0 & -1 & -1 \\ 0 & 0 & -1 & -1 \end{bmatrix}.$$

This matrix has row-echelon form

$$\begin{bmatrix} -1 & -1 & 0 & 0 \\ 0 & 0 & -1 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

and if

$$\begin{bmatrix} -1 & -1 & 0 & 0 \\ 0 & 0 & -1 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix},$$

then $c = -d$ and $a = -b$

Thus the general solution of the system $(0I_4 - A)\mathbf{X} = 0_{n \times 1}$ is

$$\begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = a \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix} + c \begin{bmatrix} 0 \\ 0 \\ 1 \\ -1 \end{bmatrix}.$$

If $\lambda = 2$, then

$$2I_4 - A = \begin{bmatrix} 1 & -1 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & -1 & 1 \end{bmatrix}.$$

This matrix has row-echelon form

$$\begin{bmatrix} -1 & 1 & 0 & 0 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

and if

$$\begin{bmatrix} 1 & -1 & 0 & 0 \\ 0 & 0 & -1 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix},$$

then $c = d$ and $a = b$

Thus the general solution of the system $(2I_4 - A)\mathbf{X} = \mathbf{0}_{n \times 1}$ is

$$\begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = a \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix} + c \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}.$$

Thus for $\lambda = 0$ we have the eigenvectors

$$\begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \\ -1 \end{bmatrix}$$

and for $\lambda = 2$ we have the eigenvectors

$$\begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}$$

657 Theorem If $\lambda = 0_{\mathbb{F}}$ is an eigenvalue of A , then A is non-invertible.

Proof: Put $p(\lambda) = \det(\lambda I_n - A)$. Then $p(0_{\mathbb{F}}) = \det(-A) = (-1)^n \det A$ is the constant term of the characteristic polynomial. If $\lambda = 0_{\mathbb{F}}$ is an eigenvalue then

$$p(0_{\mathbb{F}}) = 0_{\mathbb{F}} \implies \det A = 0_{\mathbb{F}},$$

and hence A is non-invertible by Theorem 647. \square

658 Theorem Similar matrices have the same characteristic polynomial.

Proof: We have

$$\begin{aligned}
 \det(\lambda I_n - SAS^{-1}) &= \det(\lambda SI_n S^{-1} - SAS^{-1}) \\
 &= \det S(\lambda I_n - A)S^{-1} \\
 &= (\det S)(\det(\lambda I_n - A))(\det S^{-1}) \\
 &= (\det S)(\det(\lambda I_n - A)) \left(\frac{1}{\det S} \right) \\
 &= \det(\lambda I_n - A),
 \end{aligned}$$

from where the result follows. \square

659 Problem Find the eigenvalues and eigenvectors of
 $A = \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$

660 Problem Let $A = \begin{bmatrix} 0 & 2 & -1 \\ 2 & 3 & -2 \\ -1 & -2 & 0 \end{bmatrix}$. Find

- ① The characteristic polynomial of A .
- ② The eigenvalues of A .
- ③ The corresponding eigenvectors.

8.3 Diagonalisability

In this section we find conditions for diagonalisability.

661 Theorem Let $\{v_1, v_2, \dots, v_k\} \subset V$ be the eigenvectors corresponding to the *different* eigenvalues $\{\lambda_1, \lambda_2, \dots, \lambda_k\}$ (in that order). Then these eigenvectors are linearly independent.

Proof: Let $T : V \rightarrow V$ be the underlying linear transformation. We proceed by induction. For $k=1$ the result is clear. Assume that every set of $k-1$ eigenvectors that correspond to $k-1$ distinct eigenvalues is linearly independent and let the eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_{k-1}$ have corresponding eigenvectors v_1, v_2, \dots, v_{k-1} . Let λ be a eigenvalue different from the $\lambda_1, \lambda_2, \dots, \lambda_{k-1}$ and let its corresponding eigenvector be v . If v were linearly dependent of the v_1, v_2, \dots, v_{k-1} , we would have

$$xv + x_1v_1 + x_2v_2 + \cdots + x_{k-1}v_{k-1} = 0. \quad (8.1)$$

Now

$$T(xv + x_1v_1 + x_2v_2 + \cdots + x_{k-1}v_{k-1}) = T(0) = 0,$$

by Theorem 538. This implies that

$$x\lambda v + x_1\lambda_1 v_1 + x_2\lambda_2 v_2 + \cdots + x_{k-1}\lambda_{k-1} v_{k-1} = 0. \quad (8.2)$$

From 8.2 take away λ times 8.1, obtaining

$$x_1(\lambda_1 - \lambda)v_1 + x_2(\lambda_2 - \lambda)v_2 + \cdots + x_{k-1}(\lambda_{k-1} - \lambda)v_{k-1} = 0 \quad (8.3)$$

Since $\lambda - \lambda_i \neq 0$ 8.3 is saying that the eigenvectors v_1, v_2, \dots, v_{k-1} are linearly dependent, a contradiction. Thus the claim follows for k distinct eigenvalues and the result is proven by induction. \square

662 Theorem A matrix $A \in M_n(\mathbb{F})$ is diagonalisable if and only if it possesses n linearly independent eigenvectors.

Proof: Assume first that \mathbf{A} is diagonalisable, so there exists $\mathbf{P} \in \mathbf{GL}_n(\mathbb{F})$ and

$$\mathbf{D} = \begin{bmatrix} \lambda_1 & 0 & 0 & \cdots & 0 \\ 0 & \lambda_2 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ 0 & 0 & 0 & \cdots & \lambda_n \end{bmatrix}$$

such that

$$\mathbf{P}^{-1}\mathbf{A}\mathbf{P} = \begin{bmatrix} \lambda_1 & 0 & 0 & \cdots & 0 \\ 0 & \lambda_2 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ 0 & 0 & 0 & \cdots & \lambda_n \end{bmatrix}.$$

Then

$$[\mathbf{AP}_1; \mathbf{AP}_2; \cdots; \mathbf{AP}_n] = \mathbf{AP} = \mathbf{P} \begin{bmatrix} \lambda_1 & 0 & 0 & \cdots & 0 \\ 0 & \lambda_2 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ 0 & 0 & 0 & \cdots & \lambda_n \end{bmatrix} = [\lambda_1 \mathbf{P}_1; \lambda_2 \mathbf{P}_2; \cdots; \lambda_n \mathbf{P}_n],$$

where the \mathbf{P}_k are the columns of \mathbf{P} . Since \mathbf{P} is invertible, the \mathbf{P}_k are linearly independent by virtue of Theorems 473 and 647.

Conversely, suppose now that $\mathbf{v}_1, \dots, \mathbf{v}_n$ are n linearly independent eigenvectors, with corresponding eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$. Put

$$\mathbf{P} = [\mathbf{v}_1; \dots; \mathbf{v}_n], \quad \mathbf{D} = \begin{bmatrix} \lambda_1 & 0 & 0 & \cdots & 0 \\ 0 & \lambda_2 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ 0 & 0 & 0 & \cdots & \lambda_n \end{bmatrix}.$$

Since $\mathbf{Av}_i = \lambda_i \mathbf{v}_i$ we see that $\mathbf{AP} = \mathbf{PD}$. Again \mathbf{P} is invertible by Theorems 473 and 647 since the \mathbf{v}_k are linearly independent. Left multiplying by \mathbf{P}^{-1} we deduce $\mathbf{P}^{-1}\mathbf{AP} = \mathbf{D}$, from where \mathbf{A} is diagonalisable. \square

663 Example Shew that the following matrix is diagonalisable:

$$\begin{bmatrix} 1 & -1 & -1 \\ 1 & 3 & 1 \\ -3 & 1 & -1 \end{bmatrix}$$

and find a diagonal matrix \mathbf{D} and an invertible matrix \mathbf{P} such that

$$\mathbf{A} = \mathbf{P}\mathbf{D}\mathbf{P}^{-1}.$$

Solution: Verify that the characteristic polynomial of \mathbf{A} is

$$\lambda^3 - 3\lambda^2 - 4\lambda + 12 = (\lambda - 2)(\lambda + 2)(\lambda - 3).$$

The eigenvector for $\lambda = -2$ is

$$\begin{bmatrix} 1 \\ -1 \\ 4 \end{bmatrix}.$$

The eigenvector for $\lambda = 2$ is

$$\begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}.$$

The eigenvector for $\lambda = 3$ is

$$\begin{bmatrix} -1 \\ 1 \\ 1 \end{bmatrix}.$$

We may take

$$\mathbf{D} = \begin{bmatrix} -2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}, \quad \mathbf{P} = \begin{bmatrix} 1 & -1 & 4 \\ -1 & 0 & 1 \\ -1 & 1 & 1 \end{bmatrix}.$$

We also find

$$\mathbf{P}^{-1} = \begin{bmatrix} \frac{1}{5} & -1 & \frac{1}{5} \\ 0 & -1 & 1 \\ \frac{1}{5} & 0 & \frac{1}{5} \end{bmatrix}.$$

664 Problem Let \mathbf{A} be a 2×2 matrix with eigenvalues 1 and -2 and corresponding eigenvectors $\begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $\begin{bmatrix} 1 \\ -1 \end{bmatrix}$, respectively. Determine \mathbf{A}^{10} .

One of the eigenvalues has two eigenvectors $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ and $\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$.

665 Problem Let $\mathbf{A} \in M_3(\mathbb{R})$ have characteristic polynomial

$$(\lambda + 1)^2(\lambda - 3).$$

The other eigenvalue has corresponding eigenvector $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$. Determine \mathbf{A} .

666 Problem Let

$$\mathbf{A} = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix}.$$

1. Find $\det \mathbf{A}$.
2. Find \mathbf{A}^{-1} .
3. Find $\text{rank}(\mathbf{A} - \mathbf{I}_4)$.
4. Find $\det(\mathbf{A} - \mathbf{I}_4)$.
5. Find the characteristic polynomial of \mathbf{A} .
6. Find the eigenvalues of \mathbf{A} .
7. Find the eigenvectors of \mathbf{A} .
8. Find \mathbf{A}^{10} .

667 Problem Let $\mathbf{U} \in M_n(\mathbb{R})$ be a square matrix all whose entries are equal to 1.

1. Demonstrate that $\mathbf{U}^2 = n\mathbf{U}$.
2. Find $\det \mathbf{U}$.
3. Prove that $\det(\lambda\mathbf{I}_n - \mathbf{U}) = \lambda^{n-1}(\lambda - n)$.
4. Shew that $\dim \ker(\mathbf{U}) = n - 1$.
5. Shew that

$$\mathbf{U} = \mathbf{P} \begin{bmatrix} n & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \end{bmatrix} \mathbf{P}^{-1},$$

where

$$\mathbf{P} = \begin{bmatrix} 1 & 1 & 0 & \cdots & 0 & 0 \\ 1 & 0 & 1 & \cdots & 0 & 0 \\ 1 & 0 & 0 & \ddots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 1 & 0 & 0 & \cdots & 0 & 1 \\ 1 & -1 & -1 & \cdots & -1 & -1 \end{bmatrix}.$$

8.4 The Minimal Polynomial

668 Theorem (Cayley-Hamilton) A matrix $\mathbf{A} \in M_n(\mathbb{F})$ satisfies its characteristic polynomial.

Proof: Put $\mathbf{B} = \lambda\mathbf{I}_n - \mathbf{A}$. We can write

$$\det \mathbf{B} = \det(\lambda\mathbf{I}_n - \mathbf{A}) = \lambda^n + b_1\lambda^{n-1} + b_2\lambda^{n-2} + \cdots + b_n,$$

as $\det(\lambda\mathbf{I}_n - \mathbf{A})$ is a polynomial of degree n .

Since $\text{adj}(\mathbf{B})$ is a matrix obtained by using $(n-1) \times (n-1)$ determinants from \mathbf{B} , we may write

$$\text{adj}(\mathbf{B}) = \lambda^{n-1}\mathbf{B}_{n-1} + \lambda^{n-2}\mathbf{B}_{n-2} + \cdots + \mathbf{B}_0.$$

Hence

$$\det(\lambda\mathbf{I}_n - \mathbf{A})\mathbf{I}_n = (\mathbf{B})(\text{adj}(\mathbf{B})) = (\lambda\mathbf{I}_n - \mathbf{A})(\text{adj}(\mathbf{B})),$$

from where

$$\lambda^n\mathbf{I}_n + b_1\mathbf{I}_n\lambda^{n-1} + b_2\mathbf{I}_n\lambda^{n-2} + \cdots + b_n\mathbf{I}_n = (\lambda\mathbf{I}_n - \mathbf{A})(\lambda^{n-1}\mathbf{B}_{n-1} + \lambda^{n-2}\mathbf{B}_{n-2} + \cdots + \mathbf{B}_0).$$

By equating coefficients we deduce

$$\begin{aligned}
 I_n &= B_{n-1} \\
 b_1 I_n &= -AB_{n-1} + B_{n-2} \\
 b_2 I_n &= -AB_{n-2} + B_{n-3} \\
 &\vdots \\
 b_{n-1} I_n &= -AB_1 + B_0 \\
 b_n I_n &= -AB_0.
 \end{aligned}$$

Multiply now the k -th row by A^{n-k} (the first row appearing is really the 0-th row):

$$\begin{aligned}
 A^n &= A^n B_{n-1} \\
 b_1 A^{n-1} &= -A^n B_{n-1} + A^{n-1} B_{n-2} \\
 b_2 A^{n-2} &= -A^{n-1} B_{n-2} + A^{n-2} B_{n-3} \\
 &\vdots \\
 b_{n-1} A &= -A^2 B_1 + AB_0 \\
 b_n I_n &= -AB_0.
 \end{aligned}$$

Add all the rows and through telescopic cancellation obtain

$$A^n + b_1 A^{n-1} + \cdots + b_{n-1} A + b_n I_n = 0_n,$$

from where the theorem follows. \square

669 Example From example 663 the matrix

$$\begin{bmatrix} 1 & -1 & -1 \\ 1 & 3 & 1 \\ -3 & 1 & -1 \end{bmatrix}$$

has characteristic polynomial

$$(\lambda - 3)(\lambda - 2)(\lambda + 2) = \lambda^3 - 3\lambda^2 - 4\lambda + 12,$$

hence the inverse of this matrix can be obtained by observing that

$$A^3 - 3A^2 - 4A + 12I_3 = 0_3 \implies A^{-1} = -\frac{1}{12}(A^2 - 3A - 4I_3) = \begin{bmatrix} 1/3 & 1/6 & -1/6 \\ 1/6 & 1/3 & 1/6 \\ -5/6 & -1/6 & -1/3 \end{bmatrix}.$$

Having seen that the characteristic polynomial of a square matrix A annihilates A , the question now arises of whether there exists a polynomial of minimal degree annihilating A .

Appendix A

Answers and Hints

17

$$\begin{aligned} x \in X \setminus (X \setminus A) &\iff x \in X \wedge x \notin (X \setminus A) \\ &\iff x \in X \wedge x \in A \\ &\iff x \in X \cap A. \end{aligned}$$

18

$$\begin{aligned} X \setminus (A \cup B) &\iff x \in X \wedge (x \notin (A \cup B)) \\ &\iff x \in X \wedge (x \notin A \wedge x \notin B) \\ &\iff (x \in X \wedge x \notin A) \wedge (x \in X \wedge x \notin B) \\ &\iff x \in (X \setminus A) \wedge x \in (X \setminus B) \\ &\iff x \in (X \setminus A) \cap (X \setminus B). \end{aligned}$$

21 One possible solution is

$$A \cup B \cup C = A \cup (B \setminus A) \cup (C \setminus (A \cup B)).$$

23 We have

$$|a| = |a - b + b| \leq |a - b| + |b|,$$

giving

$$|a| - |b| \leq |a - b|.$$

Similarly,

$$|b| = |b - a + a| \leq |b - a| + |a| = |a - b| + |a|,$$

gives

$$|b| - |a| \leq |a - b|.$$

The stated inequality follows from this.

38 $a \sim a$ since $\frac{a}{a} = 1 \in \mathbb{Z}$, and so the relation is reflexive. The relation is not symmetric. For $2 \sim 1$ since $\frac{2}{1} \in \mathbb{Z}$ but $1 \not\sim 2$ since $\frac{1}{2} \notin \mathbb{Z}$. The relation is transitive. For assume $a \sim b$ and $b \sim c$. Then there exist $(m, n) \in \mathbb{Z}^2$ such that $\frac{a}{b} = m$, $\frac{b}{c} = n$. This gives

$$\frac{a}{c} = \frac{a}{b} \cdot \frac{b}{c} = mn \in \mathbb{Z},$$

and so $a \sim c$.

39 Here is one possible example: put $a \sim b \Leftrightarrow \frac{a^2+a}{b} \in \mathbb{Z}$. Then clearly if $a \in \mathbb{Z} \setminus \{0\}$ we have $a \sim a$ since $\frac{a^2+a}{a} = a+1 \in \mathbb{Z}$. On the other hand, the relation is not symmetric, since $5 \sim 2$ as $\frac{5^2+5}{2} = 15 \in \mathbb{Z}$ but $2 \not\sim 5$, as $\frac{2^2+2}{5} = \frac{6}{5} \notin \mathbb{Z}$. It is not transitive either, since $\frac{5^2+5}{3} \in \mathbb{Z} \Rightarrow 5 \sim 3$ and $\frac{3^2+3}{12} \in \mathbb{Z} \Rightarrow 3 \sim 12$ but $\frac{5^2+5}{12} \notin \mathbb{Z}$ and so $5 \not\sim 12$.

41 [B] $[x] = x + \frac{1}{3}\mathbb{Z}$. [C] No.

55 Let $\omega = -\frac{1}{2} + i\frac{\sqrt{3}}{2}$. Then $\omega^2 + \omega + 1 = 0$ and $\omega^3 = 1$. Then

$$x = a^3 + b^3 + c^3 - 3abc = (a+b+c)(a+\omega b + \omega^2 c)(a+\omega^2 b + c\omega),$$

$$y = u^3 + v^3 + w^3 - 3uvw = (u+v+w)(u+\omega v + \omega^2 w)(u+\omega^2 v + \omega w).$$

Then

$$\begin{aligned} (a+b+c)(u+v+w) &= au + av + aw + bu + bv + bw + cu + cv + cw, \\ (a+\omega b + \omega^2 c)(u+\omega v + \omega^2 w) &= au + bw + cv \\ &\quad + \omega(av + bu + cw) \\ &\quad + \omega^2(aw + bv + cu), \end{aligned}$$

and

$$\begin{aligned} (a+\omega^2 b + \omega c)(u+\omega^2 v + \omega w) &= au + bw + cv \\ &\quad + \omega(aw + bv + cu) \\ &\quad + \omega^2(av + bu + cw). \end{aligned}$$

This proves that

$$\begin{aligned} xy &= (au + bw + cv)^3 + (aw + bv + cu)^3 + (av + bu + cw)^3 \\ &\quad - 3(au + bw + cv)(aw + bv + cu)(av + bu + cw), \end{aligned}$$

which proves that S is closed under multiplication.

56 We have

$$xT(yTz) = xT(y \otimes a \otimes z) = (x) \otimes (a) \otimes (y \otimes a \otimes z) = x \otimes a \otimes y \otimes a \otimes z,$$

where we may drop the parentheses since \otimes is associative. Similarly

$$(xT y)Tz = (x \otimes a \otimes y)Tz = (x \otimes a \otimes y) \otimes (a) \otimes (z) = x \otimes a \otimes y \otimes a \otimes z.$$

By virtue of having proved

$$xT(yTz) = (xT y)Tz,$$

associativity is established.

57 We proceed in order.

❶ Clearly, if a, b are rational numbers,

$$|a| < 1, |b| < 1 \implies |ab| < 1 \implies -1 < ab < 1 \implies 1 + ab > 0,$$

whence the denominator never vanishes and since sums, multiplications and divisions of rational numbers are rational, $\frac{a+b}{1+ab}$ is also rational. We must prove now that $-1 < \frac{a+b}{1+ab} < 1$ for $(a, b) \in [-1; 1]^2$. We have

$$\begin{aligned} -1 < \frac{a+b}{1+ab} < 1 &\Leftrightarrow -1 - ab < a + b < 1 + ab \\ &\Leftrightarrow -1 - ab - a - b < 0 < 1 + ab - a - b \\ &\Leftrightarrow -(a+1)(b+1) < 0 < (a-1)(b-1). \end{aligned}$$

Since $(a, b) \in [-1; 1]^2$, $(a+1)(b+1) > 0$ and so $-(a+1)(b+1) < 0$ giving the sinistral inequality. Similarly $a-1 < 0$ and $b-1 < 0$ give $(a-1)(b-1) > 0$, the dextral inequality. Since the steps are reversible, we have established that indeed $-1 < \frac{a+b}{1+ab} < 1$.

- ② Since $a \otimes b = \frac{a+b}{1+ab} = \frac{b+a}{1+ba} = b \otimes a$, commutativity follows trivially. Now

$$\begin{aligned} a \otimes (b \otimes c) &= a \otimes \left(\frac{b+c}{1+bc} \right) \\ &= \frac{a + \left(\frac{b+c}{1+bc} \right)}{1 + a \left(\frac{b+c}{1+bc} \right)} \\ &= \frac{a(1+bc) + b + c}{1 + bc + a(b + c)} = \frac{a + b + c + abc}{1 + ab + bc + ca}. \end{aligned}$$

One the other hand,

$$\begin{aligned} (a \otimes b) \otimes c &= \left(\frac{a+b}{1+ab} \right) \otimes c \\ &= \frac{\left(\frac{a+b}{1+ab} \right) + c}{1 + \left(\frac{a+b}{1+ab} \right) c} \\ &= \frac{(a+b) + c(1+ab)}{1 + ab + (a+b)c} \\ &= \frac{a + b + c + abc}{1 + ab + bc + ca}, \end{aligned}$$

whence \otimes is associative.

- ③ If $a \otimes e = a$ then $\frac{a+e}{1+ae} = a$, which gives $a+e = a+ea^2$ or $e(a^2-1)=0$. Since $a \neq \pm 1$, we must have $e=0$.
- ④ If $a \otimes b = 0$, then $\frac{a+b}{1+ab} = 0$, which means that $b=-a$.

58 We proceed in order.

- ① Since $a \otimes b = a + b - ab = b + a - ba = b \otimes a$, commutativity follows trivially. Now

$$\begin{aligned} a \otimes (b \otimes c) &= a \otimes (b + c - bc) \\ &= a + b + c - bc - a(b + c - bc) \\ &= a + b + c - ab - bc - ca + abc. \end{aligned}$$

One the other hand,

$$\begin{aligned} (a \otimes b) \otimes c &= (a + b - ab) \otimes c \\ &= a + b - ab + c - (a + b - ab)c \\ &= a + b + c - ab - bc - ca + abc, \end{aligned}$$

whence \otimes is associative.

- ② If $a \otimes e = a$ then $a + e - ae = a$, which gives $e(1-a)=0$. Since $a \neq 1$, we must have $e=0$.

- ③ If $a \otimes b = 0$, then $a + b - ab = 0$, which means that $b(1-a) = -a$. Since $a \neq 1$ we find $b = -\frac{a}{1-a}$.

$+$	$\bar{0}$	$\bar{1}$	$\bar{2}$	$\bar{3}$	$\bar{4}$	$\bar{5}$	$\bar{6}$	$\bar{7}$	$\bar{8}$	$\bar{9}$	$\bar{10}$
$\bar{0}$	$\bar{0}$	$\bar{1}$	$\bar{2}$	$\bar{3}$	$\bar{4}$	$\bar{5}$	$\bar{6}$	$\bar{7}$	$\bar{8}$	$\bar{9}$	$\bar{10}$
$\bar{1}$	$\bar{1}$	$\bar{2}$	$\bar{3}$	$\bar{4}$	$\bar{5}$	$\bar{6}$	$\bar{7}$	$\bar{8}$	$\bar{9}$	$\bar{10}$	$\bar{0}$
$\bar{2}$	$\bar{2}$	$\bar{3}$	$\bar{4}$	$\bar{5}$	$\bar{6}$	$\bar{7}$	$\bar{8}$	$\bar{9}$	$\bar{10}$	$\bar{0}$	$\bar{1}$
$\bar{3}$	$\bar{3}$	$\bar{4}$	$\bar{5}$	$\bar{6}$	$\bar{7}$	$\bar{8}$	$\bar{9}$	$\bar{10}$	$\bar{0}$	$\bar{1}$	$\bar{2}$
$\bar{4}$	$\bar{4}$	$\bar{5}$	$\bar{6}$	$\bar{7}$	$\bar{8}$	$\bar{9}$	$\bar{10}$	$\bar{0}$	$\bar{1}$	$\bar{2}$	$\bar{3}$
$\bar{5}$	$\bar{5}$	$\bar{6}$	$\bar{7}$	$\bar{8}$	$\bar{9}$	$\bar{10}$	$\bar{0}$	$\bar{1}$	$\bar{2}$	$\bar{3}$	$\bar{4}$
$\bar{6}$	$\bar{6}$	$\bar{7}$	$\bar{8}$	$\bar{9}$	$\bar{10}$	$\bar{0}$	$\bar{1}$	$\bar{2}$	$\bar{3}$	$\bar{4}$	$\bar{5}$
$\bar{7}$	$\bar{7}$	$\bar{8}$	$\bar{9}$	$\bar{10}$	$\bar{0}$	$\bar{1}$	$\bar{2}$	$\bar{3}$	$\bar{4}$	$\bar{5}$	$\bar{6}$
$\bar{8}$	$\bar{8}$	$\bar{9}$	$\bar{10}$	$\bar{0}$	$\bar{1}$	$\bar{2}$	$\bar{3}$	$\bar{4}$	$\bar{5}$	$\bar{6}$	$\bar{7}$
$\bar{9}$	$\bar{9}$	$\bar{10}$	$\bar{0}$	$\bar{1}$	$\bar{2}$	$\bar{3}$	$\bar{4}$	$\bar{5}$	$\bar{6}$	$\bar{7}$	$\bar{8}$
$\bar{10}$	$\bar{10}$	$\bar{0}$	$\bar{1}$	$\bar{2}$	$\bar{3}$	$\bar{4}$	$\bar{5}$	$\bar{6}$	$\bar{7}$	$\bar{8}$	$\bar{9}$

Table A.1: Addition table for \mathbb{Z}_{11} .

.	$\bar{0}$	$\bar{1}$	$\bar{2}$	$\bar{3}$	$\bar{4}$	$\bar{5}$	$\bar{6}$	$\bar{7}$	$\bar{8}$	$\bar{9}$	$\bar{10}$
$\bar{0}$	$\bar{0}$	$\bar{0}$	$\bar{0}$	$\bar{0}$	$\bar{0}$	$\bar{0}$	$\bar{0}$	$\bar{0}$	$\bar{0}$	$\bar{0}$	$\bar{0}$
$\bar{1}$	$\bar{0}$	$\bar{1}$	$\bar{2}$	$\bar{3}$	$\bar{4}$	$\bar{5}$	$\bar{6}$	$\bar{7}$	$\bar{8}$	$\bar{9}$	$\bar{10}$
$\bar{2}$	$\bar{0}$	$\bar{2}$	$\bar{4}$	$\bar{6}$	$\bar{8}$	$\bar{10}$	$\bar{1}$	$\bar{3}$	$\bar{5}$	$\bar{7}$	$\bar{9}$
$\bar{3}$	$\bar{0}$	$\bar{3}$	$\bar{6}$	$\bar{9}$	$\bar{1}$	$\bar{4}$	$\bar{7}$	$\bar{10}$	$\bar{2}$	$\bar{5}$	$\bar{8}$
$\bar{4}$	$\bar{0}$	$\bar{4}$	$\bar{8}$	$\bar{1}$	$\bar{5}$	$\bar{9}$	$\bar{2}$	$\bar{6}$	$\bar{10}$	$\bar{3}$	$\bar{7}$
$\bar{5}$	$\bar{0}$	$\bar{5}$	$\bar{10}$	$\bar{4}$	$\bar{9}$	$\bar{3}$	$\bar{8}$	$\bar{2}$	$\bar{7}$	$\bar{1}$	$\bar{6}$
$\bar{6}$	$\bar{0}$	$\bar{6}$	$\bar{7}$	$\bar{2}$	$\bar{8}$	$\bar{3}$	$\bar{9}$	$\bar{4}$	$\bar{10}$	$\bar{5}$	$\bar{1}$
$\bar{7}$	$\bar{0}$	$\bar{7}$	$\bar{3}$	$\bar{10}$	$\bar{6}$	$\bar{2}$	$\bar{9}$	$\bar{5}$	$\bar{1}$	$\bar{8}$	$\bar{4}$
$\bar{8}$	$\bar{0}$	$\bar{8}$	$\bar{5}$	$\bar{2}$	$\bar{10}$	$\bar{7}$	$\bar{4}$	$\bar{1}$	$\bar{9}$	$\bar{6}$	$\bar{3}$
$\bar{9}$	$\bar{0}$	$\bar{9}$	$\bar{7}$	$\bar{5}$	$\bar{3}$	$\bar{1}$	$\bar{10}$	$\bar{8}$	$\bar{6}$	$\bar{4}$	$\bar{2}$
$\bar{10}$	$\bar{0}$	$\bar{10}$	$\bar{8}$	$\bar{6}$	$\bar{4}$	$\bar{2}$	$\bar{5}$	$\bar{3}$	$\bar{1}$	$\bar{9}$	$\bar{7}$

Table A.2: Multiplication table \mathbb{Z}_{11} .

59 We have

$$\begin{aligned}
 x \circ y &= (x \circ y) \circ (x \circ y) \\
 &= [y \circ (x \circ y)] \circ x \\
 &= [(x \circ y) \circ x] \circ y \\
 &= [(y \circ x) \circ x] \circ y \\
 &= [(x \circ x) \circ y] \circ y \\
 &= (y \circ y) \circ (x \circ x) \\
 &= y \circ x,
 \end{aligned}$$

proving commutativity.

71 The tables appear in tables A.1 and A.2.

72 From example 68

$$x^2 = \bar{5}.$$

Now, the squares modulo 11 are $\bar{0}^2 = \bar{0}$, $\bar{1}^2 = \bar{1}$, $\bar{2}^2 = \bar{4}$, $\bar{3}^2 = \bar{9}$, $\bar{4}^2 = \bar{5}$, $\bar{5}^2 = \bar{3}$. Also, $(\bar{11} - \bar{4})^2 = \bar{7}^2 = \bar{5}$. Hence the solutions are $x = \bar{4}$ or $x = \bar{7}$.

81 We have

$$\begin{aligned}
 \frac{1}{\sqrt{2} + 2\sqrt{3} + 3\sqrt{6}} &= \frac{\sqrt{2} + 2\sqrt{3} - 3\sqrt{6}}{(\sqrt{2} + 2\sqrt{3})^2 - (3\sqrt{6})^2} \\
 &= \frac{\sqrt{2} + 2\sqrt{3} - 3\sqrt{6}}{2 + 12 + 4\sqrt{6} - 54} \\
 &= \frac{\sqrt{2} + 2\sqrt{3} - 3\sqrt{6}}{-40 + 4\sqrt{6}} \\
 &= \frac{(\sqrt{2} + 2\sqrt{3} - 3\sqrt{6})(-40 - 4\sqrt{6})}{40^2 - (4\sqrt{6})^2} \\
 &= \frac{(\sqrt{2} + 2\sqrt{3} - 3\sqrt{6})(-40 - 4\sqrt{6})}{1504} \\
 &= -\frac{16\sqrt{2} + 22\sqrt{3} - 30\sqrt{6} - 18}{376}
 \end{aligned}$$

82 Since

$$(-a)b^{-1} + ab^{-1} = (-a + a)b^{-1} = 0_F b^{-1} = 0_F,$$

we obtain by adding $-(ab^{-1})$ to both sides that

$$(-a)b^{-1} = -(ab^{-1}).$$

Similarly, from

$$a(-b^{-1}) + ab^{-1} = a(-b^{-1} + b^{-1}) = a0_F = 0_F,$$

we obtain by adding $-(ab^{-1})$ to both sides that

$$a(-b^{-1}) = -(ab^{-1}).$$

93 Assume $h(b) = h(a)$. Then

$$\begin{aligned} h(a) &= h(b) \implies a^3 = b^3 \\ &\implies a^3 - b^3 = 0 \\ &\implies (a - b)(a^2 + ab + b^2) = 0 \end{aligned}$$

Now,

$$b^2 + ab + a^2 = \left(b + \frac{a}{2}\right)^2 + \frac{3a^2}{4}.$$

This shews that $b^2 + ab + a^2$ is positive unless both a and b are zero. Hence $a - b = 0$ in all cases. We have shewn that $h(b) = h(a) \implies a = b$, and the function is thus injective.

94 We have

$$\begin{aligned} f(a) = f(b) &\iff \frac{6a}{2a-3} = \frac{6b}{2b-3} \\ &\iff 6a(2b-3) = 6b(2a-3) \\ &\iff 12ab - 18a = 12ab - 18b \\ &\iff -18a = -18b \\ &\iff a = b, \end{aligned}$$

proving that f is injective. Now, if

$$f(x) = y, \quad y \neq 3,$$

then

$$\frac{6x}{2x-3} = y,$$

that is $6x = y(2x - 3)$. Solving for x we find

$$x = \frac{3y}{2y-6}.$$

Since $2y - 6 \neq 0$, x is a real number, and so f is surjective. On combining the results we deduce that f is bijective.

$$\text{104 } A = \begin{bmatrix} 1 & 1 & 1 \\ 2 & 4 & 8 \\ 3 & 9 & 27 \end{bmatrix}.$$

$$\text{105 } A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \\ 3 & 6 & 9 \end{bmatrix}.$$

$$106 \quad M + N = \begin{bmatrix} a+1 & 0 & 2c \\ a & b-2a & 0 \\ 2a & 0 & -2 \end{bmatrix}, \quad 2M = \begin{bmatrix} 2a & -4a & 2c \\ 0 & -2a & 2b \\ 2a+2b & 0 & -2 \end{bmatrix}.$$

107 $x = 1$ and $y = 4$.

$$108 \quad A = \begin{bmatrix} 13 & -1 \\ 15 & 3 \end{bmatrix}, \quad B = \begin{bmatrix} 5 & 0 \\ 6 & 1 \end{bmatrix}$$

111 The set of border elements is the union of two rows and two columns. Thus we may choose at most four elements from the border, and at least one from the central 3×3 matrix. The largest element of this 3×3 matrix is 15, so any allowable choice of does not exceed 15. The choice 25, 15, 18, 1, 23, 20 shews that the largest minimum is indeed 15.

$$120 \quad \begin{bmatrix} 2 & 2 \\ 0 & -2 \end{bmatrix}$$

121

$$AB = \begin{bmatrix} a & b & c \\ c+a & a+b & b+c \\ a+b+c & a+b+c & a+b+c \end{bmatrix}, \quad BA = \begin{bmatrix} a+b+c & b+c & c \\ a+b+c & a+b & b \\ a+b+c & c+a & a \end{bmatrix}$$

$$122 \quad AB = 0_4 \text{ and } BA = \begin{bmatrix} \bar{0} & \bar{2} & \bar{0} & \bar{3} \\ \bar{0} & \bar{2} & \bar{0} & \bar{3} \\ \bar{0} & \bar{2} & \bar{0} & \bar{3} \\ \bar{0} & \bar{2} & \bar{0} & \bar{3} \end{bmatrix}.$$

123 Observe that

$$\begin{bmatrix} -4 & x \\ -x & 4 \end{bmatrix}^2 = \begin{bmatrix} -4 & x \\ -x & 4 \end{bmatrix} \begin{bmatrix} -4 & x \\ -x & 4 \end{bmatrix} = \begin{bmatrix} 16-x^2 & 0 \\ 0 & 16-x^2 \end{bmatrix},$$

and so we must have $16-x^2 = -1$ or $x = \pm\sqrt{17}$.

$$124 \quad \text{Disprove! Take } A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \text{ and } B = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}. \text{ Then } AB = B, \text{ but } BA = 0_2.$$

$$125 \quad \text{Disprove! Take for example } A = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix} \text{ and } B = \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}. \text{ Then}$$

$$A^2 - B^2 = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \neq \begin{bmatrix} -1 & 0 \\ -2 & 1 \end{bmatrix} = (A+B)(A-B).$$

127 $\begin{bmatrix} 32 & -32 \\ -32 & 32 \end{bmatrix}.$

128 $A^{2003} = \begin{bmatrix} 0 & 2^{1001}3^{1002} \\ 2^{1002}3^{1001} & 0 \end{bmatrix}.$

130 The assertion is clearly true for $n = 1$. Assume that is it true for n , that is, assume

$$A^n = \begin{bmatrix} \cos(n)\alpha & -\sin(n)\alpha \\ \sin(n)\alpha & \cos(n)\alpha \end{bmatrix}.$$

Then

$$\begin{aligned} A^{n+1} &= AA^n \\ &= \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix} \begin{bmatrix} \cos(n)\alpha & -\sin(n)\alpha \\ \sin(n)\alpha & \cos(n)\alpha \end{bmatrix} \\ &= \begin{bmatrix} \cos \alpha \cos(n)\alpha - \sin \alpha \sin(n)\alpha & -\cos \alpha \sin(n)\alpha - \sin \alpha \cos(n)\alpha \\ \sin \alpha \cos(n)\alpha + \cos \alpha \sin(n)\alpha & -\sin \alpha \sin(n)\alpha + \cos \alpha \cos(n)\alpha \end{bmatrix} \\ &= \begin{bmatrix} \cos(n+1)\alpha & -\sin(n+1)\alpha \\ \sin(n+1)\alpha & \cos(n+1)\alpha \end{bmatrix}, \end{aligned}$$

and the result follows by induction.

131 Let $A = [a_{ij}]$, $B = [b_{ij}]$ be checkered $n \times n$ matrices. Then $A + B = (a_{ij} + b_{ij})$. If $j - i$ is odd, then $a_{ij} + b_{ij} = 0 + 0 = 0$, which shows that $A + B$ is checkered. Furthermore, let $AB = [c_{ij}]$ with $c_{ij} = \sum_{k=1}^n a_{ik}b_{kj}$. If i is even and j odd, then $a_{ik} = 0$ for odd k and $b_{kj} = 0$ for even k . Thus $c_{ij} = 0$ for i even and j odd. Similarly, $c_{ij} = 0$ for odd i and even j . This proves that AB is checkered.

132 Put

$$J = \begin{bmatrix} 0 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}.$$

We first notice that

$$J^2 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad J^3 = 0_3.$$

This means that the sum in the binomial expansion

$$A^n = (I_3 + J)^n = \sum_{k=0}^n \binom{n}{k} I^{n-k} J^k$$

is a sum of zero matrices for $k \geq 3$. We thus have

$$\begin{aligned} \mathbf{A}^n &= \mathbf{I}_3^n + n\mathbf{I}_3^{n-1}\mathbf{J} + \binom{n}{2}\mathbf{I}_3^{n-2}\mathbf{J}^2 \\ &= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & n & n \\ 0 & 0 & n \\ 0 & 0 & 0 \end{bmatrix} + \begin{bmatrix} 0 & 0 & \binom{n}{2} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 1 & n & \frac{n(n+1)}{2} \\ 0 & 1 & n \\ 0 & 0 & 1 \end{bmatrix}, \end{aligned}$$

giving the result, since $\binom{n}{2} = \frac{n(n-1)}{2}$ and $n + \binom{n}{2} = \frac{n(n+1)}{2}$.

134 Argue inductively,

$$\mathbf{A}^2\mathbf{B} = \mathbf{A}(\mathbf{AB}) = \mathbf{AB} = \mathbf{B}$$

$$\mathbf{A}^3\mathbf{B} = \mathbf{A}(\mathbf{A}^2\mathbf{B}) = \mathbf{A}(\mathbf{AB}) = \mathbf{AB} = \mathbf{B}$$

⋮

$$\mathbf{A}^m\mathbf{B} = \mathbf{AB} = \mathbf{B}.$$

Hence $\mathbf{B} = \mathbf{A}^m\mathbf{B} = \mathbf{0}_n\mathbf{B} = \mathbf{0}_n$.

136 Put $\mathbf{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Using 135, deduce by iteration that

$$\mathbf{A}^k = (a + d)^{k-1}\mathbf{A}.$$

137 $\begin{bmatrix} a & b \\ c & -a \end{bmatrix}$, $bc = -a^2$

138 $\pm \mathbf{I}_2$, $\begin{bmatrix} a & b \\ c & -a \end{bmatrix}$, $a^2 = 1 - bc$

139 We complete squares by putting $\mathbf{Y} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \mathbf{X} - \mathbf{I}$. Then

$$\begin{bmatrix} a^2 + bc & b(a+d) \\ c(a+d) & bc + d^2 \end{bmatrix} = \mathbf{Y}^2 = \mathbf{X}^2 - 2\mathbf{X} + \mathbf{I} = (\mathbf{X} - \mathbf{I})^2 = \begin{bmatrix} -1 & 0 \\ 6 & 3 \end{bmatrix} + \mathbf{I} = \begin{bmatrix} 0 & 0 \\ 6 & 4 \end{bmatrix}.$$

This entails $a = 0, b = 0, cd = 6, d^2 = 4$. Using $\mathbf{X} = \mathbf{Y} + \mathbf{I}$, we find that there are two solutions,

$$\begin{bmatrix} 1 & 0 \\ 3 & 3 \end{bmatrix}, \quad \begin{bmatrix} 1 & 0 \\ -3 & -1 \end{bmatrix}.$$

150 There are infinitely many solutions. Here is one:

$$\mathbf{A} = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 3 & 1 & 2 \end{bmatrix} = \begin{bmatrix} -9 & 2 & 3 \\ 2 & 3 & 1 \\ 3 & 1 & 2 \end{bmatrix} + \begin{bmatrix} 10 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

151 If such matrices existed, then by the first equation

$$\text{tr}(\mathbf{AC}) + \text{tr}(\mathbf{DB}) = n.$$

By the second equation and by Theorem 141,

$$0 = \text{tr}(\mathbf{CA}) + \text{tr}(\mathbf{BD}) = \text{tr}(\mathbf{AC}) + \text{tr}(\mathbf{DB}) = n,$$

a contradiction, since $n \geq 1$.

152 Disprove! This is not generally true. Take $\mathbf{A} = \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix}$ and $\mathbf{B} = \begin{bmatrix} 3 & 0 \\ 0 & 1 \end{bmatrix}$. Clearly $\mathbf{A}^T = \mathbf{A}$ and $\mathbf{B}^T = \mathbf{B}$. We have

$$\mathbf{AB} = \begin{bmatrix} 3 & 1 \\ 3 & 2 \end{bmatrix}$$

but

$$(\mathbf{AB})^T = \begin{bmatrix} 3 & 3 \\ 1 & 2 \end{bmatrix}.$$

154 We have

$$\text{tr}(\mathbf{A}^2) = \text{tr}\left(\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix}\right) = \text{tr}\left(\begin{bmatrix} a^2 + bc & ab + bd \\ ca + cd & d^2 + cb \end{bmatrix}\right) = a^2 + d^2 + 2bc$$

and

$$\left(\text{tr}\left(\begin{bmatrix} a & b \\ c & d \end{bmatrix}\right)\right)^2 = (a + d)^2.$$

Thus

$$\text{tr}(\mathbf{A}^2) = (\text{tr}(\mathbf{A}))^2 \iff a^2 + d^2 + 2bc = (a + d)^2 \iff bc = ad,$$

is the condition sought.

155

$$\begin{aligned} \text{tr}((\mathbf{A} - \mathbf{I}_4)^2) &= \text{tr}(\mathbf{A}^2 - 2\mathbf{A} + \mathbf{I}_4) \\ &= \text{tr}(\mathbf{A}^2) - 2\text{tr}(\mathbf{A}) + \text{tr}(\mathbf{I}_4) \\ &= -4 - 2\text{tr}(\mathbf{A}) + 4 \\ &= -2\text{tr}(\mathbf{A}), \end{aligned}$$

and $\text{tr}(3\mathbf{I}_4) = 12$. Hence $-2\text{tr}(\mathbf{A}) = 12$ or $\text{tr}(\mathbf{A}) = -6$.

156 Disprove! Take $\mathbf{A} = \mathbf{B} = \mathbf{I}_n$ and $n > 1$. Then $\text{tr}(\mathbf{AB}) = n < n^2 = \text{tr}(\mathbf{A})\text{tr}(\mathbf{B})$.

157 Disprove! Take $\mathbf{A} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$, $\mathbf{B} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$, $\mathbf{C} = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$. Then $\text{tr}(\mathbf{ABC}) = 1$ but $\text{tr}(\mathbf{BAC}) = 0$.

158 We have

$$(\mathbf{AA}^T)^T = (\mathbf{A}^T)^T \mathbf{A}^T = \mathbf{AA}^T.$$

159 We have

$$(\mathbf{AB} - \mathbf{BA})^T = (\mathbf{AB})^T - (\mathbf{BA})^T = \mathbf{B}^T \mathbf{A}^T - \mathbf{A}^T \mathbf{B}^T = -\mathbf{BA} - \mathbf{A}(-\mathbf{B}) = \mathbf{AB} - \mathbf{BA}.$$

161 Let $\mathbf{X} = [x_{ij}]$ and put $\mathbf{XX}^T = [\mathbf{c}_{ij}]$. Then

$$0 = \mathbf{c}_{ii} = \sum_{k=1}^n x_{ik}^2 \implies x_{ik} = 0.$$

187 Here is one possible approach. If we perform $C_1 \leftrightarrow C_3$ on \mathbf{A} we obtain

$$\mathbf{A}_1 = \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 1 & -1 & 1 \\ 1 & -1 & 1 & 1 \end{bmatrix} \quad \text{so take } \mathbf{P} = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Now perform $2R_1 \rightarrow R_1$ on \mathbf{A}_1 to obtain

$$\mathbf{A}_2 = \begin{bmatrix} 2 & 0 & 2 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 1 & -1 & 1 \\ 1 & -1 & 1 & 1 \end{bmatrix} \quad \text{so take } \mathbf{D} = \begin{bmatrix} 2 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Finally, perform $R_1 + 2R_4 \rightarrow R_1$ on \mathbf{A}_2 to obtain

$$\mathbf{B} = \begin{bmatrix} 4 & -2 & 4 & 2 \\ 0 & 1 & 0 & 1 \\ 1 & 1 & -1 & 1 \\ 1 & -1 & 1 & 1 \end{bmatrix} \quad \text{so take } \mathbf{T} = \begin{bmatrix} 1 & 0 & 0 & 2 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

188 Here is one possible approach.

$$\begin{array}{ccc}
 \left[\begin{array}{ccc} a & b & c \\ d & e & f \\ g & h & i \end{array} \right] & \xrightarrow{\text{P: } \rho_3 \leftrightarrow \rho_1} & \left[\begin{array}{ccc} g & h & i \\ d & e & f \\ a & b & c \end{array} \right] \\
 & \xrightarrow{\text{P': } C_1 \leftrightarrow C_2} & \left[\begin{array}{ccc} h & g & i \\ e & d & f \\ b & a & c \end{array} \right] \\
 & \xrightarrow{\text{T: } C_1 - C_2 \rightarrow C_1} & \left[\begin{array}{ccc} h-g & g & i \\ e-d & d & f \\ b-a & a & c \end{array} \right] \\
 & \xrightarrow{\text{D: } 2\rho_3 \rightarrow \rho_3} & \left[\begin{array}{ccc} h-g & g & i \\ e-d & d & f \\ 2b-2a & 2a & 2c \end{array} \right]
 \end{array}$$

Thus we take

$$\begin{aligned}
 P &= \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}, & P' &= \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \\
 T &= \begin{bmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, & D &= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix}.
 \end{aligned}$$

189 Let $E_{ij} \in M_n(\mathbb{F})$. Then

$$AE_{ij} = \begin{bmatrix} 0 & 0 & \dots & a_{1i} & \dots & 0 \\ 0 & 0 & \vdots & a_{2i} & \vdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \vdots & a_{n-1i} & \vdots & 0 \\ 0 & 0 & \vdots & a_{ni} & \vdots & 0 \end{bmatrix},$$

where the entries appear on the j -column. Then we see that $\text{tr}(AE_{ij}) = a_{ji}$ and similarly, by considering BE_{ij} , we see that $\text{tr}(BE_{ij}) = b_{ji}$. Therefore $\forall i, j, a_{ji} = b_{ji}$, which implies that $A = B$.

190 Let $E_{st} \in M_n(\mathbb{R})$. Then

$$E_{ij}A = \begin{bmatrix} 0 & 0 & \dots & 0 \\ \vdots & \vdots & \dots & \vdots \\ a_{j1} & a_{j2} & \dots & a_{jn} \\ \vdots & \vdots & \dots & \vdots \\ 0 & 0 & \dots & 0 \end{bmatrix},$$

where the entries appear on the i -th row. Thus

$$(E_{ij}A)^2 = \begin{bmatrix} 0 & 0 & \dots & 0 \\ \vdots & \vdots & \dots & \vdots \\ a_{ji}a_{j1} & a_{ji}a_{j2} & \dots & a_{ji}a_{jn} \\ \vdots & \vdots & \dots & \vdots \\ 0 & 0 & \dots & 0 \end{bmatrix},$$

which means that $\forall i, j, a_{ji}a_{jk} = 0$. In particular, $a_{ji}^2 = 0$, which means that $\forall i, j, a_{ji} = 0$, i.e., $A = 0_n$.

216 $a = 1, b = -2$.

217 Claim: $A^{-1} = I_n - A + A^2 - A^3$. For observe that

$$(I_n + A)(I_n - A + A^2 - A^3) = I_n - A + A^2 - A^3 - A + A^2 - A^3 + A^4 = I_n,$$

proving the claim.

218 Disprove! It is enough to take $A = B = 2I_n$. Then $(A + B)^{-1} = (4I_n)^{-1} = \frac{1}{4}I_n$ but $A^{-1} + B^{-1} = \frac{1}{2}I_n + \frac{1}{2}I_n = I_n$.

223 We argue by contradiction. If exactly one of the matrices is not invertible, the identities

$$\begin{aligned} A &= AI_n = (ABC)(BC)^{-1} = 0_n, \\ B &= I_n BI_n = (A)^{-1}(ABC)C^{-1} = 0_n, \\ C &= I_n C = (AB)^{-1}(ABC) = 0_n, \end{aligned}$$

shew a contradiction depending on which of the matrices are invertible. If all the matrices are invertible then

$$0_n = 0_n C^{-1} B^{-1} A^{-1} = (ABC) C^{-1} B^{-1} A^{-1} = I_n,$$

also gives a contradiction.

224 Observe that A, B, AB are invertible. Hence

$$\begin{aligned} A^2 B^2 &= I_n = (AB)^2 \implies A ABB = ABAB \\ &\implies AB = BA, \end{aligned}$$

by cancelling A on the left and B on the right. One can also argue that $A = A^{-1}$, $B = B^{-1}$, and so $AB = (AB)^{-1} = B^{-1}A^{-1} = BA$.

225 Observe that $A = (a - b)I_n + bU$, where U is the $n \times n$ matrix with 1_F 's everywhere. Prove that

$$A^2 = (2(a - b) + nb)A - ((a - b)^2 + nb(a - b))I_n.$$

226 Compute $(A - I_n)(B - I_n)$.

227 By Theorem 141 we have $\text{tr}(\mathbf{S}\mathbf{A}\mathbf{S}^{-1}) = \text{tr}(\mathbf{S}^{-1}\mathbf{S}\mathbf{A}) = \text{tr}(\mathbf{A})$.

243 The rank is 2.

244 If \mathbf{B} is invertible, then $\text{rank}(\mathbf{AB}) = \text{rank}(\mathbf{A}) = \text{rank}(\mathbf{BA})$. Similarly, if \mathbf{A} is invertible $\text{rank}(\mathbf{AB}) = \text{rank}(\mathbf{B}) = \text{rank}(\mathbf{BA})$. Now, take $\mathbf{A} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ and $\mathbf{B} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$. Then $\mathbf{AB} = \mathbf{B}$, and so $\text{rank}(\mathbf{AB}) = 1$. But $\mathbf{BA} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$, and so $\text{rank}(\mathbf{BA}) = 0$.

245 Effecting $\mathbf{R}_3 - \mathbf{R}_1 \rightarrow \mathbf{R}_3$; $a\mathbf{R}_4 - b\mathbf{R}_2 \rightarrow \mathbf{R}_4$ successively, we obtain

$$\sim \begin{bmatrix} 1 & a & 1 & b \\ a & 1 & b & 1 \\ 1 & b & 1 & a \\ b & 1 & a & 1 \end{bmatrix} \sim \begin{bmatrix} 1 & a & 1 & b \\ a & 1 & b & 1 \\ 0 & b-a & 0 & a-b \\ 0 & a-b & a^2-b^2 & a-b \end{bmatrix}.$$

Performing $\mathbf{R}_2 - a\mathbf{R}_1 \rightarrow \mathbf{R}_2$; $\mathbf{R}_4 + \mathbf{R}_3 \rightarrow \mathbf{R}_4$ we have

$$\sim \begin{bmatrix} 1 & a & 1 & b \\ 0 & 1-a^2 & b-a & 1-ab \\ 0 & b-a & 0 & a-b \\ 0 & 0 & a^2-b^2 & 2(a-b) \end{bmatrix}.$$

Performing $(1-a^2)\mathbf{R}_3 - (b-a)\mathbf{R}_2 \rightarrow \mathbf{R}_3$ we have

$$\sim \begin{bmatrix} 1 & a & 1 & b \\ 0 & 1-a^2 & b-a & 1-ab \\ 0 & 0 & -a^2+2ab-b^2 & 2a-2b-a^3+ab^2 \\ 0 & 0 & a^2-b^2 & 2(a-b) \end{bmatrix}.$$

Performing $\mathbf{R}_3 - \mathbf{R}_4 \rightarrow \mathbf{R}_3$ we have

$$\sim \begin{bmatrix} 1 & a & 1 & b \\ 0 & 1-a^2 & b-a & 1-ab \\ 0 & 0 & -2a(a-b) & -a(a^2-b^2) \\ 0 & 0 & a^2-b^2 & 2(a-b) \end{bmatrix}.$$

Performing $2a\mathbf{R}_4 + (a+b)\mathbf{R}_3 \rightarrow \mathbf{R}_4$ we have

$$\begin{bmatrix} 1 & a & 1 & b \\ 0 & 1-a^2 & b-a & 1-ab \\ 0 & 0 & -2a(a-b) & -a(a^2-b^2) \\ 0 & 0 & 0 & 4a^2-4ab-a^4+a^2b^2-ba^3+ab^3 \end{bmatrix}.$$

Factorising, this is

$$= \begin{bmatrix} 1 & a & 1 & b \\ 0 & 1-a^2 & b-a & 1-ab \\ 0 & 0 & -2a(a-b) & -a(a-b)(a+b) \\ 0 & 0 & 0 & -a(a+2+b)(a-b)(a-2+b) \end{bmatrix}.$$

Thus the rank is 4 if $(a+2+b)(a-b)(a-2+b) \neq 0$. The rank is 3 if $a+b=2$ and $(a,b) \neq (1,1)$ or if $a+b=-2$ and $(a,b) \neq (-1,-1)$. The rank is 2 if $a=b \neq 1$ and $a \neq -1$. The rank is 1 if $a=b=\pm 1$.

246 $\text{rank } (\mathbf{A}) = 4$ if $m^3 + m^2 + 2 \neq 0$, and $\text{rank } (\mathbf{A}) = 3$ otherwise.

247 The rank is 4 if $a \neq \pm b$. The rank is 1 is $a = \pm b \neq 0$. The rank is 0 if $a = b = 0$.

248 The rank is 4 if $(a-b)(c-d) \neq 0$. The rank is 2 is $a = b, c \neq d$ or if $a \neq b, c = d$. The rank is 1 if $a = b$ and $c = d$.

249 $\text{rank } (\mathbf{ABC}) \leq 2 \implies x = 13$.

252 For the counterexample consider $\mathbf{A} = \begin{bmatrix} 1 & i \\ i & -1 \end{bmatrix}$.

261 We form the augmented matrix

$$\left[\begin{array}{ccc|ccc} \bar{1} & \bar{2} & \bar{3} & \bar{1} & \bar{0} & \bar{0} \\ \bar{2} & \bar{3} & \bar{1} & \bar{0} & \bar{1} & \bar{0} \\ \bar{3} & \bar{1} & \bar{2} & \bar{0} & \bar{0} & \bar{1} \end{array} \right]$$

From $R_2 - \bar{2}R_1 \rightarrow R_2$ and $R_3 - \bar{3}R_1 \rightarrow R_3$ we obtain

$$\sim \left[\begin{array}{ccc|ccc} \bar{1} & \bar{2} & \bar{3} & \bar{1} & \bar{0} & \bar{0} \\ \bar{0} & \bar{6} & \bar{2} & \bar{5} & \bar{1} & \bar{0} \\ \bar{0} & \bar{2} & \bar{0} & \bar{4} & \bar{0} & \bar{1} \end{array} \right].$$

From $R_2 \leftrightarrow R_3$ we obtain

$$\sim \left[\begin{array}{ccc|ccc} \bar{1} & \bar{2} & \bar{3} & \bar{1} & \bar{0} & \bar{0} \\ \bar{0} & \bar{2} & \bar{0} & \bar{4} & \bar{0} & \bar{1} \\ \bar{0} & \bar{6} & \bar{2} & \bar{5} & \bar{1} & \bar{0} \end{array} \right].$$

Now, from $R_1 - R_2 \rightarrow R_1$ and $R_3 - \bar{3}R_2 \rightarrow R_3$, we obtain

$$\sim \left[\begin{array}{ccc|ccc} \bar{1} & \bar{0} & \bar{3} & \bar{4} & \bar{0} & \bar{6} \\ \bar{0} & \bar{2} & \bar{0} & \bar{4} & \bar{0} & \bar{1} \\ \bar{0} & \bar{0} & \bar{2} & \bar{0} & \bar{1} & \bar{4} \end{array} \right].$$

From $4R_2 \rightarrow R_2$ and $4R_3 \rightarrow R_3$, we obtain

$$\sim \left[\begin{array}{ccc|ccc} \bar{1} & \bar{0} & \bar{3} & \bar{4} & \bar{0} & \bar{6} \\ \bar{0} & \bar{1} & \bar{0} & \bar{2} & \bar{0} & \bar{4} \\ \bar{0} & \bar{0} & \bar{1} & \bar{0} & \bar{4} & \bar{2} \end{array} \right].$$

Finally, from $\mathbf{R}_1 - \bar{3}\mathbf{R}_3 \rightarrow \mathbf{R}_3$ we obtain

$$\rightsquigarrow \left[\begin{array}{ccc|ccc} \bar{1} & \bar{0} & \bar{0} & \bar{4} & \bar{2} & \bar{0} \\ \bar{0} & \bar{1} & \bar{0} & \bar{2} & \bar{0} & \bar{4} \\ \bar{0} & \bar{0} & \bar{1} & \bar{0} & \bar{4} & \bar{2} \end{array} \right].$$

We deduce that

$$\left[\begin{array}{ccc} \bar{1} & \bar{2} & \bar{3} \\ \bar{2} & \bar{3} & \bar{1} \\ \bar{3} & \bar{1} & \bar{2} \end{array} \right]^{-1} = \left[\begin{array}{ccc} \bar{4} & \bar{2} & \bar{0} \\ \bar{2} & \bar{0} & \bar{4} \\ \bar{0} & \bar{4} & \bar{2} \end{array} \right].$$

262 To find the inverse of \mathbf{B} we consider the augmented matrix

$$\left[\begin{array}{ccc|ccc} 0 & 0 & -1 & 1 & 0 & 0 \\ 0 & -1 & a & 0 & 1 & 0 \\ -1 & a & b & 0 & 0 & 1 \end{array} \right].$$

Performing $\mathbf{R}_1 \leftrightarrow \mathbf{R}_3, -\mathbf{R}_3 \rightarrow \mathbf{R}_3$, in succession,

$$\left[\begin{array}{ccc|ccc} -1 & a & b & 0 & 0 & 1 \\ 0 & -1 & a & 0 & 1 & 0 \\ 0 & 0 & 1 & -1 & 0 & 0 \end{array} \right].$$

Performing $\mathbf{R}_1 + a\mathbf{R}_2 \rightarrow \mathbf{R}_1$ and $\mathbf{R}_2 - a\mathbf{R}_3 \rightarrow \mathbf{R}_2$ in succession,

$$\left[\begin{array}{ccc|ccc} -1 & 0 & b + a^2 & 0 & a & 1 \\ 0 & -1 & 0 & a & 1 & 0 \\ 0 & 0 & 1 & -1 & 0 & 0 \end{array} \right].$$

Performing $\mathbf{R}_1 - (b + a^2)\mathbf{R}_3 \rightarrow \mathbf{R}_3, -\mathbf{R}_1 \rightarrow \mathbf{R}_1$ and $-\mathbf{R}_2 \rightarrow \mathbf{R}_2$ in succession, we find

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -b - a^2 & -a & -1 \\ 0 & 1 & 0 & -a & -1 & 0 \\ 0 & 0 & 1 & -1 & 0 & 0 \end{array} \right],$$

whence

$$\mathbf{B}^{-1} = \left[\begin{array}{ccc} -b - a^2 & -a & -1 \\ -a & -1 & 0 \\ -1 & 0 & 0 \end{array} \right].$$

Now,

$$\begin{aligned}
 \mathbf{BAB}^{-1} &= \begin{bmatrix} 0 & 0 & -1 \\ 0 & -1 & a \\ -1 & a & b \end{bmatrix} \begin{bmatrix} a & b & c \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} -b - a^2 & -a & -1 \\ -a & -1 & 0 \\ -1 & 0 & 0 \end{bmatrix} \\
 &= \begin{bmatrix} 0 & -1 & 0 \\ -1 & a & 0 \\ 0 & 0 & -c \end{bmatrix} \begin{bmatrix} -b - a^2 & -a & -1 \\ -a & -1 & 0 \\ -1 & 0 & 0 \end{bmatrix} \\
 &= \begin{bmatrix} a & 1 & 0 \\ b & 0 & 1 \\ c & 0 & 0 \end{bmatrix} \\
 &= \mathbf{A}^T,
 \end{aligned}$$

which is what we wanted to prove.

263 Operating formally, and using elementary row operations, we find

$$\mathbf{B}^{-1} = \begin{bmatrix} -\frac{a^2-1}{a^2-5+2a} & \frac{a^2+2a-2}{a^2-5+2a} & \frac{a-2}{a^2-5+2a} \\ -\frac{2}{a^2-5+2a} & \frac{a+4}{a^2-5+2a} & -\frac{1}{a^2-5+2a} \\ \frac{2a}{a^2-5+2a} & -\frac{2a+5}{a^2-5+2a} & \frac{a}{a^2-5+2a} \end{bmatrix}.$$

Thus \mathbf{B} is invertible whenever $a \neq -1 \pm \sqrt{6}$.

264 Form the augmented matrix

$$\left[\begin{array}{ccc|ccc} a & 2a & 3a & 1 & 0 & 0 \\ 0 & b & 2b & 0 & 1 & 0 \\ 0 & 0 & c & 0 & 0 & 1 \end{array} \right].$$

Perform $\frac{1}{a}\mathbf{R}_1 \rightarrow \mathbf{R}_1$, $\frac{1}{b}\mathbf{R}_2 \rightarrow \mathbf{R}_2$, $\frac{1}{a}\mathbf{R}_3 \rightarrow \mathbf{R}_3$, in succession, obtaining

$$\left[\begin{array}{ccc|ccc} 1 & 2 & 3 & 1/a & 0 & 0 \\ 0 & 1 & 2 & 0 & 1/b & 0 \\ 0 & 0 & 1 & 0 & 0 & 1/c \end{array} \right].$$

Now perform $\mathbf{R}_1 - 2\mathbf{R}_2 \rightarrow \mathbf{R}_1$ and $\mathbf{R}_2 - 2\mathbf{R}_3 \rightarrow \mathbf{R}_2$ in succession, to obtain

$$\left[\begin{array}{ccc|ccc} 1 & 0 & -1 & 1/a & -2/a & 0 \\ 0 & 1 & 0 & 0 & 1/b & -2/c \\ 0 & 0 & 1 & 0 & 0 & 1/c \end{array} \right].$$

Finally, perform $R_1 + R_3 \rightarrow R_1$ to obtain

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 1/a & -2/b & 1/c \\ 0 & 1 & 0 & 0 & 1/b & -2/c \\ 0 & 0 & 1 & 0 & 0 & 1/c \end{array} \right].$$

Whence

$$\left[\begin{array}{ccc} a & 2a & 3a \\ 0 & b & 2b \\ 0 & 0 & c \end{array} \right]^{-1} = \left[\begin{array}{ccc} 1/a & -2/b & 1/c \\ 0 & 1/b & -2/c \\ 0 & 0 & 1/c \end{array} \right].$$

265 To compute the inverse matrix we proceed formally as follows. The augmented matrix is

$$\left[\begin{array}{ccc|ccc} b & a & 0 & 1 & 0 & 0 \\ c & 0 & a & 0 & 1 & 0 \\ 0 & c & b & 0 & 0 & 1 \end{array} \right].$$

Performing $bR_2 - cR_1 \rightarrow R_2$ we find

$$\left[\begin{array}{ccc|ccc} b & a & 0 & 1 & 0 & 0 \\ 0 & -ca & ab & -c & b & 0 \\ 0 & c & b & 0 & 0 & 1 \end{array} \right].$$

Performing $aR_3 + R_2 \rightarrow R_3$ we obtain

$$\left[\begin{array}{ccc|ccc} b & a & 0 & 1 & 0 & 0 \\ 0 & -ca & ab & -c & b & 0 \\ 0 & 0 & 2ab & -c & b & a \end{array} \right].$$

Performing $2R_2 - R_3 \rightarrow R_2$ we obtain

$$\left[\begin{array}{ccc|ccc} b & a & 0 & 1 & 0 & 0 \\ 0 & -2ca & 0 & -c & b & -a \\ 0 & 0 & 2ab & -c & b & a \end{array} \right].$$

Performing $2cR_1 + R_2 \rightarrow R_1$ we obtain

$$\left[\begin{array}{ccc|ccc} 2bc & 0 & 0 & c & b & -a \\ 0 & -2ca & 0 & -c & b & -a \\ 0 & 0 & 2ab & -c & b & a \end{array} \right].$$

From here we easily conclude that

$$\left[\begin{array}{ccc} b & a & 0 \\ c & 0 & a \\ 0 & c & b \end{array} \right]^{-1} = \left[\begin{array}{ccc} \frac{1}{2b} & \frac{1}{2c} & -\frac{a}{2bc} \\ \frac{1}{2a} & -\frac{b}{2ac} & \frac{1}{2c} \\ -\frac{c}{2ba} & \frac{1}{2a} & \frac{1}{2b} \end{array} \right]$$

as long as $abc \neq 0$.

266 Form the expanded matrix

$$\left[\begin{array}{ccc|ccc} 1+a & 1 & 1 & 1 & 0 & 0 \\ 1 & 1+b & 1 & 0 & 1 & 0 \\ 1 & 1 & 1+c & 0 & 0 & 1 \end{array} \right].$$

Perform $bcR_1 \rightarrow R_1$, $abR_3 \rightarrow R_3$, $caR_2 \rightarrow R_2$. The matrix turns into

$$\left[\begin{array}{ccc|ccc} bc+abc & bc & bc & bc & 0 & 0 \\ ca & ca+abc & ca & 0 & ca & 0 \\ ab & ab & ab+abc & 0 & 0 & ab \end{array} \right].$$

Perform $R_1 + R_2 + R_3 \rightarrow R_1$ the matrix turns into

$$\left[\begin{array}{ccc|ccc} ab+bc+ca+abc & ab+bc+ca+abc & ab+bc+ca+abc & bc & ca & ab \\ ca & ca+abc & ca & 0 & ca & 0 \\ ab & ab & ab+abc & 0 & 0 & ab \end{array} \right].$$

Perform $\frac{1}{ab+bc+ca+abc}R_1 \rightarrow R_1$. The matrix turns into

$$\left[\begin{array}{ccc|ccc} 1 & 1 & 1 & \frac{bc}{ab+bc+ca+abc} & \frac{ca}{ab+bc+ca+abc} & \frac{ab}{ab+bc+ca+abc} \\ ca & ca+abc & ca & 0 & ca & 0 \\ ab & ab & ab+abc & 0 & 0 & ab \end{array} \right].$$

Perform $R_2 - caR_1 \rightarrow R_2$ and $R_3 - abR_3 \rightarrow R_3$. We get

$$\left[\begin{array}{ccc|ccc} 1 & 1 & 1 & \frac{bc}{ab+bc+ca+abc} & \frac{ca}{ab+bc+ca+abc} & \frac{ab}{ab+bc+ca+abc} \\ 0 & abc & 0 & -\frac{abc^2}{ab+bc+ca+abc} & ca - \frac{c^2a^2}{ab+bc+ca+abc} & -\frac{a^2bc}{ab+bc+ca+abc} \\ 0 & 0 & abc & -\frac{ab^2c}{ab+bc+ca+abc} & -\frac{a^2bc}{ab+bc+ca+abc} & ab - \frac{a^2b^2}{ab+bc+ca+abc} \end{array} \right].$$

Perform $\frac{1}{ABC}R_2 \rightarrow R_2$ and $\frac{1}{ABC}R_3 \rightarrow R_3$. We obtain

$$\left[\begin{array}{ccc|ccc} 1 & 1 & 1 & \frac{bc}{ab+bc+ca+abc} & \frac{ca}{ab+bc+ca+abc} & \frac{ab}{ab+bc+ca+abc} \\ 0 & 1 & 0 & -\frac{c}{ab+bc+ca+abc} & \frac{1}{b} - \frac{ca}{b(ab+bc+ca+abc)} & -\frac{a}{ab+bc+ca+abc} \\ 0 & 0 & 1 & -\frac{b}{ab+bc+ca+abc} & -\frac{a}{ab+bc+ca+abc} & \frac{1}{c} - \frac{ab}{c(ab+bc+ca+abc)} \end{array} \right].$$

Finally we perform $R_1 - R_2 - R_3 \rightarrow R_1$, getting

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{a+b+c}{ab+bc+ca+abc} & -\frac{c}{ab+bc+ca+abc} & -\frac{b}{ab+bc+ca+abc} \\ 0 & 1 & 0 & -\frac{c}{ab+bc+ca+abc} & \frac{1}{b} - \frac{ca}{b(ab+bc+ca+abc)} & -\frac{a}{ab+bc+ca+abc} \\ 0 & 0 & 1 & -\frac{b}{ab+bc+ca+abc} & -\frac{a}{ab+bc+ca+abc} & \frac{1}{c} - \frac{ab}{c(ab+bc+ca+abc)} \end{array} \right].$$

We conclude that the inverse is

$$\begin{bmatrix} \frac{b+c+bc}{ab+bc+ca+abc} & -\frac{c}{ab+bc+ca+abc} & -\frac{b}{ab+bc+ca+abc} \\ -\frac{c}{ab+bc+ca+abc} & \frac{c+a+ca}{ab+bc+ca+abc} & -\frac{a}{ab+bc+ca+abc} \\ -\frac{b}{ab+bc+ca+abc} & -\frac{a}{ab+bc+ca+abc} & \frac{a+b+ab}{ab+bc+ca+abc} \end{bmatrix}$$

271 Since $\text{rank } (\mathbf{A}^2) < 5$, \mathbf{A}^2 is not invertible. But then \mathbf{A} is not invertible and hence $\text{rank } (\mathbf{A}) < 5$.

282 The free variables are z and w . We have

$$\bar{2}\mathbf{y} + \mathbf{w} = \bar{2} \implies \bar{2}\mathbf{y} = \bar{2} - \mathbf{w} \implies \mathbf{y} = \bar{1} + \mathbf{w},$$

and

$$x + \mathbf{y} + z + \mathbf{w} = \bar{0} \implies x = -\mathbf{y} - z - \mathbf{w} = \bar{2}\mathbf{y} + \bar{2}z + \bar{2}\mathbf{w}.$$

Hence

$$\begin{bmatrix} x \\ \mathbf{y} \\ z \\ \mathbf{w} \end{bmatrix} = \begin{bmatrix} \bar{0} \\ \bar{1} \\ \bar{0} \\ \bar{0} \end{bmatrix} + z \begin{bmatrix} \bar{0} \\ \bar{0} \\ \bar{1} \\ \bar{0} \end{bmatrix} + \mathbf{w} \begin{bmatrix} \bar{0} \\ \bar{0} \\ \bar{0} \\ \bar{1} \end{bmatrix}.$$

This gives the 9 solutions.

283 We have

$$\begin{bmatrix} \bar{1} & \bar{2} & \bar{3} \\ \bar{2} & \bar{3} & \bar{1} \\ \bar{3} & \bar{1} & \bar{2} \end{bmatrix} \begin{bmatrix} x \\ \mathbf{y} \\ z \end{bmatrix} = \begin{bmatrix} \bar{5} \\ \bar{6} \\ \bar{0} \end{bmatrix},$$

Hence

$$\begin{bmatrix} x \\ \mathbf{y} \\ z \end{bmatrix} = \begin{bmatrix} \bar{1} & \bar{2} & \bar{3} \\ \bar{2} & \bar{3} & \bar{1} \\ \bar{3} & \bar{1} & \bar{2} \end{bmatrix}^{-1} \begin{bmatrix} \bar{5} \\ \bar{6} \\ \bar{0} \end{bmatrix} = \begin{bmatrix} \bar{4} & \bar{2} & \bar{0} \\ \bar{2} & \bar{0} & \bar{4} \\ \bar{0} & \bar{4} & \bar{2} \end{bmatrix} \begin{bmatrix} \bar{5} \\ \bar{6} \\ \bar{0} \end{bmatrix} = \begin{bmatrix} \bar{4} \\ \bar{3} \\ \bar{3} \end{bmatrix}.$$

291 Observe that the third row is the sum of the first two rows and the fourth row is twice the third. So we have

$$\begin{array}{c|c}
 \left[\begin{array}{ccccc|c} 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 0 & 1 & 0 & 1 & -1 \\ 2 & 1 & 2 & 1 & 2 & 0 \\ 4 & 2 & 4 & 2 & 4 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 \end{array} \right] & \left[\begin{array}{ccccc|c} 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 0 & 1 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 \end{array} \right] \\
 \xrightarrow{\substack{R_3 - R_1 - R_2 \rightarrow R_3 \\ R_4 - 2R_1 - 2R_2 \rightarrow R_4}} & \\
 \left[\begin{array}{ccccc|c} 1 & 1 & 1 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 & 0 & -1 \\ 0 & 0 & 1 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 \end{array} \right] & \left[\begin{array}{ccccc|c} 0 & 1 & 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 \end{array} \right] \\
 \xrightarrow{\substack{R_2 - R_5 \rightarrow R_2 \\ R_1 - R_5 \rightarrow R_1}} & \\
 \left[\begin{array}{ccccc|c} 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right] &
 \end{array}$$

Rearranging the rows we obtain

$$\left[\begin{array}{ccccc|c} 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right].$$

Hence d and f are free variables. We obtain

$$\begin{aligned} c &= -1, \\ b &= 1 - c - d = 2 - d, \\ a &= -f. \end{aligned}$$

The solution is

$$\begin{bmatrix} a \\ b \\ c \\ d \\ f \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \\ -1 \\ 0 \\ 0 \end{bmatrix} + d \begin{bmatrix} 0 \\ -1 \\ 0 \\ 1 \\ 0 \end{bmatrix} + f \begin{bmatrix} -1 \\ 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}.$$

292 The unique solution is

$$\begin{bmatrix} 1 \\ 1 \\ -1 \\ 1 \\ 1 \end{bmatrix}.$$

293 The augmented matrix of the system is

$$\left[\begin{array}{ccc|c} 2m & 1 & 1 & 2 \\ 1 & 2m & 1 & 4m \\ 1 & 1 & 2m & 2m^2 \end{array} \right].$$

Performing $R_1 \leftrightarrow R_2$.

$$\left[\begin{array}{ccc|c} 1 & 2m & 1 & 4m \\ 2m & 1 & 1 & 2 \\ 1 & 1 & 2m & 2m^2 \end{array} \right].$$

Performing $R_2 \leftrightarrow R_3$.

$$\left[\begin{array}{ccc|c} 1 & 2m & 1 & 4m \\ 1 & 1 & 2m & 2m^2 \\ 2m & 1 & 1 & 2 \end{array} \right].$$

Performing $R_2 - R_1 \rightarrow R_1$ and $R_3 - 2mR_1 \rightarrow R_3$ we obtain

$$\left[\begin{array}{ccc|c} 1 & 2m & 1 & 4m \\ 0 & 1-2m & 2m-1 & 2m^2-4m \\ 0 & 1-4m^2 & 1-2m & 2-8m^2 \end{array} \right].$$

If $m = \frac{1}{2}$ the matrix becomes

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 2 \\ 0 & 0 & 0 & -\frac{3}{2} \\ 0 & 0 & 0 & 0 \end{array} \right]$$

and hence it does not have a solution. If $m \neq \frac{1}{2}$, by performing $\frac{1}{1-2m}R_2 \rightarrow R_2$ and $\frac{1}{1-2m}R_3 \rightarrow R_3$, the matrix becomes

$$\left[\begin{array}{ccc|c} 1 & 2m & 1 & 4m \\ 0 & 1 & -1 & \frac{2m(m-2)}{1-2m} \\ 0 & 1+2m & 1 & 2(1+2m) \end{array} \right].$$

Performing $R_3 - (1+2m)R_2 \rightarrow R_3$ we obtain

$$\left[\begin{array}{ccc|c} 1 & 2m & 1 & 4m \\ 0 & 1 & -1 & \frac{2m(m-2)}{1-2m} \\ 0 & 0 & 2+2m & \frac{2(1+2m)(1-m^2)}{1-2m} \end{array} \right].$$

If $m = -1$ then the matrix reduces to

$$\left[\begin{array}{ccc|c} 1 & -2 & 1 & -4 \\ 0 & 1 & -1 & 2 \\ 0 & 0 & 0 & 0 \end{array} \right].$$

The solution in this case is

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} z \\ 2+z \\ z \end{bmatrix}.$$

If $m \neq -1, m \neq -\frac{1}{2}$ we have the solutions

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} \frac{m-1}{1-2m} \\ \frac{1-3m}{1-2m} \\ \frac{(1+2m)(1-m)}{1-2m} \end{bmatrix}.$$

294 By performing the elementary row operations, we obtain the following triangular form:

$$\begin{aligned} ax + y - 2z &= 1, \\ (a-1)^2y + (1-a)(a-2)z &= 1-a, \\ (a-2)z &= 0. \end{aligned}$$

If $a = 2$, there is an infinity of solutions:

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1+t \\ -1 \\ t \end{bmatrix} \quad t \in \mathbb{R}.$$

Assume $a \neq 2$. Then $z = 0$ and the system becomes

$$\begin{aligned} ax + y &= 1, \\ (a-1)^2y &= 1-a, \\ 2x + (3-a)y &= 1. \end{aligned}$$

We see that if $a = 1$, the system becomes

$$\begin{aligned} x + y &= 1, \\ 2x + 2y &= 1, \end{aligned}$$

and so there is no solution. If $(a-1)(a-2) \neq 0$, we obtain the unique solution

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} \frac{1}{a-1} \\ -\frac{1}{a-1} \\ 0 \end{bmatrix}.$$

295 The system is solvable if $m \neq 0, m \neq \pm 2$. If $m \neq 2$ there is the solution

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} \frac{1}{m-2} \\ \frac{m+3}{m-2} \\ \frac{m+2}{m-2} \end{bmatrix}.$$

296 There is the unique solution

$$\begin{bmatrix} x \\ y \\ z \\ t \end{bmatrix} = \begin{bmatrix} a+d+b-c \\ -c-d-b+a \\ d+c-b+a \\ c-d+b+a \end{bmatrix}.$$

297 The system can be written as

$$\begin{bmatrix} b & a & 0 \\ c & 0 & a \\ 0 & c & b \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} c \\ b \\ a \end{bmatrix}.$$

The system will have the unique solution

$$\begin{aligned} \begin{bmatrix} x \\ y \\ z \end{bmatrix} &= \begin{bmatrix} b & a & 0 \\ c & 0 & a \\ 0 & c & b \end{bmatrix}^{-1} \begin{bmatrix} c \\ b \\ a \end{bmatrix} \\ &= \begin{bmatrix} \frac{1}{2b} & \frac{1}{2c} & -\frac{a}{2bc} \\ \frac{1}{2a} & -\frac{b}{2ac} & \frac{1}{2c} \\ -\frac{c}{2ba} & \frac{1}{2a} & \frac{1}{2b} \end{bmatrix} \begin{bmatrix} c \\ b \\ a \end{bmatrix}, \\ &= \begin{bmatrix} \frac{b^2 + c^2 - a^2}{2bc} \\ \frac{a^2 + c^2 - b^2}{2ac} \\ \frac{a^2 + b^2 - c^2}{2ab} \end{bmatrix} \end{aligned}$$

as long as the inverse matrix exists, which is as long as $abc \neq 0$

298

$$x = 2^{-2}3^6$$

$$y = 2^{-3}3^{12}$$

$$z = 2^23^{-7}.$$

299 Denote the addition operations applied to the rows by a_1, a_2, a_3, a_4 and the subtraction operations to the columns by b_1, b_2, b_3, b_4 . Comparing A and A^T we obtain 7 equations in 8 unknowns. By inspecting the diagonal entries, and the entries of the first row of A and A^T , we deduce the following equations

$$a_1 = b_1,$$

$$a_2 = b_2,$$

$$a_3 = b_3,$$

$$a_4 = b_4,$$

$$a_1 - b_2 = 3,$$

$$a_1 - b_3 = 6,$$

$$a_1 - b_4 = 9.$$

This is a system of 7 equations in 8 unknowns. We may let $a_4 = 0$ and thus obtain $a_1 = b_1 = 9$, $a_2 = b_2 = 6$, $a_3 = b_3 = 3$, $a_4 = b_4 = 0$.

300 The augmented matrix of this system is

$$\left[\begin{array}{ccccc|c} -y & 1 & 0 & 0 & 1 & 0 \\ 1 & -y & 1 & 0 & 0 & 0 \\ 0 & 1 & -y & 1 & 0 & 0 \\ 0 & 0 & 1 & -y & 1 & 0 \\ 1 & 0 & 0 & 1 & -y & 0 \end{array} \right].$$

Permute the rows to obtain

$$\left[\begin{array}{ccccc|c} 1 & 0 & 0 & 1 & -y & 0 \\ 0 & 1 & -y & 1 & 0 & 0 \\ 0 & 0 & 1 & -y & 1 & 0 \\ 1 & -y & 1 & 0 & 0 & 0 \\ -y & 1 & 0 & 0 & 1 & 0 \end{array} \right].$$

Performing $R_5 + yR_1 \rightarrow R_5$ and $R_4 - R_1 \rightarrow R_4$ we get

$$\left[\begin{array}{ccccc|c} 1 & 0 & 0 & 1 & -y & 0 \\ 0 & 1 & -y & 1 & 0 & 0 \\ 0 & 0 & 1 & -y & 1 & 0 \\ 0 & -y & 1 & -1 & y & 0 \\ 0 & 1 & 0 & y & 1-y^2 & 0 \end{array} \right].$$

Performing $R_5 - R_2 \rightarrow R_5$ and $R_4 + yR_2 \rightarrow R_4$ we get

$$\left[\begin{array}{ccccc|c} 1 & 0 & 0 & 1 & -y & 0 \\ 0 & 1 & -y & 1 & 0 & 0 \\ 0 & 0 & 1 & -y & 1 & 0 \\ 0 & 0 & 1-y^2 & y-1 & y & 0 \\ 0 & 0 & y & y-1 & 1-y^2 & 0 \end{array} \right].$$

Performing $R_5 - yR_3 \rightarrow R_5$ and $R_4 + (y^2 - 1)R_3 \rightarrow R_4$ we get

$$\left[\begin{array}{ccccc|c} 1 & 0 & 0 & 1 & -y & 0 \\ 0 & 1 & -y & 1 & 0 & 0 \\ 0 & 0 & 1 & -y & 1 & 0 \\ 0 & 0 & 0 & -y^3 + 2y - 1 & y^2 + y - 1 & 0 \\ 0 & 0 & 0 & y^2 + y - 1 & 1 - y - y^2 & 0 \end{array} \right].$$

Performing $R_5 + R_4 \rightarrow R_5$ we get

$$\left[\begin{array}{ccccc} 1 & 0 & 0 & 1 & -y \\ 0 & 1 & -y & 1 & 0 \\ 0 & 0 & 1 & -y & 1 \\ 0 & 0 & 0 & -y^3 + 2y - 1 & y^2 + y - 1 \\ 0 & 0 & 0 & -y^3 + y^2 + 3y - 2 & 0 \end{array} \right] .$$

Upon factoring, the matrix is equivalent to

$$\left[\begin{array}{ccccc} 1 & 0 & 0 & 1 & -y \\ 0 & 1 & -y & 1 & 0 \\ 0 & 0 & 1 & -y & 1 \\ 0 & 0 & 0 & -(y-1)(y^2+y-1) & y^2+y-1 \\ 0 & 0 & 0 & -(y-2)(y^2+y-1) & 0 \end{array} \right] .$$

Thus $(y-2)(y^2+y-1)x_4 = 0$. If $y = 2$ then the system reduces to

$$\left[\begin{array}{ccccc} 1 & 0 & 0 & 1 & -2 \\ 0 & 1 & -2 & 1 & 0 \\ 0 & 0 & 1 & -2 & 1 \\ 0 & 0 & 0 & -5 & 5 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right] .$$

In this case x_5 is free and by backwards substitution we obtain

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = \begin{bmatrix} t \\ t \\ t \\ t \\ t \end{bmatrix}, \quad t \in \mathbb{R}.$$

If $y^2 + y - 1 = 0$ then the system reduces to

$$\left[\begin{array}{ccccc} 1 & 0 & 0 & 1 & -y \\ 0 & 1 & -y & 1 & 0 \\ 0 & 0 & 1 & -y & 1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right] .$$

In this case x_4, x_5 are free, and

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = \begin{bmatrix} yt - s \\ y^2s - yt - s \\ ys - t \\ s \\ t \end{bmatrix}, \quad (s, t) \in \mathbb{R}^2.$$

Since $y^2s - s = (y^2 + y - 1)s - ys$, this last solution can be also written as

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = \begin{bmatrix} yt - s \\ -ys - yt \\ ys - t \\ s \\ t \end{bmatrix}, \quad (s, t) \in \mathbb{R}^2.$$

Finally, if $(y - 2)(y^2 + y - 1) \neq 0$, then $x_4 = 0$, and we obtain

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}.$$

323 $\sqrt{2a^2 - 2a + 1}$

324 $\|\lambda v\| = \frac{1}{2} \implies \sqrt{(\lambda)^2 + (-\lambda)^2} = \frac{1}{2} \implies 2\lambda^2 = \frac{1}{4} \implies \lambda = \pm \frac{1}{\sqrt{8}}$.

325 0

326 $a = \pm 1$ or $a = -8$.

327 [A] $2(x + y) - \frac{1}{2}z$, [B] $x + y - \frac{1}{2}z$, [C] $-(x + y + z)$

328 [A]. 0, [B]. 0, [C]. 0, [D]. 0, [E]. $2c (= 2d)$

329 [F]. 0, [G]. b, [H]. 20, [I]. 0.

330 Let the skew quadrilateral be ABCD and let P, Q, R, S be the midpoints of [A, B], [B, C], [C, D], [D, A], respectively. Put $x = OX$, where $X \in \{A, B, C, D, P, Q, R, S\}$. Using the Section Formula 4.4 we have

$$p = \frac{a+b}{2}, \quad q = \frac{b+c}{2}, \quad r = \frac{c+d}{2}, \quad s = \frac{d+a}{2}.$$

This gives

$$p - q = \frac{a - c}{2}, \quad s - r = \frac{a - c}{2}.$$

This means that $\overrightarrow{QP} = \overrightarrow{RS}$ and so PQRS is a parallelogram since one pair of sides are equal and parallel.

331 We have $2\vec{BC} = \vec{BE} + \vec{EC}$. By Chasles' Rule $\vec{AC} = \vec{AE} + \vec{EC}$, and $\vec{BD} = \vec{BE} + \vec{ED}$. We deduce that

$$\vec{AC} + \vec{BD} = \vec{AE} + \vec{EC} + \vec{BE} + \vec{ED} = \vec{AD} + \vec{BC}.$$

But since $ABCD$ is a parallelogram, $\vec{AD} = \vec{BC}$. Hence

$$\vec{AC} + \vec{BD} = \vec{AD} + \vec{BC} = 2\vec{BC}.$$

332 We have $\vec{IA} = -3\vec{IB} \iff \vec{IA} = -3(\vec{IA} + \vec{AB}) = -3\vec{IA} - 3\vec{AB}$. Thus we deduce

$$\begin{aligned}\vec{IA} + 3\vec{IA} &= -3\vec{AB} \iff 4\vec{IA} = -3\vec{AB} \\ &\iff 4\vec{AI} = 3\vec{AB} \\ &\iff \vec{AI} = \frac{3}{4}\vec{AB}.\end{aligned}$$

Similarly

$$\begin{aligned}\vec{JA} &= -\frac{1}{3}\vec{JB} \iff 3\vec{JA} = -\vec{JB} \\ &\iff 3\vec{JA} = -\vec{JA} - \vec{AB} \\ &\iff 4\vec{JA} = -\vec{AB} \\ &\iff \vec{AJ} = \frac{1}{4}\vec{AB}\end{aligned}$$

Thus we take I such that $\vec{AI} = \frac{3}{4}\vec{AB}$ and J such that $\vec{AJ} = \frac{1}{4}\vec{AB}$.

Now

$$\begin{aligned}\vec{MA} + 3\vec{MB} &= \vec{MI} + \vec{IA} + 3\vec{IB} \\ &= 4\vec{MI} + \vec{IA} + 3\vec{IB} \\ &= 4\vec{MI},\end{aligned}$$

and

$$\begin{aligned}3\vec{MA} + \vec{MB} &= 3\vec{MJ} + 3\vec{JA} + \vec{MJ} + \vec{JB} \\ &= 4\vec{MJ} + 3\vec{JA} + \vec{JB} \\ &= 4\vec{MJ}.\end{aligned}$$

333 Let G , O and P denote vectors from an arbitrary origin to the gallows, oak, and pine, respectively. The conditions of the problem define X and Y , thought of similarly as vectors from the origin, by $X = O + R(O - G)$, $Y = P + R(P - G)$, where R is the 90° rotation to the right, a linear transformation on vectors in the plane; the fact that $-R$ is 90° leftward rotation has been used in writing Y . Anyway, then

$$\frac{\vec{X} + \vec{Y}}{2} = \frac{\vec{O} + \vec{P}}{2} + \frac{R(\vec{O} - \vec{G}) + R(\vec{P} - \vec{G})}{2}$$

is independent of the position of the gallows. This gives a simple algorithm for treasure-finding: take P as the (hitherto) arbitrary origin, then the treasure is at $\frac{\vec{O} + R(\vec{O})}{2}$.

352 $a = \frac{1}{2}$

354

$$\vec{p} = \begin{bmatrix} 4 \\ 5 \end{bmatrix} = 2 \begin{bmatrix} -1 \\ 1 \end{bmatrix} + 3 \begin{bmatrix} 2 \\ 1 \end{bmatrix} = 2\vec{r} + 3\vec{s}.$$

355 Since $\mathbf{a}_1 = \mathbf{a} \bullet \mathbf{i}$, $\mathbf{a}_2 = \mathbf{a} \bullet \mathbf{j}$, we may write

$$\mathbf{a} = (\mathbf{a} \bullet \mathbf{i})\mathbf{i} + (\mathbf{a} \bullet \mathbf{j})\mathbf{j}$$

from where the assertion follows.

356

$$\begin{aligned}\alpha \mathbf{a} + \beta \mathbf{b} = \mathbf{0} &\implies \mathbf{a} \bullet (\alpha \mathbf{a} + \beta \mathbf{b}) = \mathbf{a} \bullet \mathbf{0} \\ &\implies \alpha(\mathbf{a} \bullet \mathbf{a}) = 0 \\ &\implies \alpha \|\mathbf{a}\|^2 = 0.\end{aligned}$$

Since $\mathbf{a} \neq \mathbf{0}$, we must have $\|\mathbf{a}\| \neq 0$ and thus $\alpha = 0$. But if $\alpha = 0$ then

$$\begin{aligned}\alpha \mathbf{a} + \beta \mathbf{b} = \mathbf{0} &\implies \beta \mathbf{b} = \mathbf{0} \\ &\implies \beta = 0,\end{aligned}$$

since $\mathbf{b} \neq \mathbf{0}$.

357 We must shew that

$$(2\mathbf{x} + 3\mathbf{y}) \bullet (2\mathbf{x} - 3\mathbf{y}) = 0.$$

But

$$(2\mathbf{x} + 3\mathbf{y}) \bullet (2\mathbf{x} - 3\mathbf{y}) = 4\|\mathbf{x}\|^2 - 9\|\mathbf{y}\|^2 = 4\left(\frac{9}{4}\|\mathbf{y}\|^2\right) - 9\|\mathbf{y}\|^2 = 0.$$

358 We have $\forall \mathbf{v} \in \mathbb{R}^2$, $\mathbf{v} \bullet (\mathbf{a} - \mathbf{b}) = 0$. In particular, choosing $\mathbf{v} = \mathbf{a} - \mathbf{b}$, we gather

$$(\mathbf{a} - \mathbf{b}) \bullet (\mathbf{a} - \mathbf{b}) = \|\mathbf{a} - \mathbf{b}\|^2 = 0.$$

But the norm of a vector is 0 if and only if the vector is the 0 vector. Therefore $\mathbf{a} - \mathbf{b} = \mathbf{0}$, i.e., $\mathbf{a} = \mathbf{b}$.

359 We have

$$\begin{aligned}\|\mathbf{a} \pm \mathbf{b}\|^2 &= (\mathbf{a} \pm \mathbf{b}) \bullet (\mathbf{a} \pm \mathbf{b}) \\ &= \mathbf{a} \bullet \mathbf{a} \pm 2\mathbf{a} \bullet \mathbf{b} + \mathbf{b} \bullet \mathbf{b} \\ &= \|\mathbf{a}\|^2 \pm 2\mathbf{a} \bullet \mathbf{b} + \|\mathbf{b}\|^2,\end{aligned}$$

whence the result follows.

360 We have

$$\begin{aligned}\|\mathbf{u} + \mathbf{v}\|^2 - \|\mathbf{u} - \mathbf{v}\|^2 &= (\mathbf{u} + \mathbf{v}) \bullet (\mathbf{u} + \mathbf{v}) - (\mathbf{u} - \mathbf{v}) \bullet (\mathbf{u} - \mathbf{v}) \\ &= \mathbf{u} \bullet \mathbf{u} + 2\mathbf{u} \bullet \mathbf{v} + \mathbf{v} \bullet \mathbf{v} - (\mathbf{u} \bullet \mathbf{u} - 2\mathbf{u} \bullet \mathbf{v} + \mathbf{v} \bullet \mathbf{v}) \\ &= 4\mathbf{u} \bullet \mathbf{v},\end{aligned}$$

giving the result.

361 By definition

$$\begin{aligned}\text{proj}_{\mathbf{a}}^{\text{proj}_{\mathbf{x}}^{\mathbf{a}}} &= \frac{\text{proj}_{\mathbf{x}}^{\mathbf{a}} \bullet \mathbf{a}}{\|\mathbf{a}\|^2} \mathbf{a} \\ &= \frac{\frac{\mathbf{a} \bullet \mathbf{x}}{\|\mathbf{x}\|^2} \mathbf{x} \bullet \mathbf{a}}{\|\mathbf{a}\|^2} \mathbf{a} \\ &= \frac{(\mathbf{a} \bullet \mathbf{x})^2}{\|\mathbf{x}\|^2 \|\mathbf{a}\|^2} \mathbf{a},\end{aligned}$$

Since $0 \leq \frac{(\mathbf{a} \bullet \mathbf{x})^2}{\|\mathbf{x}\|^2 \|\mathbf{a}\|^2} \leq 1$ by the CBS Inequality, the result follows.

362 Clearly, if $\mathbf{a} = \mathbf{0}$ and $\lambda \neq 0$ then there are no solutions. If both $\mathbf{a} = \mathbf{0}$ and $\lambda = 0$, then the solution set is the whole space \mathbb{R}^2 . So assume that $\mathbf{a} \neq \mathbf{0}$. By Theorem 347, we may write $\mathbf{x} = \mathbf{u} + \mathbf{v}$ with $\text{proj}_{\mathbf{a}}^{\mathbf{x}} = \mathbf{u} \parallel \mathbf{a}$ and $\mathbf{v} \perp \mathbf{a}$. Thus there are infinitely many solutions, each of the form

$$\mathbf{x} = \mathbf{u} + \mathbf{v} = \frac{\mathbf{x} \cdot \mathbf{a}}{\|\mathbf{a}\|^2} \mathbf{a} + \mathbf{v} = \frac{\lambda}{\|\mathbf{a}\|^2} \mathbf{a} + \mathbf{v},$$

where $\mathbf{v} \in \mathbf{a}^\perp$.

372 Since $\mathbf{a} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$ is normal to $2x - y = 1$ and $\mathbf{b} = \begin{bmatrix} 1 \\ -3 \end{bmatrix}$ is normal to $x - 3y = 1$, the desired angle can be obtained by finding the angle between the normal vectors:

$$\widehat{(\mathbf{a}, \mathbf{b})} = \arccos \frac{\mathbf{a} \cdot \mathbf{b}}{\|\mathbf{a}\| \|\mathbf{b}\|} = \arccos \frac{5}{\sqrt{5} \cdot \sqrt{10}} = \arccos \frac{1}{\sqrt{2}} = \frac{\pi}{4}.$$

373 $2(x - 1) + (y + 1) = 0$ or $2x + y = 1$.

374 By Chasles' Rule $\overrightarrow{AA'} = \overrightarrow{AG} + \overrightarrow{GA'}$, $\overrightarrow{BB'} = \overrightarrow{BG} + \overrightarrow{GB'}$, and $\overrightarrow{CC'} = \overrightarrow{CG} + \overrightarrow{GC'}$. Thus

$$\begin{aligned} 0 &= \overrightarrow{AA'} + \overrightarrow{BB'} + \overrightarrow{CC'} \\ &= \overrightarrow{AG} + \overrightarrow{GA'} + \overrightarrow{BG} + \overrightarrow{GB'} + \overrightarrow{CG} + \overrightarrow{GC'} \\ &= -(\overrightarrow{GA} + \overrightarrow{GB} + \overrightarrow{GC}) + (\overrightarrow{GA'} + \overrightarrow{GB'} + \overrightarrow{GC'}) \\ &= \overrightarrow{GA'} + \overrightarrow{GB'} + \overrightarrow{GC'}, \end{aligned}$$

whence the result.

375 We have:

- ① The points F, A, D are collinear, and so \overrightarrow{FA} is parallel to \overrightarrow{FD} , meaning that there is $k \in \mathbb{R} \setminus \{0\}$ such that $\overrightarrow{FA} = k\overrightarrow{FD}$. Since the lines (AB) and (DC) are parallel, we obtain through Thales' Theorem that $\overrightarrow{FI} = k\overrightarrow{FJ}$ and $\overrightarrow{FB} = k\overrightarrow{FC}$. This gives

$$\overrightarrow{FA} - \overrightarrow{FI} = k(\overrightarrow{FD} - \overrightarrow{FJ}) \implies \overrightarrow{IA} = k\overrightarrow{JD}.$$

Similarly

$$\overrightarrow{FB} - \overrightarrow{FI} = k(\overrightarrow{FC} - \overrightarrow{FJ}) \implies \overrightarrow{IB} = k\overrightarrow{JC}.$$

Since I is the midpoint of $[A, B]$, $\overrightarrow{IA} + \overrightarrow{IB} = \mathbf{0}$, and thus $k(\overrightarrow{JC} + \overrightarrow{JD}) = \mathbf{0}$. Since $k \neq 0$, we have $\overrightarrow{JC} + \overrightarrow{JD} = \mathbf{0}$, meaning that J is the midpoint of $[C, D]$. Therefore the midpoints of $[A, B]$ and $[C, D]$ are aligned with F .

- ② Let J' be the intersection of the lines (EI) and (DC) . Let us prove that $J' = J$.

Since the points E, A, C are collinear, there is $l \neq 0$ such that $\overrightarrow{EA} = l\overrightarrow{EC}$. Since the lines (ab) and (DC) are parallel, we obtain via Thales' Theorem that $\overrightarrow{EI} = l\overrightarrow{EJ'}$ and $\overrightarrow{EB} = l\overrightarrow{ED}$. These equalities give

$$\begin{aligned} \overrightarrow{EA} - \overrightarrow{EI} &= l(\overrightarrow{EC} - \overrightarrow{EJ'}) \implies \overrightarrow{IA} = l\overrightarrow{J'C}, \\ \overrightarrow{EB} - \overrightarrow{EI} &= l(\overrightarrow{ED} - \overrightarrow{EJ'}) \implies \overrightarrow{IB} = l\overrightarrow{J'D}. \end{aligned}$$

Since I is the midpoint of $[A, B]$, $\overrightarrow{IA} + \overrightarrow{IB} = \mathbf{0}$, and thus $l(\overrightarrow{J'C} + \overrightarrow{J'D}) = \mathbf{0}$. Since $l \neq 0$, we deduce $\overrightarrow{J'C} + \overrightarrow{J'D} = \mathbf{0}$, that is, J' is the midpoint of $[C, D]$, and so $J' = J$.

376 We have:

- ① By Chasles' Rule

$$\overrightarrow{AE} = \frac{1}{4}\overrightarrow{AC} \iff \overrightarrow{AB} + \overrightarrow{BE} = \frac{1}{4}\overrightarrow{AC},$$

and

$$\overrightarrow{AF} = \frac{3}{4}\overrightarrow{AC} \iff \overrightarrow{AD} + \overrightarrow{DF} = \frac{3}{4}\overrightarrow{AC}.$$

Adding, and observing that since $ABCD$ is a parallelogram, $\vec{AB} = \vec{CD}$,

$$\begin{aligned}\vec{AB} + \vec{BE} + \vec{AD} + \vec{DF} - \vec{AC} &\iff \vec{BE} + \vec{DF} = \vec{AC} - \vec{AB} - \vec{AD} \\ &\iff \vec{BE} + \vec{DF} = \vec{AD} + \vec{DC} - \vec{AB} - \vec{AD} . \\ &\iff \vec{BE} = -\vec{DF}.\end{aligned}$$

The last equality shews that the lines (BE) and (DF) are parallel.

- ② Observe that $\vec{BJ} = \frac{1}{2}\vec{BC} = \frac{1}{2}\vec{AD} = \vec{AI} = -\vec{IA}$. Hence

$$\vec{IJ} = \vec{IA} + \vec{AB} + \vec{BJ} = \vec{AB},$$

proving that the lines (AB) and (IJ) are parallel.

Observe that

$$\vec{IE} = \vec{IA} + \vec{AE} = \frac{1}{2}\vec{DA} + \frac{1}{4}\vec{AC} = \frac{1}{2}\vec{CB} + \vec{FC} = \vec{CJ} + \vec{FC} = \vec{FC} + \vec{CJ} = \vec{FJ},$$

whence IEJF is a parallelogram.

- 377 Since $\vec{IE} = \frac{1}{3}\vec{ID}$ and $[I, D]$ is a median of $\triangle ABD$, E is the centre of gravity of $\triangle ABD$. Let M be the midpoint of $[B, D]$, and observe that M is the centre of the parallelogram, and so $2\vec{AM} = \vec{AB} + \vec{AD}$. Thus

$$\vec{AE} = \frac{2}{3}\vec{AM} = \frac{1}{3}(2\vec{AM}) = \frac{1}{3}(\vec{AB} + \vec{AD}).$$

To shew that A, C, E are collinear it is enough to notice that $\vec{AE} = \frac{1}{3}\vec{AC}$.

- 378 Suppose A, B, C are collinear and that $\frac{\| [A, B] \|}{\| [B, C] \|} = \frac{\lambda}{\mu}$. Then by the Section Formula 4.4,

$$\mathbf{b} = \frac{\lambda \mathbf{c} + \mu \mathbf{a}}{\lambda + \mu},$$

whence $\mu \mathbf{a} - (\lambda + \mu) \mathbf{b} + \lambda \mathbf{c} = 0$ and clearly $\mu - (\lambda + \mu) + \lambda = 0$. Thus we may take $\alpha = \mu$, $\beta = \lambda + \mu$, and $\gamma = \lambda$. Conversely, suppose that

$$\alpha \mathbf{a} + \beta \mathbf{b} + \gamma \mathbf{c} = 0, \quad \alpha + \beta + \gamma = 0$$

for some real numbers α, β, γ , not all zero. Assume without loss of generality that $\gamma \neq 0$. Otherwise we simply change the roles of γ , and α and β . Then $\gamma = -(\alpha + \beta) \neq 0$. Hence

$$\alpha \mathbf{a} + \beta \mathbf{b} = (\alpha + \beta) \mathbf{c} \implies \mathbf{c} = \frac{\alpha \mathbf{a} + \beta \mathbf{b}}{\alpha + \beta},$$

and thus $[O, C]$ divides $[A, B]$ into the ratio $\frac{\beta}{\alpha}$, and therefore, A, B, C are collinear.

- 379 Put $\vec{OX} = \mathbf{x}$ for $X \in \{A, A', B, B', C, C', L, M, N, V\}$. Using problem 378 we deduce

$$\mathbf{v} + \alpha \mathbf{a} + \alpha' \mathbf{a}' = 0, \quad 1 + \alpha + \alpha' = 0, \tag{A.1}$$

$$\mathbf{v} + \beta \mathbf{a} + \beta' \mathbf{a}' = 0, \quad 1 + \beta + \beta' = 0, \tag{A.2}$$

$$\mathbf{v} + \gamma \mathbf{a} + \gamma' \mathbf{a}' = 0, \quad 1 + \gamma + \gamma' = 0. \tag{A.3}$$

From A.2, A.3, and the Section Formula 4.4 we find

$$\frac{\beta \mathbf{b} - \gamma \mathbf{c}}{\beta - \gamma} = \frac{\beta' \mathbf{b}' - \gamma' \mathbf{c}'}{\beta' - \gamma'} = 1,$$

whence $(\beta - \gamma)\mathbf{l} = \beta \mathbf{b} - \gamma \mathbf{c}$. In a similar fashion, we deduce

$$(\gamma - \alpha)\mathbf{m} = \gamma \mathbf{c} - \alpha \mathbf{a},$$

$$(\alpha - \beta)\mathbf{n} = \alpha \mathbf{a} - \beta \mathbf{b}.$$

This gives

$$(\beta - \gamma)\mathbf{l} + (\gamma - \alpha)\mathbf{m} + (\alpha - \beta)\mathbf{n} = 0,$$

$$(\beta - \gamma) + (\gamma - \alpha) + (\alpha - \beta) = 0,$$

and appealing to problem 378 once again, we deduce that L, M, N are collinear.

391 [A] \overrightarrow{AS} , [B] \overrightarrow{AB} .

392 Put

$$\mathbf{a} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \times \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = (\mathbf{i} + \mathbf{j} + \mathbf{k}) \times (\mathbf{i} + \mathbf{j}) = \mathbf{j} - \mathbf{i} = \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}.$$

Then either

$$\frac{3\mathbf{a}}{\|\mathbf{a}\|} = \frac{3\mathbf{a}}{\sqrt{2}} = \begin{bmatrix} -\frac{3}{\sqrt{2}} \\ \frac{3}{\sqrt{2}} \\ 0 \end{bmatrix},$$

or

$$-\frac{3\mathbf{a}}{\|\mathbf{a}\|} = \begin{bmatrix} \frac{3}{\sqrt{2}} \\ -\frac{3}{\sqrt{2}} \\ 0 \end{bmatrix}$$

will satisfy the requirements.

393 The desired area is

$$\left\| \overrightarrow{PQ} \times \overrightarrow{PR} \right\| = \left\| \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix} \times \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} \right\| = \left\| \begin{bmatrix} -1 \\ -1 \\ -1 \end{bmatrix} \right\| = \sqrt{3}.$$

394 It is not associative, since $\mathbf{i} \times (\mathbf{i} \times \mathbf{j}) = \mathbf{i} \times \mathbf{k} = -\mathbf{j}$ but $(\mathbf{i} \times \mathbf{i}) \times \mathbf{j} = \mathbf{0} \times \mathbf{j} = \mathbf{0}$.

395 We have $\mathbf{x} \times \mathbf{x} = -\mathbf{x} \times \mathbf{x}$ by letting $\mathbf{y} = \mathbf{x}$ in 4.15. Thus $2\mathbf{x} \times \mathbf{x} = \mathbf{0}$ and hence $\mathbf{x} \times \mathbf{x} = \mathbf{0}$.

396 $2\mathbf{a} \times \mathbf{b}$

397

$$\mathbf{a} \times (\mathbf{x} \times \mathbf{b}) = \mathbf{b} \times (\mathbf{x} \times \mathbf{a}) \iff (\mathbf{a} \bullet \mathbf{b})\mathbf{x} - (\mathbf{a} \bullet \mathbf{x})\mathbf{b} = (\mathbf{b} \bullet \mathbf{a})\mathbf{x} - (\mathbf{b} \bullet \mathbf{x})\mathbf{a} \iff \mathbf{a} \bullet \mathbf{x} = \mathbf{b} \bullet \mathbf{x} = 0.$$

The answer is thus $\{\mathbf{x} : \mathbf{x} \in \mathbb{R}\mathbf{a} \times \mathbf{b}\}$.

398

$$\mathbf{x} = \frac{(\mathbf{a} \bullet \mathbf{b})\mathbf{a} + 6\mathbf{b} + 2\mathbf{a} \times \mathbf{c}}{12 + 2\|\mathbf{a}\|^2}$$

$$\mathbf{y} = \frac{(\mathbf{a} \bullet \mathbf{c})\mathbf{a} + 6\mathbf{c} + 3\mathbf{a} \times \mathbf{b}}{18 + 3\|\mathbf{a}\|^2}$$

399 Assume contrariwise that \mathbf{a} , \mathbf{b} , \mathbf{c} are three unit vectors in \mathbb{R}^3 such that the angle between any two of them is $> \frac{2\pi}{3}$. Then $\mathbf{a} \bullet \mathbf{b} < -\frac{1}{2}$, $\mathbf{b} \bullet \mathbf{c} < -\frac{1}{2}$, and $\mathbf{c} \bullet \mathbf{a} < -\frac{1}{2}$. Thus

$$\begin{aligned} \|\mathbf{a} + \mathbf{b} + \mathbf{c}\|^2 &= \|\mathbf{a}\|^2 + \|\mathbf{b}\|^2 + \|\mathbf{c}\|^2 \\ &\quad + 2\mathbf{a} \bullet \mathbf{b} + 2\mathbf{b} \bullet \mathbf{c} + 2\mathbf{c} \bullet \mathbf{a} \\ &< 1 + 1 + 1 - 1 - 1 - 1 \\ &= 0, \end{aligned}$$

which is impossible, since a norm of vectors is always ≥ 0 .

410 The vectors

$$\begin{bmatrix} \mathbf{a} - (-\mathbf{a}) \\ 0 - 1 \\ \mathbf{a} - 0 \end{bmatrix} = \begin{bmatrix} 2\mathbf{a} \\ -1 \\ \mathbf{a} \end{bmatrix}$$

and

$$\begin{bmatrix} 0 - (-\mathbf{a}) \\ 1 - 1 \\ 2\mathbf{a} - 0 \end{bmatrix} = \begin{bmatrix} \mathbf{a} \\ 0 \\ 2\mathbf{a} \end{bmatrix}$$

are coplanar. A vector normal to the plane is

$$\begin{bmatrix} 2\mathbf{a} \\ -1 \\ \mathbf{a} \end{bmatrix} \times \begin{bmatrix} \mathbf{a} \\ 0 \\ 2\mathbf{a} \end{bmatrix} = \begin{bmatrix} -2\mathbf{a} \\ -3\mathbf{a}^2 \\ \mathbf{a} \end{bmatrix}.$$

The equation of the plane is thus given by

$$\begin{bmatrix} -2\mathbf{a} \\ -3\mathbf{a}^2 \\ \mathbf{a} \end{bmatrix} \cdot \begin{bmatrix} x - \mathbf{a} \\ y - 0 \\ z - \mathbf{a} \end{bmatrix} = 0,$$

that is,

$$2ax + 3a^2y - az = a^2.$$

411 The vectorial form of the equation of the line is

$$\mathbf{r} = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} + t \begin{bmatrix} 1 \\ -2 \\ -1 \end{bmatrix}.$$

Since the line follows the direction of $\begin{bmatrix} 1 \\ -2 \\ -1 \end{bmatrix}$, this means that $\begin{bmatrix} 1 \\ -2 \\ -1 \end{bmatrix}$ is normal to the plane, and thus the equation of the desired plane is

$$(x - 1) - 2(y - 1) - (z - 1) = 0.$$

412 Observe that $(0, 0, 0)$ (as $0 = 2(0) = 3(0)$) is on the line, and hence on the plane. Thus the vector

$$\begin{bmatrix} 1 - 0 \\ -1 - 0 \\ -1 - 0 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \\ -1 \end{bmatrix}$$

lies on the plane. Now, if $x = 2y = 3z = t$, then $x = t, y = t/2, z = t/3$. Hence, the vectorial form of the equation of the line is

$$\mathbf{r} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} + t \begin{bmatrix} 1 \\ 1/2 \\ 1/3 \end{bmatrix} = t \begin{bmatrix} 1 \\ 1/2 \\ 1/3 \end{bmatrix}.$$

This means that $\begin{bmatrix} 1 \\ 1/2 \\ 1/3 \end{bmatrix}$ also lies on the plane, and thus

$$\begin{bmatrix} 1 \\ -1 \\ -1 \end{bmatrix} \times \begin{bmatrix} 1 \\ 1/2 \\ 1/3 \end{bmatrix} = \begin{bmatrix} 1/6 \\ -4/3 \\ 3/2 \end{bmatrix}$$

is normal to the plane. The desired equation is thus

$$\frac{1}{6}x - \frac{4}{3}y + \frac{3}{2}z = 0.$$

413 Put $ax = by = cz = t$, so $x = t/a; y = t/b; z = t/c$. The parametric equation of the line is

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = t \begin{bmatrix} 1/a \\ 1/b \\ 1/c \end{bmatrix}, \quad t \in \mathbb{R}.$$

Thus the vector $\begin{bmatrix} 1/a \\ 1/b \\ 1/c \end{bmatrix}$ is perpendicular to the plane. Therefore, the equation of the plane is

$$\begin{bmatrix} 1/a \\ 1/b \\ 1/c \end{bmatrix} \cdot \begin{bmatrix} x-1 \\ y-1 \\ z-1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix},$$

or

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = \frac{1}{a} + \frac{1}{b} + \frac{1}{c}.$$

We may also write this as

$$bcx + cay + abz = ab + bc + ca.$$

$$\begin{bmatrix} a \\ a^2 \\ a^2 \end{bmatrix}$$

414 A vector normal to the plane is $\begin{bmatrix} a \\ a^2 \\ a^2 \end{bmatrix}$. The line sought has the same direction as this vector, thus the equation of the

line is

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} + t \begin{bmatrix} a \\ a^2 \\ a^2 \end{bmatrix}, \quad t \in \mathbb{R}.$$

415 We have

$$x - z - y = 1 \implies -1 - y = 1 \implies y = -2.$$

Hence if $z = t$,

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} t - 1 \\ -2 \\ t \end{bmatrix} = \begin{bmatrix} -1 \\ -2 \\ 0 \end{bmatrix} + t \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}.$$

416 The vector

$$\begin{bmatrix} 2 - 1 \\ 1 - 0 \\ 1 - (-1) \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$$

lies on the plane. The vector

$$\begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} \times \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}$$

is normal to the plane. Hence the equation of the plane is

$$\begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix} \cdot \begin{bmatrix} x - 1 \\ y \\ z + 1 \end{bmatrix} = 0 \implies x + y - z = 2.$$

417 We have $\mathbf{c} \times \mathbf{a} = -\mathbf{i} + 2\mathbf{j}$ and $\mathbf{a} \times \mathbf{b} = 2\mathbf{k} - 3\mathbf{i}$. By Theorem 408, we have

$$\mathbf{b} \times \mathbf{c} = -\mathbf{a} \times \mathbf{b} - \mathbf{c} \times \mathbf{a} = -2\mathbf{k} + 3\mathbf{i} + \mathbf{i} - 2\mathbf{j} = 4\mathbf{i} - 2\mathbf{j} - 2\mathbf{k}.$$

418 $4x + 6y = 1$

419 There are 7 vertices ($V_0 = (0, 0, 0)$, $V_1 = (11, 0, 0)$, $V_2 = (0, 9, 0)$, $V_3 = (0, 0, 8)$, $V_4 = (0, 3, 8)$, $V_5 = (9, 0, 2)$, $V_6 = (4, 7, 0)$) and 11 edges (V_0V_1 , V_0V_2 , V_0V_3 , V_1V_5 , V_1V_6 , V_2V_4 , V_3V_4 , V_3V_5 , V_4V_5 , and V_4V_6).

427 Expand $\left| \sum_{i=1}^n \mathbf{a}_i \right|^2 = 0$.

428 Observe that $\sum_{k=1}^n 1 = n$. Then we have

$$n^2 = \left(\sum_{k=1}^n 1 \right)^2 = \left(\sum_{k=1}^n (a_k) \left(\frac{1}{a_k} \right) \right)^2 \leq \left(\sum_{k=1}^n a_k^2 \right) \left(\sum_{k=1}^n \frac{1}{a_k^2} \right),$$

giving the result.

429 This follows at once from the CBS Inequality by putting

$$\mathbf{v} = \begin{bmatrix} \frac{a_1}{1} \\ \frac{a_2}{2} \\ \dots \\ \frac{a_n}{n} \end{bmatrix}, \quad \mathbf{u} = \begin{bmatrix} 1 \\ 2 \\ \dots \\ n \end{bmatrix}$$

and noticing that

$$\sum_{k=1}^n k^2 = \frac{n(n+1)(2n+1)}{6}.$$

442 No, since $\mathbf{1}_F \mathbf{v} = \mathbf{v}$ is not fulfilled. For example

$$1 \cdot \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \cdot 1 \\ 0 \end{bmatrix} \neq \begin{bmatrix} 1 \\ 1 \end{bmatrix}.$$

443 We expand $(\mathbf{1}_F + \mathbf{1}_F)(\mathbf{a} + \mathbf{b})$ in two ways, first using 5.7 first and then 5.8, obtaining

$$(\mathbf{1}_F + \mathbf{1}_F)(\mathbf{a} + \mathbf{b}) = (\mathbf{1}_F + \mathbf{1}_F)\mathbf{a} + (\mathbf{1}_F + \mathbf{1}_F)\mathbf{b} = \mathbf{a} + \mathbf{a} + \mathbf{b} + \mathbf{b},$$

and then using 5.8 first and then 5.7, obtaining

$$(\mathbf{1}_F + \mathbf{1}_F)(\mathbf{a} + \mathbf{b}) = \mathbf{1}_F(\mathbf{a} + \mathbf{b}) + \mathbf{1}_F(\mathbf{a} + \mathbf{b}) = \mathbf{a} + \mathbf{b} + \mathbf{a} + \mathbf{b}.$$

We thus have the equality

$$\mathbf{a} + \mathbf{a} + \mathbf{b} + \mathbf{b} = \mathbf{a} + \mathbf{b} + \mathbf{a} + \mathbf{b}.$$

Cancelling \mathbf{a} from the left and \mathbf{b} from the right, we obtain

$$\mathbf{a} + \mathbf{b} = \mathbf{b} + \mathbf{a},$$

which is what we wanted to shew.

444 We must prove that each of the axioms of a vector space are satisfied. Clearly if $(x, y, \alpha) \in \mathbb{R}^+ \times \mathbb{R}^+ \times \mathbb{R}$ then $x \oplus y = xy > 0$ and $\alpha \otimes x = x^\alpha > 0$, so V is closed under vector addition and scalar multiplication. Commutativity and associativity of vector addition are obvious.

Let \mathbf{A} be additive identity. Then we need

$$x \oplus \mathbf{A} = x \implies x\mathbf{A} = x \implies \mathbf{A} = \mathbf{1}.$$

Thus the additive identity is $\mathbf{1}$. Suppose \mathbf{I} is the additive inverse of x . Then

$$x \oplus \mathbf{I} = \mathbf{1} \implies x\mathbf{I} = \mathbf{1} \implies \mathbf{I} = \frac{1}{x}.$$

Hence the additive inverse of x is $\frac{1}{x}$.

Now

$$\alpha \otimes (x \oplus y) = (xy)^\alpha = x^\alpha y^\alpha = x^\alpha \oplus y^\alpha = (\alpha \otimes x) \oplus (\alpha \otimes y),$$

and

$$(\alpha + \beta) \otimes x = x^{\alpha+\beta} = x^\alpha x^\beta = (x^\alpha) \oplus (x^\beta) = (\alpha \otimes x) \oplus (\beta \otimes x),$$

whence the distributive laws hold.

Finally,

$$\mathbf{1} \otimes x = x^1 = x,$$

and

$$\alpha \otimes (\beta \otimes x) = (\beta \otimes x)^\alpha = (x^\beta)^\alpha = x^{\alpha\beta} = (\alpha\beta) \otimes x,$$

and the last two axioms also hold.

445 \mathbb{C} is a vector space over \mathbb{R} , the proof is trivial. But \mathbb{R} is not a vector space over \mathbb{C} , since, for example taking i as a scalar (from \mathbb{C}) and 1 as a vector (from \mathbb{R}) the scalar multiple $i \cdot 1 = i \notin \mathbb{R}$ and so there is no closure under scalar multiplication.

446 One example is

$$(\mathbb{Z}_2)^3 = \left\{ \begin{bmatrix} \bar{0} \\ \bar{0} \\ \bar{0} \end{bmatrix}, \begin{bmatrix} \bar{0} \\ \bar{0} \\ \bar{1} \end{bmatrix}, \begin{bmatrix} \bar{0} \\ \bar{1} \\ \bar{0} \end{bmatrix}, \begin{bmatrix} \bar{0} \\ \bar{1} \\ \bar{1} \end{bmatrix}, \begin{bmatrix} \bar{1} \\ \bar{0} \\ \bar{0} \end{bmatrix}, \begin{bmatrix} \bar{1} \\ \bar{0} \\ \bar{1} \end{bmatrix}, \begin{bmatrix} \bar{1} \\ \bar{1} \\ \bar{0} \end{bmatrix}, \begin{bmatrix} \bar{1} \\ \bar{1} \\ \bar{1} \end{bmatrix} \right\}.$$

Addition is the natural element-wise addition and scalar multiplication is ordinary element-wise scalar multiplication.

447 One example is

$$(\mathbb{Z}_3)^2 = \left\{ \begin{bmatrix} \bar{0} \\ \bar{0} \end{bmatrix}, \begin{bmatrix} \bar{0} \\ \bar{1} \end{bmatrix}, \begin{bmatrix} \bar{0} \\ \bar{2} \end{bmatrix}, \begin{bmatrix} \bar{1} \\ \bar{0} \end{bmatrix}, \begin{bmatrix} \bar{1} \\ \bar{1} \end{bmatrix}, \begin{bmatrix} \bar{1} \\ \bar{2} \end{bmatrix}, \begin{bmatrix} \bar{2} \\ \bar{0} \end{bmatrix}, \begin{bmatrix} \bar{2} \\ \bar{1} \end{bmatrix}, \begin{bmatrix} \bar{2} \\ \bar{2} \end{bmatrix} \right\}.$$

Addition is the natural element-wise addition and scalar multiplication is ordinary element-wise scalar multiplication.

454 Take $\alpha \in \mathbb{R}$ and

$$\mathbf{x} = \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} \in \mathbf{X}, \quad a - b - 3d = 0, \quad \mathbf{y} = \begin{bmatrix} a' \\ b' \\ c' \\ d' \end{bmatrix} \in \mathbf{X}, \quad a' - b' - 3d' = 0.$$

Then

$$\mathbf{x} + \alpha \mathbf{y} = \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} + \alpha \begin{bmatrix} a' \\ b' \\ c' \\ d' \end{bmatrix} = \begin{bmatrix} a + \alpha a' \\ b + \alpha b' \\ c + \alpha c' \\ d + \alpha d' \end{bmatrix}.$$

Observe that

$$(a + \alpha a') - (b + \alpha b') - 3(d + \alpha d') = (a - b - 3d) + \alpha(a' - b' - 3d') = 0 + \alpha 0 = 0,$$

meaning that $\mathbf{x} + \alpha \mathbf{y} \in \mathbf{X}$, and so \mathbf{X} is a vector subspace of \mathbb{R}^4 .

455 Take

$$\mathbf{u} = \begin{bmatrix} a_1 \\ 2a_1 - 3b_1 \\ 5b_1 \\ a_1 + 2b_1 \\ a_1 \end{bmatrix}, \quad \mathbf{v} = \begin{bmatrix} a_2 \\ 2a_2 - 3b_2 \\ 5b_2 \\ a_2 + 2b_2 \\ a_2 \end{bmatrix}, \quad \alpha \in \mathbb{R}.$$

Put $s = \mathbf{a}_1 + \alpha\mathbf{a}_2$, $t = \mathbf{b}_1 + \alpha\mathbf{b}_2$. Then

$$\mathbf{u} + \alpha\mathbf{v} = \begin{bmatrix} \mathbf{a}_1 + \alpha\mathbf{a}_2 \\ 2(\mathbf{a}_1 + \alpha\mathbf{a}_2) - 3(\mathbf{b}_1 + \alpha\mathbf{b}_2) \\ 5(\mathbf{b}_1 + \alpha\mathbf{b}_2) \\ (\mathbf{a}_1 + \alpha\mathbf{a}_2) + 2(\mathbf{b}_1 + \alpha\mathbf{b}_2) \\ \mathbf{a}_1 + \alpha\mathbf{a}_2 \end{bmatrix} = \begin{bmatrix} s \\ 2s - 3t \\ 5t \\ s + 2t \\ s \end{bmatrix} \in X,$$

since this last matrix has the basic shape of matrices in X . This shows that X is a vector subspace of \mathbb{R}^5 .

456 Take $(\mathbf{u}, \mathbf{v}) \in X^2$ and $\alpha \in \mathbb{R}$. Then

$$\mathbf{a} \bullet (\mathbf{u} + \alpha\mathbf{v}) = \mathbf{a} \bullet \mathbf{u} + \alpha\mathbf{a} \bullet \mathbf{v} = 0 + 0 = 0,$$

proving that X is a vector subspace of \mathbb{R}^n .

457 Take $(\mathbf{u}, \mathbf{v}) \in X^2$ and $\alpha \in \mathbb{R}$. Then

$$\mathbf{a} \times (\mathbf{u} + \alpha\mathbf{v}) = \mathbf{a} \times \mathbf{u} + \alpha\mathbf{a} \times \mathbf{v} = 0 + \alpha 0 = 0,$$

proving that X is a vector subspace of \mathbb{R}^n .

462 We show that some of the properties in the definition of vector subspace fail to hold in these sets.

❶ Take $\mathbf{x} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$, $\alpha = 2$. Then $\mathbf{x} \in V$ but $2\mathbf{x} = \begin{bmatrix} 0 \\ 2 \\ 0 \end{bmatrix} \notin V$ as $0^2 + 2^2 = 4 \neq 1$. So V is not closed under scalar multiplication.

❷ Take $\mathbf{x} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$, $\mathbf{y} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$. Then $\mathbf{x} \in W, \mathbf{y} \in W$ but $\mathbf{x} + \mathbf{y} = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \notin W$ as $1 \cdot 1 = 1 \neq 0$. Hence W is not closed under vector addition.

❸ Take $\mathbf{x} = \begin{bmatrix} -1 & 1 \\ 0 & 0 \end{bmatrix}$. Then $\mathbf{x} \in Z$ but $-\mathbf{x} = -\begin{bmatrix} -1 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & -1 \\ 0 & 0 \end{bmatrix} \notin Z$ as $1 + (-1)^2 = 2 \neq 0$. So Z is not closed under scalar multiplication.

463 Assume $U_1 \not\subseteq U_2$ and $U_2 \not\subseteq U_1$. Take $\mathbf{v} \in U_2 \setminus U_1$ (which is possible because $U_2 \not\subseteq U_1$) and $\mathbf{u} \in U_1 \setminus U_2$ (which is possible because $U_1 \not\subseteq U_2$). If $\mathbf{u} + \mathbf{v} \in U_1$, then—as $-\mathbf{u}$ is also in U_1 —the sum of two vectors in U_1 must also be in U_1 giving

$$\mathbf{u} + \mathbf{v} - \mathbf{u} = \mathbf{v} \in U_1,$$

a contradiction. Similarly if $\mathbf{u} + \mathbf{v} \in U_2$, then—as $-\mathbf{v}$ also in U_2 —the sum of two vectors in U_2 must also be in U_1 giving

$$\mathbf{u} + \mathbf{v} - \mathbf{u} = \mathbf{v} \in U_2,$$

another contradiction. Hence either $U_1 \subseteq U_2$ or $U_2 \subseteq U_1$ (or possibly both).

464 Assume contrariwise that $V = U_1 \cup U_2 \cup \dots \cup U_k$ is the shortest such list. Since the U_i are proper subspaces, $k > 1$. Choose $\mathbf{x} \in U_1, \mathbf{x} \notin U_2 \cup \dots \cup U_k$ and choose $\mathbf{y} \notin U_1$. Put $L = \{\mathbf{y} + \alpha\mathbf{x} | \alpha \in \mathbb{F}\}$. Claim: $L \cap U_1 = \emptyset$. For if $\mathbf{u} \in L \cap U_1$ then $\exists \alpha_0 \in \mathbb{F}$ with $\mathbf{u} = \mathbf{y} + \alpha_0\mathbf{x}$ and so $\mathbf{y} = \mathbf{u} - \alpha_0\mathbf{x} \in U_1$, a contradiction. So L and U_1 are disjoint.

We now shew that L has at most one vector in common with U_j , $2 \leq j \leq k$. For, if there were two elements of \mathbb{F} , $a \neq b$ with $y + ax, y + bx \in U_j, j \geq 2$ then

$$(a - b)x = (y + ax) - (y + bx) \in U_j,$$

contrary to the choice of x .

Conclusion: since \mathbb{F} is infinite, L is infinite. But we have shewn that L can have at most one element in common with the U_j . This means that there are not enough U_j to go around to cover the whole of L . So V cannot be a finite union of proper subspaces.

465 Take $\mathbb{F} = \mathbb{Z}_2$, $V = \mathbb{F} \times \mathbb{F}$. Then V has the four elements

$$\begin{bmatrix} \bar{0} \\ \bar{0} \end{bmatrix}, \begin{bmatrix} \bar{0} \\ \bar{1} \end{bmatrix}, \begin{bmatrix} \bar{1} \\ \bar{0} \end{bmatrix}, \begin{bmatrix} \bar{1} \\ \bar{1} \end{bmatrix},$$

with the following subspaces

$$V_1 = \left\{ \begin{bmatrix} \bar{0} \\ \bar{0} \end{bmatrix}, \begin{bmatrix} \bar{0} \\ \bar{1} \end{bmatrix} \right\}, \quad V_2 = \left\{ \begin{bmatrix} \bar{0} \\ \bar{0} \end{bmatrix}, \begin{bmatrix} \bar{1} \\ \bar{0} \end{bmatrix} \right\}, \quad V_3 = \left\{ \begin{bmatrix} \bar{0} \\ \bar{0} \end{bmatrix}, \begin{bmatrix} \bar{1} \\ \bar{1} \end{bmatrix} \right\}.$$

It is easy to verify that these subspaces satisfy the conditions of the problem.

475 If

$$a \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + b \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + c \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = 0,$$

then

$$\begin{bmatrix} a+b+c \\ b+c \\ c \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}.$$

This clearly entails that $c = b = a = 0$, and so the family is free.

476 Assume

$$a \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + b \begin{bmatrix} 1 \\ 1 \\ -1 \\ -1 \end{bmatrix} + c \begin{bmatrix} 1 \\ -1 \\ 1 \\ -1 \end{bmatrix} + d \begin{bmatrix} 1 \\ 1 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}.$$

Then

$$a + b + c + d = 0,$$

$$a + b - c + d = 0,$$

$$a - b + c = 0,$$

$$a - b - c + d = 0.$$

Subtracting the second equation from the first, we deduce $2c = 0$, that is, $c = 0$. Subtracting the third equation from the fourth, we deduce $-2c + d = 0$ or $d = 0$. From the first and third equations, we then deduce $a + b = 0$ and $a - b = 0$, which entails $a = b = 0$. In conclusion, $a = b = c = d = 0$.

Now, put

$$x \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + y \begin{bmatrix} 1 \\ 1 \\ -1 \\ -1 \end{bmatrix} + z \begin{bmatrix} 1 \\ -1 \\ 1 \\ -1 \end{bmatrix} + w \begin{bmatrix} 1 \\ 1 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 1 \\ 1 \end{bmatrix}.$$

Then

$$x + y + z + w = 1,$$

$$x + y - z + w = 2,$$

$$x - y + z = 1,$$

$$x - y - z + w = 1.$$

Solving as before, we find

$$2 \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + \frac{1}{2} \begin{bmatrix} 1 \\ 1 \\ -1 \\ -1 \end{bmatrix} - \frac{1}{2} \begin{bmatrix} 1 \\ -1 \\ 1 \\ -1 \end{bmatrix} - \begin{bmatrix} 1 \\ 1 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ 1 \\ 1 \end{bmatrix}.$$

477 Since \mathbf{a}, \mathbf{b} are linearly independent, none of them is 0. Assume that there are $(\alpha, \beta, \gamma) \in \mathbb{R}^3$ such that

$$\alpha \mathbf{a} + \beta \mathbf{b} + \gamma \mathbf{a} \times \mathbf{b} = 0. \quad (\text{A.4})$$

Since $\mathbf{a} \cdot (\mathbf{a} \times \mathbf{b}) = 0$, taking the dot product of A.4 with \mathbf{a} yields $\alpha \|\mathbf{a}\|^2 = 0$, which means that $\alpha = 0$, since $\|\mathbf{a}\| \neq 0$. Similarly, we take the dot product with \mathbf{b} and $\mathbf{a} \times \mathbf{b}$ obtaining respectively, $\beta = 0$ and $\gamma = 0$. This establishes linear independence.

478 Assume that

$$\lambda_1 \mathbf{a}_1 + \dots + \lambda_k \mathbf{a}_k = 0.$$

Taking the dot product with \mathbf{a}_j and using the fact that $\mathbf{a}_i \cdot \mathbf{a}_j = 0$ for $i \neq j$ we obtain

$$0 = \mathbf{0} \cdot \mathbf{a}_j = \lambda_j \mathbf{a}_j \cdot \mathbf{a}_j = \lambda_j \|\mathbf{a}_j\|^2.$$

Since $\mathbf{a}_j \neq 0 \implies \|\mathbf{a}_j\|^2 \neq 0$, we must have $\lambda_j = 0$. Thus the only linear combination giving the zero vector is the trivial linear combination, which proves that the vectors are linearly independent.

481 We have

$$(\mathbf{v}_1 + \mathbf{v}_2) - (\mathbf{v}_2 + \mathbf{v}_3) + (\mathbf{v}_3 + \mathbf{v}_4) - (\mathbf{v}_4 + \mathbf{v}_1) = 0,$$

a non-trivial linear combination of these vectors equalling the zero-vector.

483 Yes. Suppose that $a + b\sqrt{2} = 0$ is a non-trivial linear combination of 1 and $\sqrt{2}$ with rational numbers a and b . If one of a, b is different from 0 then so is the other. Hence

$$a + b\sqrt{2} = 0 \implies \sqrt{2} = -\frac{b}{a}.$$

The sinistral side of the equality $\sqrt{2} = -\frac{b}{a}$ is irrational whereas the dextral side is rational, a contradiction.

484 No. The representation $2 \cdot 1 + (-\sqrt{2})\sqrt{2} = 0$ is a non-trivial linear combination of 1 and $\sqrt{2}$.

485 1. Assume that

$$a + b\sqrt{2} + c\sqrt{3} = 0, \quad a, b, c \in \mathbb{Q}, \quad a^2 + b^2 + c^2 \neq 0.$$

If $ac \neq 0$, then

$$b\sqrt{2} = -a - c\sqrt{3} \Leftrightarrow 2b^2 = a^2 + 2ac\sqrt{3} + 3c^2 \Leftrightarrow \frac{2b^2 - a^2 - 3c^2}{2ac} = \sqrt{3}.$$

The dextral side of the last implication is irrational, whereas the sinistral side is rational. Thus it must be the case that $ac = 0$. If $a = 0, c \neq 0$ then

$$b\sqrt{2} + c\sqrt{3} = 0 \Leftrightarrow -\frac{b}{c} = \sqrt{\frac{3}{2}},$$

and again the dextral side is irrational and the sinistral side is rational. Thus if $a = 0$ then also $c = 0$. We can similarly prove that $c = 0$ entails $a = 0$. Thus we have

$$b\sqrt{2} = 0,$$

which means that $b = 0$. Therefore

$$a + b\sqrt{2} + c\sqrt{3} = 0, a, b, c \in \mathbb{Q}, \Leftrightarrow a = b = c = 0.$$

This proves that $\{1, \sqrt{2}, \sqrt{3}\}$ are linearly independent over \mathbb{Q} .

2. Rationalising denominators,

$$\begin{aligned} \frac{1}{1-\sqrt{2}} + \frac{2}{\sqrt{12}-2} &= \frac{1+\sqrt{2}}{1-2} + \frac{2\sqrt{12}+4}{12-4} \\ &= -1-\sqrt{2} + \frac{1}{2}\sqrt{3} + \frac{1}{2} \\ &= -\frac{1}{2}-\sqrt{2} + \frac{1}{2}\sqrt{3}. \end{aligned}$$

486 Assume that

$$ae^x + be^{2x} + ce^{3x} = 0.$$

Then

$$c = -ae^{-2x} - be^{-x}.$$

Letting $x \rightarrow +\infty$, we obtain $c = 0$. Thus

$$ae^x + be^{2x} = 0,$$

and so

$$b = -ae^{-x}.$$

Again, letting $x \rightarrow +\infty$, we obtain $b = 0$. This yields

$$ae^x = 0.$$

Since the exponential function never vanishes, we deduce that $a = 0$. Thus $a = b = c = 0$ and the family is linearly independent over \mathbb{R} .

487 This follows at once from the identity

$$\cos 2x = \cos^2 x - \sin^2 x,$$

which implies

$$\cos 2x - \cos^2 x + \sin^2 x = 0.$$

500 Given an arbitrary polynomial

$$p(x) = a + bx + cx^2 + dx^3,$$

we must shew that there are real numbers s, t, u, v such that

$$p(x) = s + t(1+x) + u(1+x)^2 + v(1+x)^3.$$

In order to do this we find the Taylor expansion of p around $x = -1$. Letting $x = -1$ in this last equality,

$$s = p(-1) = a - b + c - d \in \mathbb{R}.$$

Now,

$$p'(x) = b + 2cx + 3dx^2 = t + 2u(1+x) + 3v(1+x)^2.$$

Letting $x = -1$ we find

$$t = p'(-1) = b - 2c + 3d \in \mathbb{R}.$$

Again,

$$p''(x) = 2c + 6dx = 2u + 6v(1+x).$$

Letting $x = -1$ we find

$$u = p''(-1) = c - 3d \in \mathbb{R}.$$

Finally,

$$p'''(x) = 6d = 6v,$$

so we let $v = d \in \mathbb{R}$. In other words, we have

$$p(x) = a + bx + cx^2 + dx^3 = (a - b + c - d) + (b - 2c + 3d)(1+x) + (c - 3d)(1+x)^2 + d(1+x)^3.$$

501 Assume contrariwise that

$$\begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix} = \mathbf{a} \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} + \mathbf{b} \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix}.$$

Then we must have

$$\mathbf{a} = 1,$$

$$\mathbf{b} = 1,$$

$$-\mathbf{a} - \mathbf{b} = -1,$$

which is impossible. Thus $\begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}$ is not a linear combination of $\begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix}$ and hence is not in $\text{span} \left(\begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix} \right)$.

502 It is

$$\mathbf{a} \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + \mathbf{b} \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} + \mathbf{c} \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} = \begin{bmatrix} \mathbf{a} & \mathbf{c} \\ -\mathbf{c} & \mathbf{b} \end{bmatrix},$$

i.e., this family spans the set of all skew-symmetric 2×2 matrices over \mathbb{R} .

515 We have

$$\begin{bmatrix} \mathbf{a} \\ 2\mathbf{a} - 3\mathbf{b} \\ 5\mathbf{b} \\ \mathbf{a} + 2\mathbf{b} \\ \mathbf{a} \end{bmatrix} = \mathbf{a} \begin{bmatrix} 1 \\ 2 \\ 0 \\ 1 \\ 1 \end{bmatrix} + \mathbf{b} \begin{bmatrix} 0 \\ -3 \\ 5 \\ 2 \\ 0 \end{bmatrix},$$

so clearly the family

$$\left\{ \begin{bmatrix} 1 \\ 2 \\ 0 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ -3 \\ 5 \\ 2 \\ 0 \end{bmatrix} \right\}$$

spans the subspace. To shew that this is a linearly independent family, assume that

$$\mathbf{a} \begin{bmatrix} 1 \\ 2 \\ 0 \\ 1 \\ 1 \end{bmatrix} + \mathbf{b} \begin{bmatrix} 0 \\ -3 \\ 5 \\ 2 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}.$$

Then it follows clearly that $\mathbf{a} = \mathbf{b} = \mathbf{0}$, and so this is a linearly independent family. Conclusion:

$$\left\{ \begin{bmatrix} 1 \\ 2 \\ 0 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ -3 \\ 5 \\ 2 \\ 0 \end{bmatrix} \right\}$$

is a basis for the subspace.

516 Suppose

$$\begin{aligned} \mathbf{0} &= \mathbf{a}(\mathbf{v}_1 + \mathbf{v}_2) + \mathbf{b}(\mathbf{v}_2 + \mathbf{v}_3) + \mathbf{c}(\mathbf{v}_3 + \mathbf{v}_4) + \mathbf{d}(\mathbf{v}_4 + \mathbf{v}_5) + \mathbf{f}(\mathbf{v}_5 + \mathbf{v}_1) \\ &= (\mathbf{a} + \mathbf{f})\mathbf{v}_1 + (\mathbf{a} + \mathbf{b})\mathbf{v}_2 + (\mathbf{b} + \mathbf{c})\mathbf{v}_3 + (\mathbf{c} + \mathbf{d})\mathbf{v}_4 + (\mathbf{d} + \mathbf{f})\mathbf{v}_5. \end{aligned}$$

Since $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_5\}$ are linearly independent, we have

$$\begin{aligned} \mathbf{a} + \mathbf{f} &= \mathbf{0}, \\ \mathbf{a} + \mathbf{b} &= \mathbf{0} \\ \mathbf{b} + \mathbf{c} &= \mathbf{0} \\ \mathbf{c} + \mathbf{d} &= \mathbf{0} \\ \mathbf{d} + \mathbf{f} &= \mathbf{0}. \end{aligned}$$

Solving we find $\mathbf{a} = \mathbf{b} = \mathbf{c} = \mathbf{d} = \mathbf{f} = \mathbf{0}$, which means that the

$$\{\mathbf{v}_1 + \mathbf{v}_2, \mathbf{v}_2 + \mathbf{v}_3, \mathbf{v}_3 + \mathbf{v}_4, \mathbf{v}_4 + \mathbf{v}_5, \mathbf{v}_5 + \mathbf{v}_1\}$$

are linearly independent. Since the dimension of \mathbf{V} is 5, and we have 5 linearly independent vectors, they must also be a basis for \mathbf{V} .

517 The matrix of coefficients is already in echelon form. The dimension of the solution space is $n - 1$ and the following vectors in \mathbb{R}^{2n} form a basis for the solution space

$$\mathbf{a}_1 = \begin{bmatrix} -1 \\ 1 \\ 0 \\ \vdots \\ 0 \\ -1 \\ 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \quad \mathbf{a}_2 = \begin{bmatrix} -1 \\ 0 \\ 1 \\ \vdots \\ 0 \\ -1 \\ 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \dots, \quad \mathbf{a}_{n-1} = \begin{bmatrix} -1 \\ 0 \\ \dots \\ 1 \\ -1 \\ 0 \\ \vdots \\ 0 \\ 1 \end{bmatrix}.$$

(The “second” -1 occurs on the n -th position. The 1 ’s migrate from the 2nd and $n + 1$ -th position on \mathbf{a}_1 to the $n - 1$ -th and $2n$ -th position on \mathbf{a}_{n-1} .)

518 Take $(\mathbf{u}, \mathbf{v}) \in X^2$ and $\alpha \in \mathbb{R}$. Then

$$\mathbf{u} = \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix}, \quad b + 2c = 0, \quad \mathbf{v} = \begin{bmatrix} a' \\ b' \\ c' \\ d' \end{bmatrix}, \quad b' + 2c' = 0.$$

We have

$$\mathbf{u} + \alpha \mathbf{v} = \begin{bmatrix} a + \alpha a' \\ b + \alpha b' \\ c + \alpha c' \\ d + \alpha d' \end{bmatrix},$$

and to demonstrate that $\mathbf{u} + \alpha \mathbf{v} \in X$ we need to shew that $(b + \alpha b') + 2(c + \alpha c') = 0$. But this is easy, as

$$(b + \alpha b') + 2(c + \alpha c') = (b + 2c) + \alpha(b' + 2c') = 0 + \alpha 0 = 0.$$

Now

$$\begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} a \\ -2c \\ c \\ d \end{bmatrix} = a \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + c \begin{bmatrix} 0 \\ -2 \\ 1 \\ 0 \end{bmatrix} + d \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}$$

It is clear that

$$\begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ -2 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}$$

are linearly independent and span X . They thus constitute a basis for X .

519 As a basis we may take the $\frac{n(n+1)}{2}$ matrices $E_{ij} \in M_n(\mathbb{F})$ for $1 \leq i \leq j \leq n$.

520 $\dim X = 2$, as basis one may take $\{\mathbf{v}_1, \mathbf{v}_2\}$.

521 $\dim X = 3$, as basis one may take $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$.

522 $\dim X = 3$, as basis one may take $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$.

523 Since $a \perp a \times x = b$, there are no solutions if $a \cdot b \neq 0$. Neither are there solutions if $a = 0$ and $b \neq 0$. If both $a = b = 0$, then the solution set is the whole of \mathbb{R}^3 . Assume thus that $a \cdot b = 0$ and that a and b are linearly independent. Then $a, b, a \times b$ are linearly independent, and so they constitute a basis for \mathbb{R}^3 . Any $x \in \mathbb{R}^3$ can be written in the form

$$x = \alpha a + \beta b + \gamma a \times b.$$

We then have

$$\begin{aligned}
 \mathbf{b} &= \mathbf{a} \times \mathbf{x} \\
 &= \beta \mathbf{a} \times \mathbf{b} + \gamma \mathbf{a} \times (\mathbf{a} \times \mathbf{b}) \\
 &= \beta \mathbf{a} \times \mathbf{b} + \gamma ((\mathbf{a} \cdot \mathbf{b})\mathbf{a} - (\mathbf{a} \cdot \mathbf{a})\mathbf{b}) \\
 &= \beta \mathbf{a} \times \mathbf{b} - \gamma (\mathbf{a} \cdot \mathbf{a})\mathbf{b} \\
 &= \beta \mathbf{a} \times \mathbf{b} - \gamma \|\mathbf{a}\|^2 \mathbf{b},
 \end{aligned}$$

from where

$$\beta \mathbf{a} \times \mathbf{b} + (-\gamma \|\mathbf{a}\|^2 - 1)\mathbf{b} = 0,$$

which means that $\beta = 0$ and $\gamma = -\frac{1}{\|\mathbf{a}\|^2}$, since $\mathbf{a}, \mathbf{b}, \mathbf{a} \times \mathbf{b}$ are linearly independent. Thus

$$\mathbf{x} = \alpha \mathbf{a} - \frac{1}{\|\mathbf{a}\|^2} \mathbf{a} \times \mathbf{b}$$

in this last case.

531 1. It is enough to prove that the matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & -1 & 1 \end{bmatrix}$$

is invertible. But an easy computation shews that

$$\mathbf{A}^2 = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & -1 & 1 \end{bmatrix}^2 = 4\mathbf{I}_4,$$

whence the inverse sought is

$$\mathbf{A}^{-1} = \frac{1}{4}\mathbf{A} = \frac{1}{4} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & -1 & 1 \end{bmatrix} = \begin{bmatrix} 1/4 & 1/4 & 1/4 & 1/4 \\ 1/4 & 1/4 & -1/4 & -1/4 \\ 1/4 & -1/4 & 1/4 & -1/4 \\ 1/4 & -1/4 & -1/4 & 1/4 \end{bmatrix}.$$

2. Since the \mathbf{a}_k are four linearly independent vectors in \mathbb{R}^4 and $\dim \mathbb{R}^4 = 4$, they form a basis for \mathbb{R}^4 . Now, we want to solve

$$\mathbf{A} \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & -1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 1 \\ 1 \end{bmatrix}$$

and so

$$\begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} = \mathbf{A}^{-1} \begin{bmatrix} 1 \\ 2 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1/4 & 1/4 & 1/4 & 1/4 \\ 1/4 & 1/4 & -1/4 & -1/4 \\ 1/4 & -1/4 & 1/4 & -1/4 \\ 1/4 & -1/4 & -1/4 & 1/4 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 5/4 \\ 1/4 \\ -1/4 \\ -1/4 \end{bmatrix}.$$

It follows that

$$\begin{bmatrix} 1 \\ 2 \\ 1 \\ 1 \end{bmatrix} = \frac{5}{4} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + \frac{1}{4} \begin{bmatrix} 1 \\ 1 \\ -1 \\ -1 \end{bmatrix} - \frac{1}{4} \begin{bmatrix} 1 \\ -1 \\ 1 \\ -1 \end{bmatrix} - \frac{1}{4} \begin{bmatrix} 1 \\ -1 \\ -1 \\ 1 \end{bmatrix}.$$

The coordinates sought are

$$\left(\frac{5}{4}, \frac{1}{4}, -\frac{1}{4}, -\frac{1}{4} \right).$$

3. Since we have

$$\begin{bmatrix} 1 \\ 2 \\ 1 \\ 1 \end{bmatrix} = \frac{5}{4} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} - \frac{1}{4} \begin{bmatrix} 1 \\ -1 \\ 1 \\ -1 \end{bmatrix} + \frac{1}{4} \begin{bmatrix} 1 \\ 1 \\ -1 \\ -1 \end{bmatrix} - \frac{1}{4} \begin{bmatrix} 1 \\ -1 \\ -1 \\ 1 \end{bmatrix},$$

the coordinates sought are

$$\left(\frac{5}{4}, -\frac{1}{4}, \frac{1}{4}, -\frac{1}{4} \right).$$

532 [1] $a = 1$, [2] $(\mathbf{A}(a))^{-1} = \begin{bmatrix} \frac{1}{a-1} & 0 & 0 & -\frac{1}{a-1} \\ -1 & 1-a & -1 & a+1 \\ -\frac{1}{a-1} & -1 & 0 & \frac{a}{a-1} \\ 1 & a & 1 & -a-1 \end{bmatrix}$ [3]

$$\begin{bmatrix} 0 & \frac{1}{a-1} & \frac{1}{a-1} & \frac{1}{a-1} \\ 0 & -a-1 & -a & -1 \\ 0 & -\frac{a}{a-1} & -\frac{a}{a-1} & -\frac{1}{a-1} \\ 1 & 2+a & a+1 & 1 \end{bmatrix}$$

539 Let $\alpha \in \mathbb{R}$. Then

$$\begin{aligned} L \begin{bmatrix} x + \alpha a \\ y + \alpha b \\ z + \alpha c \end{bmatrix} &= \begin{bmatrix} (x + \alpha a) - (y + \alpha b) - (z + \alpha c) \\ (x + \alpha a) + (y + \alpha b) + (z + \alpha c) \\ z + \alpha c \end{bmatrix} \\ &= \begin{bmatrix} x - y - z \\ x + y + z \\ z \end{bmatrix} + \alpha \begin{bmatrix} a - b - c \\ a + b + c \\ c \end{bmatrix} \\ &= L \begin{bmatrix} x \\ y \\ z \end{bmatrix} + \alpha L \begin{bmatrix} a \\ b \\ c \end{bmatrix}, \end{aligned}$$

proving that L is a linear transformation.

540 Let x, y, x', y' be vectors in \mathbb{R}^3 and let $\alpha \in \mathbb{R}$ be a scalar. Then

$$\begin{aligned} L((x, y) + \alpha(x', y')) &= L(x + \alpha x', y + \alpha y') \\ &= (x + \alpha x') \times k + h \times (y + \alpha y') \\ &= x \times k + \alpha x' \times k + h \times y + h \times \alpha y' \\ &= L(x, y) + \alpha L(x', y') \end{aligned}$$

541

$$\begin{aligned} L(H + \alpha H') &= -A^{-1}(H + \alpha H')A^{-1} \\ &= -A^{-1}HA^{-1} + \alpha(-A^{-1}H'A^{-1}) \\ &= L(H) + \alpha L(H'), \end{aligned}$$

proving that L is linear.

542 Let S be convex and let $a, b \in T(S)$. We must prove that $\forall \alpha \in [0; 1], (1 - \alpha)a + \alpha b \in T(S)$. But since a, b belong to $T(S)$, $\exists x \in S, y \in S$ with $T(x) = a, T(y) = b$. Since S is convex, $(1 - \alpha)x + \alpha y \in S$. Thus

$$T((1 - \alpha)x + \alpha y) \in T(S),$$

which means that

$$(1 - \alpha)T(x) + \alpha T(y) \in T(S),$$

that is,

$$(1 - \alpha)a + \alpha b \in T(S),$$

as we wished to show.

550 Assume $\begin{bmatrix} x \\ y \\ z \end{bmatrix} \in \ker(L)$. Then

$$L \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix},$$

that is

$$x - y - z = 0,$$

$$x + y + z = 0,$$

$$z = 0.$$

This implies that $x - y = 0$ and $x + y = 0$, and so $x = y = z = 0$. This means that

$$\ker(L) = \left\{ \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \right\},$$

and L is injective.

By the Dimension Theorem 546, $\dim \text{Im}(L) = \dim V - \dim \ker(L) = 3 - 0 = 3$, which means that

$$\text{Im}(L) = \mathbb{R}^3$$

and L is surjective.

551 Assume that $\begin{bmatrix} a \\ b \\ c \end{bmatrix} \in \ker(T)$,

$$\begin{bmatrix} a \\ b \\ c \end{bmatrix} = (a - b) \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + b \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + c \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}.$$

Then

$$\begin{aligned}
 \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} &= T \begin{bmatrix} a \\ b \\ c \end{bmatrix} \\
 &= (a - b)T \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + bT \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + cT \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \\
 &= (a - b) \begin{bmatrix} 1 \\ 0 \\ -1 \\ 0 \end{bmatrix} + b \begin{bmatrix} 2 \\ -1 \\ 0 \\ 0 \end{bmatrix} + c \begin{bmatrix} 1 \\ -1 \\ 1 \\ 0 \end{bmatrix} \\
 &= \begin{bmatrix} a+b+c \\ -b-c \\ -a+b+c \\ 0 \end{bmatrix}.
 \end{aligned}$$

It follows that $a = 0$ and $b = -c$. Thus

$$\ker(T) = \left\{ c \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix} : c \in \mathbb{R} \right\},$$

and so $\dim \ker(T) = 1$.

By the Dimension Theorem 546,

$$\dim \text{Im}(T) = \dim V - \dim \ker(T) = 3 - 1 = 2.$$

We readily see that

$$\begin{bmatrix} 2 \\ -1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ -1 \\ 0 \end{bmatrix} + \begin{bmatrix} 1 \\ -1 \\ 1 \\ 0 \end{bmatrix},$$

and so

$$\text{Im } (\mathbf{T}) = \text{span} \left(\begin{bmatrix} 1 \\ 0 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \\ 0 \end{bmatrix} \right).$$

552 Assume that

$$\mathbf{L} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x + 2y \\ x + 2y \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}.$$

Then $x = -2y$ and so

$$\begin{bmatrix} x \\ y \end{bmatrix} = y \begin{bmatrix} -2 \\ 1 \end{bmatrix}.$$

This means that $\dim \ker (\mathbf{L}) = 1$ and $\ker (\mathbf{L})$ is the line through the origin and $(-2, 1)$. Observe that \mathbf{L} is not injective.

$$\begin{bmatrix} a \\ b \\ c \end{bmatrix}$$

By the Dimension Theorem 546, $\dim \text{Im } (\mathbf{L}) = \dim V - \dim \ker (\mathbf{L}) = 2 - 1 = 1$. Assume that

$\exists (x, y) \in \mathbb{R}^2$ such that

$$\mathbf{L} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x + 2y \\ x + 2y \\ 0 \end{bmatrix} = \begin{bmatrix} a \\ b \\ c \end{bmatrix}.$$

This means that

$$\begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} x + 2y \\ x + 2y \\ 0 \end{bmatrix} = (x + 2y) \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}.$$

Observe that \mathbf{L} is not surjective.

553 Assume that

$$\mathbf{L} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x - y \\ x + y \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}.$$

Then $x + y = 0 = x - y$, that is, $x = y = 0$, meaning that

$$\ker (\mathbf{L}) = \left\{ \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right\},$$

and so L is injective.

By the Dimension Theorem 546, $\dim \text{Im}(L) = \dim V - \dim \ker(L) = 2 - 0 = 2$. Assume that $\begin{bmatrix} a \\ b \\ c \end{bmatrix} \in \text{Im}(L)$. Then

$\exists(x, y) \in \mathbb{R}^2$ such that

$$L \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x-y \\ x+y \\ 0 \end{bmatrix} = \begin{bmatrix} a \\ b \\ c \end{bmatrix}.$$

This means that

$$\begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} x-y \\ x+y \\ 0 \end{bmatrix} = x \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + y \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}.$$

Since

$$\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$$

are linearly independent, they span a subspace of dimension 2 in \mathbb{R}^3 , that is, a plane containing the origin. Observe that L is not surjective.

554 Assume that

$$L \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} x-y-z \\ y-2z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

Then $y = 2z; x = y + z = 3z$. This means that $\ker(L) = \left\{ z \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix} : z \in \mathbb{R} \right\}$. Hence $\dim \ker(L) = 1$, and so L is not injective.

Now, if

$$L \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} x-y-z \\ y-2z \end{bmatrix} = \begin{bmatrix} a \\ b \end{bmatrix}.$$

Then

$$\begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} x-y-z \\ y-2z \end{bmatrix} = x \begin{bmatrix} 1 \\ 0 \end{bmatrix} + y \begin{bmatrix} -1 \\ 1 \end{bmatrix} + z \begin{bmatrix} -1 \\ -2 \end{bmatrix}.$$

Now,

$$-3 \begin{bmatrix} 1 \\ 0 \end{bmatrix} - 2 \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} -1 \\ -2 \end{bmatrix}$$

and

$$\begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

are linearly independent. Since $\dim \text{Im } (\mathbf{L}) = 2$, we have $\text{Im } (\mathbf{L}) = \mathbb{R}^2$, and so \mathbf{L} is surjective.

555 Assume that

$$0 = \text{tr} \left(\begin{bmatrix} a & b \\ c & d \end{bmatrix} \right) = a + d.$$

Then $a = -d$ and so,

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} -d & b \\ c & d \end{bmatrix} = d \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} + b \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} + c \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix},$$

and so $\dim \ker (\mathbf{L}) = 3$. Thus \mathbf{L} is not injective. \mathbf{L} is surjective, however. For if $\alpha \in \mathbb{R}$, then

$$\alpha = \text{tr} \left(\begin{bmatrix} \alpha & 0 \\ 0 & 0 \end{bmatrix} \right).$$

556 1. Let $(\mathbf{A}, \mathbf{B})^T \in \mathbf{M}_2(\mathbb{R})$, $\alpha \in \mathbb{R}$. Then

$$\begin{aligned} \mathbf{L}(\mathbf{A} + \alpha \mathbf{B}) &= (\mathbf{A} + \alpha \mathbf{B})^T + (\mathbf{A} + \alpha \mathbf{B}) \\ &= \mathbf{A}^T + \mathbf{B}^T + \mathbf{A} + \alpha \mathbf{B} \\ &= \mathbf{A}^T + \mathbf{A} + \alpha \mathbf{B}^T + \alpha \mathbf{B} \\ &= \mathbf{L}(\mathbf{A}) + \alpha \mathbf{L}(\mathbf{B}), \end{aligned}$$

proving that \mathbf{L} is linear.

2. Assume that $\mathbf{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \ker (\mathbf{L})$. Then

$$\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = \mathbf{L}(\mathbf{A}) = \begin{bmatrix} a & b \\ c & d \end{bmatrix} + \begin{bmatrix} a & c \\ b & d \end{bmatrix} = \begin{bmatrix} 2a & b+c \\ b+c & 2d \end{bmatrix},$$

whence $a = d = 0$ and $b = -c$. Hence

$$\ker (\mathbf{L}) = \text{span} \left(\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \right),$$

and so $\dim \ker (\mathbf{L}) = 1$.

3. By the Dimension Theorem, $\dim \text{Im}(\mathbf{L}) = 4 - 1 = 3$. As above,

$$\begin{aligned}\mathbf{L}(\mathbf{A}) &= \begin{bmatrix} 2\mathbf{a} & \mathbf{b} + \mathbf{c} \\ \mathbf{b} + \mathbf{c} & 2\mathbf{d} \end{bmatrix} \\ &= \mathbf{a} \begin{bmatrix} 2 & 0 \\ 0 & 0 \end{bmatrix} + (\mathbf{b} + \mathbf{c}) \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} + \mathbf{d} \begin{bmatrix} 0 & 0 \\ 0 & 2 \end{bmatrix},\end{aligned}$$

from where

$$\text{Im}(\mathbf{L}) = \text{span} \left(\begin{bmatrix} 2 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 2 \end{bmatrix} \right).$$

557 ① Observe that

$$(\mathbf{I} - \mathbf{T})^2 = \mathbf{I} - 2\mathbf{T} + \mathbf{T}^2 = \mathbf{I} - 2\mathbf{T} + \mathbf{T} = \mathbf{I} - \mathbf{T},$$

proving the result.

② The inverse is $\mathbf{I} - \frac{1}{2}\mathbf{T}$, for

$$(\mathbf{I} + \mathbf{T})(\mathbf{I} - \frac{1}{2}\mathbf{T}) = \mathbf{I} + \mathbf{T} - \frac{1}{2}\mathbf{T} - \frac{1}{2}\mathbf{T}^2 = \mathbf{I} + \mathbf{T} - \frac{1}{2}\mathbf{T} - \frac{1}{2}\mathbf{T} = \mathbf{I},$$

proving the claim.

③ We have

$$\begin{aligned}\mathbf{x} \in \ker(\mathbf{T}) &\iff \mathbf{x} - \mathbf{T}(\mathbf{x}) \in \ker(\mathbf{T}) \\ &\iff \mathbf{I}(\mathbf{x}) - \mathbf{T}(\mathbf{x}) \in \ker(\mathbf{T}) \\ &\iff (\mathbf{I} - \mathbf{T})(\mathbf{x}) \in \ker(\mathbf{T}) \\ &\iff \mathbf{x} \in \text{Im}(\mathbf{I} - \mathbf{T}).\end{aligned}$$

562 1. Since the image of \mathbf{T} is the plane $x + y + z = 0$, we must have

$$\mathbf{a} + \mathbf{0} + \mathbf{1} = \mathbf{0} \implies \mathbf{a} = -\mathbf{1},$$

$$3 + \mathbf{b} - 5 = \mathbf{0} \implies \mathbf{b} = 2,$$

$$-\mathbf{1} + 2 + \mathbf{c} = \mathbf{0} \implies \mathbf{c} = -\mathbf{1}.$$

2. Observe that $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \in \ker(\mathbf{T})$ and so

$$\mathbf{T} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}.$$

Thus

$$\mathbf{T} \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = \mathbf{T} \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} - \mathbf{T} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 \\ 2 \\ -5 \end{bmatrix},$$

$$\mathbf{T} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \mathbf{T} \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} - \mathbf{T} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \\ -1 \end{bmatrix},$$

$$\mathbf{T} \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \mathbf{T} \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix} - \mathbf{T} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}.$$

The required matrix is therefore

$$\begin{bmatrix} 3 & -1 & -1 \\ 2 & 2 & 0 \\ -5 & -1 & 1 \end{bmatrix}.$$

- 563** 1. Let $\alpha \in \mathbb{R}$. We have

$$\begin{aligned} \mathbf{T} \left(\begin{bmatrix} x \\ y \end{bmatrix} + \alpha \begin{bmatrix} u \\ v \end{bmatrix} \right) &= \mathbf{T} \left(\begin{bmatrix} x + \alpha u \\ y + \alpha v \end{bmatrix} \right) \\ &= \begin{bmatrix} x + \alpha u + y + \alpha v \\ x + \alpha u - y - \alpha v \\ 2(x + \alpha u) + 3(y + \alpha v) \end{bmatrix} \\ &= \begin{bmatrix} x + y \\ x - y \\ 2x + 3y \end{bmatrix} + \alpha \begin{bmatrix} u + v \\ u - v \\ 2u + 3v \end{bmatrix} \\ &= \mathbf{T} \left(\begin{bmatrix} x \\ y \end{bmatrix} \right) + \alpha \mathbf{T} \left(\begin{bmatrix} u \\ v \end{bmatrix} \right), \end{aligned}$$

proving that \mathbf{T} is linear.

2. We have

$$\mathbf{T} \left(\begin{bmatrix} x \\ y \end{bmatrix} \right) = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \iff \begin{bmatrix} x + y \\ x - y \\ 2x + 3y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \iff x = y = 0,$$

$\dim \ker(\mathbf{T}) = 0$, and whence \mathbf{T} is injective.

3. By the Dimension Theorem, $\dim \text{Im}(\mathbf{T}) = 2 - 0 = 2$. Now, since

$$\mathbf{T} \left(\begin{bmatrix} x \\ y \end{bmatrix} \right) = \begin{bmatrix} x + y \\ x - y \\ 2x + 3y \end{bmatrix} = x \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix} + y \begin{bmatrix} 1 \\ -1 \\ 3 \end{bmatrix},$$

whence

$$\text{Im } (\mathbf{T}) = \text{span} \left(\begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 3 \end{bmatrix} \right).$$

4. We have

$$\mathbf{T} \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix} = \begin{bmatrix} 3 \\ -1 \\ 8 \end{bmatrix} = \frac{11}{2} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} - \frac{5}{2} \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} - \frac{13}{2} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 11/2 \\ -5/2 \\ -13/2 \end{bmatrix}_{\mathcal{B}},$$

and

$$\mathbf{T} \begin{pmatrix} 1 \\ 3 \\ 11 \end{pmatrix} = \begin{bmatrix} 4 \\ -2 \\ 11 \end{bmatrix} = \frac{15}{2} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} - \frac{7}{2} \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} - \frac{19}{2} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 15/2 \\ -7/2 \\ -19/2 \end{bmatrix}_{\mathcal{B}}.$$

The required matrix is

$$\begin{bmatrix} 11/2 & 15/2 \\ -5/2 & -7/2 \\ -13/2 & -19/2 \end{bmatrix}_{\mathcal{B}}.$$

564 The matrix will be a 2×3 matrix. In each case, we find the action of \mathbf{L} on the basis elements of \mathbb{R}^3 and express the result in the given basis for \mathbb{R}^3 .

1. We have

$$\mathbf{L} \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = \begin{bmatrix} 1 \\ 1 \\ 3 \end{bmatrix}, \mathbf{L} \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} = \begin{bmatrix} 2 \\ 0 \\ 0 \end{bmatrix}, \mathbf{L} \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} = \begin{bmatrix} 0 \\ 0 \\ -1 \end{bmatrix}.$$

The required matrix is

$$\begin{bmatrix} 1 & 2 & 0 \\ 3 & 0 & -1 \end{bmatrix}.$$

2. We have

$$\mathbf{L} \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = \begin{bmatrix} 1 \\ 1 \\ 3 \end{bmatrix}, \mathbf{L} \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} = \begin{bmatrix} 3 \\ 3 \\ 0 \end{bmatrix}, \mathbf{L} \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}.$$

The required matrix is

$$\begin{bmatrix} 1 & 3 & 3 \\ 3 & 3 & 2 \end{bmatrix}.$$

3. We have

$$\mathbf{L} \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 3 \\ 3 \end{pmatrix} = -2 \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} + 3 \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} -2 \\ 3 \\ 3 \end{pmatrix}_{\mathcal{A}},$$

$$\mathbf{L} \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 3 \\ 3 \\ 3 \end{pmatrix} = 0 \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} + 3 \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 3 \\ 3 \end{pmatrix}_{\mathcal{A}},$$

$$\mathbf{L} \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \\ 2 \end{pmatrix} = 1 \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} + 2 \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \\ 2 \end{pmatrix}_{\mathcal{A}}.$$

The required matrix is

$$\begin{bmatrix} -2 & 0 & 1 \\ 3 & 3 & 2 \end{bmatrix}.$$

565 Observe that $\begin{bmatrix} 2 \\ 3 \end{bmatrix} \in \text{Im } (\mathbf{T}) = \ker (\mathbf{T})$ and so

$$\mathbf{T} \begin{bmatrix} 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

Now

$$\mathbf{T} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \mathbf{T} \left(3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} - \begin{bmatrix} 2 \\ 3 \end{bmatrix} \right) = 3\mathbf{T} \begin{bmatrix} 1 \\ 1 \end{bmatrix} - \mathbf{T} \begin{bmatrix} 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 6 \\ 9 \end{bmatrix},$$

and

$$\mathbf{T} \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \mathbf{T} \left(\begin{bmatrix} 2 \\ 3 \end{bmatrix} - 2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right) = \mathbf{T} \begin{bmatrix} 2 \\ 3 \end{bmatrix} - 2\mathbf{T} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} -4 \\ -6 \end{bmatrix}.$$

The required matrix is thus

$$\begin{bmatrix} 6 & -4 \\ 9 & -6 \end{bmatrix}.$$

566 The matrix will be a 1×4 matrix. We have

$$\text{tr} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} = 1,$$

$$\text{tr} \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} = 0,$$

$$\text{tr} \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} = 0,$$

$$\text{tr} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} = 1.$$

Thus

$$\mathbf{M}_L = (1 \ 0 \ 0 \ 1).$$

567 First observe that $\ker(\mathbf{B}) \subseteq \ker(\mathbf{AB})$ since $\forall \mathbf{X} \in \mathbf{M}_{q \times 1}(\mathbb{R})$,

$$\mathbf{BX} = \mathbf{0} \implies (\mathbf{AB})\mathbf{X} = \mathbf{A}(\mathbf{BX}) = \mathbf{0}.$$

Now

$$\begin{aligned} \dim \ker(\mathbf{B}) &= q - \dim \text{Im}(\mathbf{B}) \\ &= q - \text{rank}(\mathbf{B}) \\ &= q - \text{rank}(\mathbf{AB}) \\ &= q - \dim \text{Im}(\mathbf{AB}) \\ &= \dim \ker(\mathbf{AB}). \end{aligned}$$

Thus $\ker(\mathbf{B}) = \ker(\mathbf{AB})$. Similarly, we can demonstrate that $\ker(\mathbf{ABC}) = \ker(\mathbf{BC})$. Thus

$$\begin{aligned} \text{rank}(\mathbf{ABC}) &= \dim \text{Im}(\mathbf{ABC}) \\ &= r - \dim \ker(\mathbf{ABC}) \\ &= r - \dim \ker(\mathbf{BC}) \\ &= \dim \text{Im}(\mathbf{BC}) \\ &= \text{rank}(\mathbf{BC}). \end{aligned}$$

594 This is clearly $(1 \ 2 \ 3 \ 4)(6 \ 8 \ 7)$ of order $\text{lcm}(4, 3) = 12$.

621 Multiplying the first column of the given matrix by a , its second column by b , and its third column by c , we obtain

$$abc\Omega = \begin{bmatrix} abc & abc & abc \\ a^2 & b^2 & c^2 \\ a^3 & b^3 & c^3 \end{bmatrix}.$$

We may factor out abc from the first row of this last matrix thereby obtaining

$$abc\Omega = abc \det \begin{bmatrix} 1 & 1 & 1 \\ a^2 & b^2 & c^2 \\ a^3 & b^3 & c^3 \end{bmatrix}.$$

Upon dividing by abc ,

$$\Omega = \det \begin{bmatrix} 1 & 1 & 1 \\ a^2 & b^2 & c^2 \\ a^3 & b^3 & c^3 \end{bmatrix}.$$

622 Performing $R_1 + R_2 + R_3 \rightarrow R_1$ we have

$$\begin{aligned}\Omega &= \det \begin{bmatrix} a - b - c & 2a & 2a \\ 2b & b - c - a & 2b \\ 2c & 2c & c - a - b \end{bmatrix} \\ &= \det \begin{bmatrix} a + b + c & a + b + c & a + b + c \\ 2b & b - c - a & 2b \\ 2c & 2c & c - a - b \end{bmatrix}.\end{aligned}$$

Factorising $(a + b + c)$ from the first row of this last determinant, we have

$$\Omega = (a + b + c) \det \begin{bmatrix} 1 & 1 & 1 \\ 2b & b - c - a & 2b \\ 2c & 2c & c - a - b \end{bmatrix}.$$

Performing $C_2 - C_1 \rightarrow C_2$ and $C_3 - C_1 \rightarrow C_3$,

$$\Omega = (a + b + c) \det \begin{bmatrix} 1 & 0 & 0 \\ 2b & -b - c - a & 0 \\ 2c & 0 & -c - a - b \end{bmatrix}.$$

This last matrix is triangular, hence

$$\Omega = (a + b + c)(-b - c - a)(-c - a - b) = (a + b + c)^3,$$

as wanted.

623 $\det A_1 = \det A = -540$ by multilinearity. $\det A_2 = -\det A_1 = 540$ by alternancy. $\det A_3 = 3 \det A_2 = 1620$ by both multilinearity and homogeneity from one column. $\det A_4 = \det A_3 = 1620$ by multilinearity, and $\det A_5 = 2 \det A_4 = 3240$ by homogeneity from one column.

624 From the given data, $\det B = -2$. Hence

$$\det ABC = (\det A)(\det B)(\det C) = -12,$$

$$\det 5AC = 5^3 \det AC = (125)(\det A)(\det C) = 750,$$

$$(\det A^3 B^{-3} C^{-1}) = \frac{(\det A)^3}{(\det B)^3 (\det C)} = -\frac{27}{16}.$$

625 Pick $\lambda \in \mathbb{R} \setminus \{0, a_{11}, a_{22}, \dots, a_{nn}\}$. Put

$$X = \begin{bmatrix} a_{11} - \lambda & 0 & 0 & \cdots & 0 \\ a_{21} & a_{22} - \lambda & 0 & \cdots & 0 \\ a_{31} & a_{32} & a_{33} - \lambda & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ a_{n1} & a_{n2} & a_{n3} & \cdots & a_{nn} - \lambda \end{bmatrix}$$

and

$$Y = \begin{bmatrix} \lambda & a_{12} & a_{13} & \cdots & a_{1n} \\ 0 & \lambda & a_{23} & \cdots & a_{2n} \\ 0 & 0 & \lambda & \cdots & a_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \lambda \end{bmatrix}$$

Clearly $A = X + Y$, $\det X = (a_{11} - \lambda)(a_{22} - \lambda) \cdots (a_{nn} - \lambda) \neq 0$, and $\det Y = \lambda^n \neq 0$. This completes the proof.

626 No.

636 We have

$$\begin{aligned} \det A &= 2(-1)^{1+2} \det \begin{bmatrix} 4 & 6 \\ 7 & 9 \end{bmatrix} + 5(-1)^{2+2} \det \begin{bmatrix} 1 & 3 \\ 7 & 9 \end{bmatrix} + 8(-1)^{2+3} \det \begin{bmatrix} 1 & 3 \\ 4 & 6 \end{bmatrix} \\ &= -2(36 - 42) + 5(9 - 21) - 8(6 - 12) = 0. \end{aligned}$$

637 Since the second column has three 0's, it is advantageous to expand along it, and thus we are reduced to calculate

$$-3(-1)^{3+2} \det \begin{bmatrix} 1 & -1 & 1 \\ 2 & 0 & 1 \\ 1 & 0 & 1 \end{bmatrix}$$

Expanding this last determinant along the second column, the original determinant is thus

$$-3(-1)^{3+2}(-1)(-1)^{1+2} \det \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix} = -3(-1)(-1)(-1)(1) = 3.$$

638 Expanding along the first column,

$$\begin{aligned}
 0 &= \det \begin{bmatrix} 1 & 1 & 1 & 1 \\ x & a & 0 & 0 \\ x & 0 & b & 0 \\ x & 0 & 0 & c \end{bmatrix} \\
 &= \det \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix} - x \det \begin{bmatrix} 1 & 1 & 1 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix} \\
 &\quad + x \det \begin{bmatrix} 1 & 1 & 1 \\ a & 0 & 0 \\ 0 & 0 & c \end{bmatrix} - x \det \begin{bmatrix} 1 & 1 & 1 \\ a & 0 & 0 \\ 0 & b & 0 \end{bmatrix} \\
 &= xabc - xbc + x \det \begin{bmatrix} 1 & 1 & 1 \\ a & 0 & 0 \\ 0 & 0 & c \end{bmatrix} - x \det \begin{bmatrix} 1 & 1 & 1 \\ a & 0 & 0 \\ 0 & b & 0 \end{bmatrix}.
 \end{aligned}$$

Expanding these last two determinants along the third row,

$$\begin{aligned}
 0 &= abc - xbc + x \det \begin{bmatrix} 1 & 1 & 1 \\ a & 0 & 0 \\ 0 & 0 & c \end{bmatrix} - x \det \begin{bmatrix} 1 & 1 & 1 \\ a & a & 0 \\ 0 & b & 0 \end{bmatrix} \\
 &= abc - xbc + xc \det \begin{bmatrix} 1 & 1 \\ a & 0 \end{bmatrix} + xb \det \begin{bmatrix} 1 & 1 \\ a & 0 \end{bmatrix} \\
 &= abc - xbc - xca - xab.
 \end{aligned}$$

It follows that

$$abc = x(bc + ab + ca),$$

whence

$$\frac{1}{x} = \frac{bc + ab + ca}{abc} = \frac{1}{a} + \frac{1}{b} + \frac{1}{c},$$

as wanted.

639 Expanding along the first row the determinant equals

$$\begin{aligned}
 -a \det \begin{bmatrix} a & b & 0 \\ 0 & 0 & b \\ 1 & 1 & 1 \end{bmatrix} + b \det \begin{bmatrix} a & 0 & 0 \\ 0 & a & b \\ 1 & 1 & 1 \end{bmatrix} &= ab \det \begin{bmatrix} a & b \\ 1 & 1 \end{bmatrix} + ab \det \begin{bmatrix} a & b \\ 1 & 1 \end{bmatrix} \\
 &= 2ab(a - b),
 \end{aligned}$$

as wanted.

640 Expanding along the first row, the determinant equals

$$a \det \begin{bmatrix} a & 0 & b \\ 0 & d & 0 \\ c & 0 & d \end{bmatrix} + b \det \begin{bmatrix} 0 & a & b \\ c & 0 & 0 \\ 0 & c & d \end{bmatrix}.$$

Expanding the resulting two determinants along the second row, we obtain

$$ad \det \begin{bmatrix} a & b \\ c & d \end{bmatrix} + b(-c) \det \begin{bmatrix} a & b \\ c & d \end{bmatrix} = ad(ad - bc) - bc(ad - bc) = (ad - bc)^2,$$

as wanted.

641 For $n = 1$ we have $\det(1) = 1 = (-1)^{1+1}$. For $n = 2$ we have

$$\det \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} = -1 = (-1)^{2+1}.$$

Assume that the result is true for $n - 1$. Expanding the determinant along the first column

$$\begin{aligned} \det \begin{bmatrix} 1 & 1 & 1 & \cdots & 1 & 1 \\ 1 & 0 & 0 & \vdots & 0 & 0 \\ 0 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \cdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & 0 \end{bmatrix} &= 1 \det \begin{bmatrix} 0 & 0 & \cdots & 0 & 0 \\ 1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ \vdots & \cdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & 0 \end{bmatrix} \\ &\quad -1 \det \begin{bmatrix} 1 & 1 & \cdots & 1 & 1 \\ 1 & 0 & \cdots & \vdots & 0 \\ 0 & 1 & \cdots & \cdots & 0 \\ 0 & 0 & \cdots & \cdots & 0 \\ \vdots & \vdots & \cdots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & 0 \end{bmatrix} \\ &= 1(0) - (1)(-1)^n \\ &= (-1)^{n+1}, \end{aligned}$$

giving the result.

642 Perform $C_k - C_1 \rightarrow C_k$ for $k \in [2; n]$. Observe that these operations do not affect the value of the determinant. Then

$$\det A = \det \begin{bmatrix} 1 & n-1 & n-1 & n-1 & \cdots & n-1 \\ n & 2-n & 0 & 0 & \vdots & 0 \\ n & 0 & 3-n & 0 & \cdots & 0 \\ n & 0 & 0 & 4-n & \cdots & 0 \\ \vdots & \vdots & \vdots & \cdots & \vdots & \vdots \\ n & 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

Expand this last determinant along the n -th row, obtaining,

$$\begin{aligned} \det A &= (-1)^{1+n} n \det \begin{bmatrix} n-1 & n-1 & n-1 & \cdots & n-1 & n-1 \\ 2-n & 0 & 0 & \vdots & 0 & 0 \\ 0 & 3-n & 0 & \cdots & 0 & 0 \\ 0 & 0 & 4-n & \cdots & 0 & 0 \\ \vdots & \vdots & \cdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & -1 & 0 \end{bmatrix} \\ &= (-1)^{1+n} n(n-1)(2-n)(3-n) \\ &\quad \cdots (-2)(-1) \det \begin{bmatrix} 1 & 1 & 1 & \cdots & 1 & 1 \\ 1 & 0 & 0 & \vdots & 0 & 0 \\ 0 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \cdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & 0 \end{bmatrix} \\ &= -(n!) \det \begin{bmatrix} 1 & 1 & 1 & \cdots & 1 & 1 \\ 1 & 0 & 0 & \vdots & 0 & 0 \\ 0 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \cdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & 0 \end{bmatrix} \\ &= -(n!)(-1)^n \\ &= (-1)^{n+1} n!, \end{aligned}$$

upon using the result of problem 641.

643 Recall that $\binom{n}{k} = \binom{n}{n-k}$,

$$\sum_{k=0}^n \binom{n}{k} = 2^n$$

and

$$\sum_{k=0}^n (-1)^k \binom{n}{k} = 0, \quad \text{if } n > 0.$$

Assume that n is odd. Observe that then there are $n+1$ (an even number) of columns and that on the same row, $\binom{n}{k}$ is on a column of opposite parity to that of $\binom{n}{n-k}$. By performing $C_1 - C_2 + C_3 - C_4 + \dots + C_n - C_{n+1} \rightarrow C_1$, the first column becomes all 0's, whence the determinant is 0 if n is odd.

659 We have

$$\det(\lambda I_2 - A) = \det \begin{bmatrix} \lambda - 1 & 1 \\ 1 & \lambda - 1 \end{bmatrix} = (\lambda - 1)^2 - 1 = \lambda(\lambda - 2),$$

whence the eigenvalues are 0 and 2. For $\lambda = 0$ we have

$$0I_2 - A = \begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix}.$$

This has row-echelon form

$$\begin{bmatrix} 1 & -1 \\ 0 & 0 \end{bmatrix}.$$

If

$$\begin{bmatrix} 1 & -1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

then $a = b$. Thus

$$\begin{bmatrix} a \\ b \end{bmatrix} = a \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

and we can take $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$ as the eigenvector corresponding to $\lambda = 0$. Similarly, for $\lambda = 2$,

$$2I_2 - A = \begin{bmatrix} 1 & 3 \\ 1 & 3 \end{bmatrix},$$

which has row-echelon form

$$\begin{bmatrix} 1 & 3 \\ 0 & 0 \end{bmatrix}.$$

If

$$\begin{bmatrix} 1 & 3 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

then $\mathbf{a} = -3\mathbf{b}$. Thus

$$\begin{bmatrix} \mathbf{a} \\ \mathbf{b} \end{bmatrix} = \mathbf{a} \begin{bmatrix} 1 \\ -3 \end{bmatrix}$$

and we can take $\begin{bmatrix} 1 \\ -3 \end{bmatrix}$ as the eigenvector corresponding to $\lambda = 2$.

660 ❶ We have

$$\begin{aligned} \det(\lambda \mathbf{I}_3 - \mathbf{A}) &= \det \begin{bmatrix} \lambda & -2 & 1 \\ -2 & \lambda - 3 & 2 \\ 1 & 2 & \lambda \end{bmatrix} \\ &= \lambda \det \begin{bmatrix} \lambda - 3 & 2 \\ 2 & \lambda \end{bmatrix} + 2 \det \begin{bmatrix} -2 & 2 \\ 1 & \lambda \end{bmatrix} + \det \begin{bmatrix} -2 & \lambda - 3 \\ 1 & 2 \end{bmatrix} \\ &= \lambda(\lambda^2 - 3\lambda - 4) + 2(-2\lambda - 2) + (-\lambda - 1) \\ &= \lambda(\lambda - 4)(\lambda + 1) - 5(\lambda + 1) \\ &= (\lambda^2 - 4\lambda - 5)(\lambda + 1) \\ &= (\lambda + 1)^2(\lambda - 5) \end{aligned}$$

❷ The eigenvalues are $-1, -1, 5$.

❸ If $\lambda = -1$,

$$\begin{aligned} (-\mathbf{I}_3 - \mathbf{A}) &= \begin{bmatrix} -1 & -2 & 1 \\ -2 & -4 & 2 \\ 1 & 2 & -1 \end{bmatrix} \begin{bmatrix} \mathbf{a} \\ \mathbf{b} \\ \mathbf{c} \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \iff \mathbf{a} = -2\mathbf{b} + \mathbf{c} \\ &\iff \begin{bmatrix} \mathbf{a} \\ \mathbf{b} \\ \mathbf{c} \end{bmatrix} = \mathbf{b} \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} + \mathbf{c} \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}. \end{aligned}$$

We may take as eigenvectors $\begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$, which are clearly linearly independent.

If $\lambda = 5$,

$$(5\mathbf{I}_3 - \mathbf{A}) = \begin{bmatrix} 5 & -2 & 1 \\ -2 & 2 & 2 \\ 1 & 2 & 5 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \iff a = -c, b = -2c,$$

$$\iff \begin{bmatrix} a \\ b \\ c \end{bmatrix} = c \begin{bmatrix} -1 \\ -2 \\ 1 \end{bmatrix}.$$

We may take as eigenvector $\begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}$.

664 Solution: Put

$$\mathbf{D} = \begin{bmatrix} 1 & 0 \\ 0 & -2 \end{bmatrix}, \quad , \quad \mathbf{P} = \begin{bmatrix} 1 & 0 \\ 1 & -1 \end{bmatrix}.$$

We find

$$\mathbf{P}^{-1} = \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix}.$$

Since $\mathbf{A} = \mathbf{PDP}^{-1}$

$$\mathbf{A}^{10} = \mathbf{P}\mathbf{D}^{10}\mathbf{P}^{-1} = \begin{bmatrix} 1 & 0 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1024 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix} = \begin{bmatrix} 1 & -1023 \\ 0 & 1024 \end{bmatrix}.$$

665 Put

$$\mathbf{D} = \begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 3 \end{bmatrix}, \quad \mathbf{X} = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}.$$

Then we know that $\mathbf{A} = \mathbf{XDX}^{-1}$ and so we need to find \mathbf{X}^{-1} . But this is readily obtained by performing $R_1 - R_2 \rightarrow R_1$ and $R_2 - R_3 \rightarrow R_3$ in the augmented matrix

$$\left[\begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right],$$

getting

$$\mathbf{X}^{-1} = \begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{bmatrix}.$$

Thus

$$\begin{aligned}\mathbf{A} &= \mathbf{XDX}^{-1} \\ &= \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 3 \end{bmatrix} \begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{bmatrix} \\ &= \begin{bmatrix} -1 & 0 & 4 \\ 0 & -1 & 4 \\ 0 & 0 & 3 \end{bmatrix}.\end{aligned}$$

666 The determinant is 1, $\mathbf{A} = \mathbf{A}^{-1}$, and the characteristic polynomial is $(\lambda^2 - 1)^2$.

Bibliography

- [Cul] CULLEN, C., **Matrices and Linear Transformations**, 2nd ed., New York: Dover Publications, 1990.
- [Del] DELODE, C., **Géométrie Affine et Euclidienne**, Paris: Dunod, 2000.
- [Fad] FADDÉEV, D., SOMINSKY, I., **Recueil d'Exercices d'Algèbre Supérieure**, 7th ed., Paris: Ellipses, 1994.
- [Hau] HAUSNER, M., **A Vector Space Approach to Geometry**, New York: Dover, 1998.
- [Lan] LANG, S., **Introduction to Linear Algebra**, 2nd ed., New York: Springer-Verlag, 1986.
- [Pro] PROSKURYAKOV, I. V., **Problems in Linear Algebra**, Moscow: Mir Publishers, 1978.
- [Riv] RIVAUD, J., **Ejercicios de álgebra**, Madrid: Aguilar, 1968.
- [Tru] TRUFFAULT, B., **Géométrie Élémentaire: Cours et exercices**, Paris: Ellipses, 2001.