



# The effect of asymmetric Ekman damping on energy and enstrophy injection in two-layer quasi-geostrophic turbulence

Eleftherios Gkioulekas <sup>(1)</sup> and Ka-Kit Tung <sup>(2)</sup>

<sup>(1)</sup> Department of Mathematics, University of Texas-Pan American

<sup>(2)</sup> Department of Applied Mathematics, University of Washington

# Overview

- Review of 2D turbulence.
- The Nastrom-Gage spectrum of atmospheric turbulence.
- The Tung-Orlando theory of double cascade.
- The two-layer quasi-geostrophic model.

# 2D Navier-Stokes equations

- In 2D turbulence, the scalar vorticity  $\zeta(x, y, t)$  is governed by

$$\frac{\partial \zeta}{\partial t} + J(\psi, \zeta) = d + f, \quad (1)$$

where  $\psi(x, y, t)$  is the streamfunction, and  $\zeta(x, y, t) = -\nabla^2 \psi(x, y, t)$ , and

$$d = -[\nu(-\Delta)^\kappa + \nu_1(-\Delta)^{-m}] \zeta \quad (2)$$

- The Jacobian term  $J(\psi, \zeta)$  describes the advection of  $\zeta$  by  $\psi$ , and is defined as

$$J(a, b) = \frac{\partial a}{\partial x} \frac{\partial b}{\partial y} - \frac{\partial b}{\partial x} \frac{\partial a}{\partial y}. \quad (3)$$

# Energy and enstrophy spectrum. I

- ➊ Two conserved quadratic invariants: energy  $E$  and enstrophy  $G$  defined as

$$E(t) = -\frac{1}{2} \int \psi(x, y, t) \zeta(x, y, t) \, dx dy \quad G(t) = \frac{1}{2} \int \zeta^2(x, y, t) \, dx dy. \quad (4)$$

- ➋ Let  $a^{<k}(\mathbf{x})$  be the field obtained from  $a(\mathbf{x})$  by setting to zero, in Fourier space, the components corresponding to wavenumbers with norm greater than  $k$ :

$$a^{<k}(\mathbf{x}) = \int d\mathbf{y} P(k|\mathbf{x} - \mathbf{y}) a(\mathbf{y}) \quad (5)$$

$$= \int_{\mathbb{R}^2} d\mathbf{x}_0 \int_{\mathbb{R}^2} d\mathbf{k}_0 \frac{H(k - \|\mathbf{k}_0\|)}{4\pi^2} \exp(i\mathbf{k}_0 \cdot (\mathbf{x} - \mathbf{x}_0)) a(\mathbf{x}_0) \quad (6)$$

- ➌ Filtered inner product:

$$\langle a, b \rangle_k = \frac{d}{dk} \int_{\mathbb{R}^2} d\mathbf{x} a^{<k}(\mathbf{x}) b^{<k}(\mathbf{x}) \quad (7)$$

# Energy and enstrophy spectrum. II

- Energy spectrum:  $E(k) = \langle \psi, \zeta \rangle_k$
- Enstrophy spectrum  $G(k) = \langle \zeta, \zeta \rangle_k$
- Consider the conservation laws for  $E(k)$  and  $G(k)$ :

$$\frac{\partial E(k)}{\partial t} + \frac{\partial \Pi_E(k)}{\partial k} = D_E(k) + F_E(k) \quad (8)$$


$$\frac{\partial G(k)}{\partial t} + \frac{\partial \Pi_E(k)}{\partial k} = D_G(k) + F_G(k) \quad (9)$$

- In two-dimensional turbulence, the energy flux  $\Pi_E(k)$  and the enstrophy flux  $\Pi_G(k)$  are constrained by

$$k^2 \Pi_E(k) - \Pi_G(k) < 0 \quad (10)$$

for all  $k$  not in the forcing range.

# KLB theory.



$k_f$  = forcing wavenumber

$k_{ir}$  = IR dissipation wavenumber

$k_{uv}$  = UV dissipation wavenumber

$\varepsilon_{ir}$  = upscale energy flux

$\eta_{uv}$  = downscale enstrophy flux

# The need for a large-scale sink

- ➊ It can also be shown that:  $F_E(k) = k^{-2}F_G(k)$ . It follows that
    - ➋ At the forcing range:  $(\eta/\varepsilon) \sim k_f^2$
    - ➋ At the dissipation range:  $(\eta_{uv}/\varepsilon_{uv}) \sim k_d^2 \gg k_f^2$
  - ➋ It follows that most of the energy injected at the forcing range along with the enstrophy cannot be dissipated at small scales.
  - ➋ A large scale sink that can reduce predominantly  $\varepsilon_{uv}$  is needed.
  - ➋ In the papers:
    - ➋ E. Gkioulekas and K.K. Tung (2005), *DCDS B*, **5**, 79-102
    - ➋ E. Gkioulekas and K.K. Tung (2005), *DCDS B*, **5**, 103-124.
- we have theorized that if it were possible to have  $(\eta_{uv}/\varepsilon_{uv}) \gg k_d^2$ , then within the enstrophy cascade we would see a transition from  $k^{-3}$  scaling to  $k^{-5/3}$  scaling at the transition wavenumber  $k_t \sim (\eta_{uv}/\varepsilon_{uv})$ .

# Superposition principle. I

Specifically, Gkioulekas and Tung (2005) have argued that in 2D turbulence a leading downscale enstrophy cascade *and a subleading downscale energy cascade* contribute linearly to the total energy spectrum:

$$E(k) = E_{uv}^{(\varepsilon)}(k) + E_{uv}^{(\eta)}(k) + E_{uv}^{(p)}(k), \quad \forall k\ell_0 \gg 1, \quad (11)$$

where  $E_{uv}^{(\varepsilon)}(k)$ ,  $E_{uv}^{(\eta)}(k)$  are the contributions of the downscale energy and enstrophy cascade, given by

$$\begin{aligned} E_{uv}^{(\varepsilon)}(k) &= a_{uv}\varepsilon_{uv}^{2/3}k^{-5/3}\mathcal{D}_{uv}^{(\varepsilon)}(k\ell_{uv}^{(\varepsilon)}) \\ E_{uv}^{(\eta)}(k) &= b_{uv}\eta_{uv}^{2/3}k^{-3}[\chi + \ln(k\ell_0)]^{-1/3}\mathcal{D}_{uv}^{(\eta)}(k\ell_{uv}^{(\eta)}), \end{aligned} \quad (12)$$

Thus, in the inertial range where the effect of forcing and dissipation can be ignored, the energy spectrum will take the simple form

$$E(k) \approx a_{uv}\varepsilon_{uv}^{2/3}k^{-5/3} + b_{uv}\eta_{uv}^{2/3}k^{-3}[\chi + \ln(k\ell_0)]^{-1/3}. \quad (13)$$

# Superposition principle. II

- ➊ To recap: the downscale enstrophy cascade contributes a dominant  $k^{-3}$  term to  $E(k)$  and the downscale energy cascade a subdominant  $k^{-5/3}$  term.
- ➋ Why is the  $k^{-5/3}$  term hidden in two-dimensional turbulence?
- ➌ In two-dimensional turbulence, the energy flux  $\Pi_E(k)$  and the enstrophy flux  $\Pi_G(k)$  are constrained by

$$k^2 \Pi_E(k) - \Pi_G(k) < 0, \quad (14)$$


for all wavenumbers  $k$  outside of the forcing range.

- ➍ The transition wavenumber  $k_t$  where the break from  $k^{-3}$  to  $k^{-5/3}$  should occur, approaches the dissipation scale from the dissipation range, thus a transition cannot be seen visually in two-dimensional turbulence.
- ➎ Thus, the contribution of the downscale energy cascade to the energy spectrum is overwhelmed by the contribution of the downscale enstrophy cascade.


# The Nastrom-Gage spectrum. I

- ➊ A transition from  $k^{-3}$  scaling to  $k^{-5/3}$  scaling has been observed in the energy spectrum of large-scale atmospheric turbulence.
  - ➌ G.D. Nastrom and K.S. Gage (1984), *J. Atmos. Sci.* **42**, 950–960.
  - ➌ K.S. Gage and G.D. Nastrom (1986), *J. Atmos. Sci.* **43**, 729–740.
- ➋ This energy spectrum is known as the **Nastrom-Gage spectrum**.
- ➌ The QG model (quasi-geostrophic) describes large-scale atmospheric turbulence down to a scale of 100km.
- ➌ Charney claimed that the QG model is isomorphic to two-dimensional turbulence:
  - ➌ J.G. Charney (1971), *J. Atmos. Sci.* **28**, 1087–1095.
- ➌ This result by Charney has caused considerable confusion when efforts were made to explain the Nastrom-Gage spectrum in terms of 2D turbulence.

# The Nastrom-Gage spectrum. II



# Nastrom-Gage spectrum schematic



$k^{-3} \rightarrow 3000\text{km} - 800\text{km}$   
 $k^{-5/3} \rightarrow 600\text{km} - \ll 1\text{km}$   
 $k_t \approx 700\text{km} \approx k_R$

# Interpretation of NG spectrum. I

- The  $k^{-3}$  range is interpreted as downscale enstrophy cascade.
- The  $k^{-5/3}$  range used to be interpreted as an 2D inverse energy cascade forced at small scales by thunderstorms.
- This tortured interpretation followed from the perceived need to explain the Nastrom-Gage spectrum in terms of 2D turbulence.
- Tung and Orlando (formerly: Welch) challenged the Charney QG-2D equivalence:
  - K.K. Tung and W.T. Welch (2001), *J. Atmos. Sci.* **58**, 2009-2012.
  - K.K. Tung and W.W. Orlando (2003a), *J. Atmos. Sci.* **60**, 824-835.
  - K.K. Tung and W.W. Orlando (2003b), *DCDS B*, **3**, 145-162.
- New interpretation: A double downscale cascade of enstrophy and energy with enstrophy flux  $\eta_{uv}$  and energy flux  $\varepsilon_{uv}$  and transition from  $k^{-3}$  scaling to  $k^{-5/3}$  scaling at the transition wavenumber  $k_t \sim \sqrt{\eta_{uv}/\varepsilon_{uv}}$ .

# Interpretation of NG spectrum. II

- Recent measurements and data analysis by Cho and Lindborg (2001) have confirmed the existence of a *downscale* energy flux and estimate

$$\eta_{uv} \approx 2 \times 10^{-15} \text{ s}^{-3} \quad (15)$$

$$\varepsilon_{uv} \approx 6 \times 10^{-11} \text{ km}^2 \text{ s}^{-3} \quad (16)$$


- From these estimates we find the mean value of the transition scale

$$k_t = \sqrt{\eta_{uv}/\varepsilon_{uv}} \approx 0.57 \times 10^{-2} \text{ km}^{-1} \implies \lambda_t = 2\pi/k_t \approx 1 \times 10^3 \text{ km} \quad (17)$$

which has the correct order of magnitude.

- Tung and Orlando (2003) have also demonstrated numerically that a two-layer quasi-geostrophic channel model with thermal forcing, Ekman damping, and hyperdiffusion can reproduce the atmospheric energy spectrum.

# Tung and Orlando spectrum



# More on Tung-Orlando theory

- ➊ Smith-Tung debate: Transition not possible in 2D turbulence
  - ➌ K.S. Smith (2004), *J. Atmos. Sci.* **61**, 937-942
  - ➌ K.K. Tung (2004), *J. Atmos. Sci.* **61**, 943-948.
  - ➌ E. Gkioulekas and K.K. Tung (2007), *DCDS B* **7**, 293-314
- ➋ Smith argues: In 2D turbulence, transition from  $k^{-3}$  to  $k^{-5/3}$  cannot occur in the inertial range.
- ➌ Objections:
  - ➌ 2D turbulence is not QG turbulence.
  - ➌ In 2D turbulence we can rule out transition in  $S_3(r)$  but not in  $E(k)$ .
  - ➌ Very high-order hyperdiffusion can amplify ratio of Kolmogorov constants in 2D turbulence thereby allowing a transition in  $E(k)$ .
  - ➌ In QG turbulence, asymmetric dissipation can violate the constraint of vanishing downscale energy flux.

# The two-layer model. I

The governing equations for the two-layer quasi-geostrophic model are

$$\frac{\partial \zeta_1}{\partial t} + J(\psi_1, \zeta_1 + f) = -\frac{2f}{h}\omega + d_1 \quad (18)$$

$$\frac{\partial \zeta_2}{\partial t} + J(\psi_2, \zeta_2 + f) = +\frac{2f}{h}\omega + d_2 + 2e_2 \quad (19)$$

$$\frac{\partial T}{\partial t} + \frac{1}{2}[J(\psi_1, T) + J(\psi_2, T)] = -\frac{N^2}{f}\omega + Q_0 \quad (20)$$

where  $\zeta_1 = \nabla^2 \psi_1$ ;  $\zeta_2 = \nabla^2 \psi_2$ ;  $T = (2/h)(\psi_1 - \psi_2)$ .  $f$  is the Coriolis term;  $N$  the Brunt-Väisälä frequency;  $Q_0$  is the thermal forcing on the temperature equation;  $d_1$ ,  $d_2$ ,  $e_2$  the dissipation terms:

$$d_1 = (-1)^{\kappa+1} \nu \nabla^{2\kappa} \zeta_1 \quad (21)$$

$$d_2 = (-1)^{\kappa+1} \nu \nabla^{2\kappa} \zeta_2 \quad (22)$$

$$e_2 = -\nu_E \zeta_2 \quad (23)$$

# The two-layer model. II



The potential vorticity is defined as

$$q_1 = \nabla^2 \psi_1 + f + \frac{k_R^2}{2}(\psi_2 - \psi_1) \quad (24)$$

$$q_2 = \nabla^2 \psi_2 + f - \frac{k_R^2}{2}(\psi_2 - \psi_1) \quad (25)$$

with  $k_R \equiv 2\sqrt{2}f/(hN)$  and it satisfies

$$\frac{\partial q_1}{\partial t} + J(\psi_1, q_1) = f_1 + d_1 \quad (26)$$

$$\frac{\partial q_2}{\partial t} + J(\psi_2, q_2) = f_2 + d_2 + e_2 \quad (27)$$

with  $f_1 = -(k_R^2 Q)/(2f)$  and  $f_2 = (k_R^2 Q)/(2f)$  where  $Q = (1/4)k_R^2 h Q_0$ .

# The two-layer model. III

- The two-layer model conserves total energy  $E$  and potential layer enstrophies  $G_1$  and  $G_2$ :

$$E = \int [-\psi_1(x, y, t)q_1(x, y, t) - \psi_2(x, y, t)q_2(x, y, t)] dx dy \quad (28)$$

$$G_1 = \int q_1^2(x, y, t) dx dy \quad (29)$$

$$G_2 = \int q_2^2(x, y, t) dx dy \quad (30)$$

- We define:
  - The energy spectrum  $E(k) = \langle \psi_1, q_1 \rangle_k + \langle \psi_2, q_2 \rangle_k$
  - Top-layer potential enstrophy spectrum  $G_1(k) = \langle q_1, q_1 \rangle_k$
  - Bottom-layer potential enstrophy spectrum  $G_2(k) = \langle q_2, q_2 \rangle_k$
  - Total potential enstrophy spectrum  $G(k) = G_1(k) + G_2(k)$

# The two-layer model. IV



Consider the generalized form of an n-layer model:

$$\frac{\partial q_\alpha}{\partial t} + J(\psi_\alpha, q_\alpha) = \sum_\beta \mathcal{D}_{\alpha\beta} \psi_\beta + f_\alpha \quad (31)$$

$$\hat{q}_\alpha(\mathbf{k}, t) = \sum_\beta L_{\alpha\beta}(\|\mathbf{k}\|) \hat{\psi}_\beta(\mathbf{k}, t) \quad (32)$$



The energy spectrum  $E(k)$  and the potential enstrophy spectrum  $G(k)$  are given by:

$$E(k) = \sum_\alpha \langle \psi_\alpha, q_\alpha \rangle_k = \sum_{\alpha\beta} L_{\alpha\beta}(k) C_{\alpha\beta}(k) \quad (33)$$

$$G(k) = \sum_\alpha \langle q_\alpha, q_\alpha \rangle_k = \sum_{\alpha\beta\gamma} L_{\alpha\beta}(k) L_{\alpha\gamma}(k) C_{\beta\gamma}(k) \quad (34)$$

with  $C_{\alpha\beta}(k) = \langle \psi_\alpha, \psi_\beta \rangle_k$ .

# The two-layer model. V

- Let  $\phi_{\alpha\beta}(k) = \langle f_\alpha, \psi_\beta \rangle_k$ , and  $D_{\alpha\beta}(k)$  be the spectrum of the operator  $\mathcal{D}_{\alpha\beta}$
- The energy forcing spectrum  $F_E(k)$  and the potential enstrophy forcing spectrum  $F_G(k)$  are given by:

$$F_E(k) = 2 \sum_{\alpha} \phi_{\alpha\alpha}(k) \quad (35)$$

$$F_G(k) = 2 \sum_{\alpha\beta} L_{\alpha\beta}(k) \phi_{\alpha\beta}(k) \quad (36)$$

- The energy dissipation spectrum  $D_E(k)$  and the potential dissipation enstrophy spectrum  $D_G(k)$  are given by:

$$D_E(k) = 2 \sum_{\alpha\beta} D_{\alpha\beta}(k) C_{\alpha\beta}(k) \quad (37)$$

$$D_G(k) = 2 \sum_{\alpha\beta\gamma} L_{\alpha\beta}(k) D_{\alpha\gamma}(k) C_{\beta\gamma}(k) \quad (38)$$

# The energy flux constraint

- Can the 2-layer QG model violate the energy flux constraint  
 $k^2 \Pi_E(k) - \Pi_G(k) < 0$ ?
  - E. Gkioulekas and K.K. Tung (2007), *Discr. Contin. Dyn. Syst. Ser. B*, **7**, 293-314
- Recall that in the two-layer model, the dissipation terms read:

$$d_1 = \nu(-\Delta)^{\kappa+1} \psi_1, \quad (39)$$

$$d_2 = \nu(-\Delta)^{\kappa+1} \psi_2 - \nu_E \Delta \psi_2 \quad (40)$$

- Only the **asymmetric** presence of  $\nu_E \Delta \psi_2$  can break the energy flux constraint.
- Then the transition wavenumber  $k_t$  can occur in the inertial range.
- Necessary condition:

$$\nu_E > 4\nu k_{\max}^{2p} \left( \frac{k_{\max}}{k_R} \right)^2 \quad (41)$$

# Forcing spectrum. I.

- For antisymmetric forcing  $f_1 = f$  and  $f_2 = -f$ :

$$F_E(k) = 2(\Phi_1(k) - \Phi_2(k)) \quad (42)$$

$$F_G(k) = (k^2 + k_R^2)F_E(k) \quad (43)$$

with  $\Phi_1(k) = \langle f, \psi_1 \rangle_k$  and  $\Phi_2(k) = \langle f, \psi_2 \rangle_k$ .

- It follows that for  $k_f \ll k_R$ :  $(\eta/\varepsilon) \sim k_R^2$ .
- If both are dissipated at small scales, then  $k_t \sim k_R$ .
- In 2D turbulence Ekman damping dissipates most of injected energy and some of injected enstrophy.
- Not true in two-layer QG model because the Ekman term appears only on bottom layer.

# Forcing spectrum. II.

- For Ekman-damped forcing:  $f_1 = f$  and  $f_2 = -f - \nu_E \Delta \psi_2$ , incorporating the damping effect to the forcing spectrum gives:

$$F_E(k) = 2(\Phi_1(k) - \Phi_2(k)) + 2\nu_E k^2 U_2(k) \quad (44)$$

$$F_G(k) = (k^2 + k_R^2)F_E(k) - \nu_E k^2 k_R^2 (U_2(k) + C_{12}(k)) \quad (45)$$

with  $U_1(k) = \langle \psi_1, \psi_1 \rangle_k$ ,  $U_2(k) = \langle \psi_2, \psi_2 \rangle_k$ , and  $C_{12}(k) = \langle \psi_1, \psi_2 \rangle_k$ .

- Note the  $F_E(k)$  increases, because  $U_2(k) > 0$ .
- If  $E_K(k) \geq E_P(k) \implies C_{12}(k) \geq 0 \implies F_G(k)$  decreases.
- Thus, the tendency is to decrease  $k_t$ .
- Layer interaction makes it non-obvious whether  $F_G(k)$  increases or decreases.

# Forcing spectrum. IV.

- Claim: Suppressing bottom-layer forcing tends to decrease  $k_t$ .
- Let  $f_1 = f$  and  $f_2 = -\mu f$  with  $\mu \in (0, 1)$  (suppression factor).
- Assume  $f$  is random Gaussian with

$$\langle f_\alpha(\mathbf{x}_1, t_1) f_\beta(\mathbf{x}_2, t_2) \rangle = 2Q_{\alpha\beta}(\mathbf{x}_1, \mathbf{x}_2) \delta(t_1 - t_2) \quad (46)$$

- Define the forcing spectrum:

$$Q_{\alpha\beta}(k) = \frac{d}{dk} \int d\mathbf{x} d\mathbf{y} d\mathbf{z} P(k|\mathbf{x} - \mathbf{y}) P(k|\mathbf{x} - \mathbf{z}) Q_{\alpha\beta}(\mathbf{y}, \mathbf{z}) \quad (47)$$

- It follows that the streamfunction-forcing spectrum reads:

$$\varphi_{\alpha\beta}(k) = \langle f_\alpha, \psi_\beta \rangle_k = \sum_\gamma L_{\beta\gamma}^{-1}(k) Q_{\alpha\gamma}(k) \quad (48)$$

# Forcing spectrum. V.

- Energy forcing spectrum:

$$F_E(k) = 2 \sum_{\alpha} \varphi_{\alpha\alpha}(k) = \frac{2\mathcal{Q}(k)[2(1 + \mu^2)k^2 + (1 - \mu)^2 k_R^2]}{2k^2(k^2 + k_R^2)} \quad (49)$$

- Enstrophy forcing spectrum:

$$F_G(k) = 2 \sum_{\alpha} L_{\alpha\beta}(k) \varphi_{\alpha\beta}(k) = 2(1 + \mu^2)\mathcal{Q}(k) \quad (50)$$

- Consider the forcing range limit:  $k \ll k_R$ .
- For  $\mu = 1$ :  $F_G(k) \sim k_R^2 F_E(k) \implies k_t \sim k_R$
- Thus: Antisymmetric forcing indicates transition at  $k_R$ .
- For  $\mu = 0$ :  $F_G(k) \sim 2k^2 F_E(k) \implies k_t \sim 2k_f$
- Thus: Baroclinically damped forcing contributes to inertial range transition.

# Conclusion

- The energy flux constraint prevents a transition from  $k^{-3}$  scaling to  $k^{-5/3}$  scaling in 2D turbulence.
- The energy flux constraint can be broken in two-layer QG turbulence under asymmetric dissipation.
- In the two-layer QG model, the rates of enstrophy over energy injection satisfy:  
$$(\eta/\varepsilon) = k_R^2$$
- Asymmetric Ekman dissipation increases  $\varepsilon$  and may decrease  $\eta$ ,
- Suppressing the bottom layer forcing directly always decreases the ratio  $(\eta/\varepsilon)$ .
- Open question: Can the injected energy and enstrophy be dissipated?